# Difference Equations:
# *From Rabbits to Chaos*
# DRAFT

Paul Cull
Department of Computer Science
Oregon State University

Mary Flahive
Department of Mathematics
Oregon State University

Robby Robson
Department of Computer Science
Oregon State University

June 18, 2003

# Contents

**Preface** **ix**
Notational Preliminaries . . . . . xi

**1 Fibonacci Numbers** **1**
1.1 The Rabbit Problem . . . . . 1
1.2 The Fibonacci Sequence . . . . . 2
1.2.1 Computing Fibonacci numbers . . . . . 3
1.2.2 A Formula for the Fibonacci Numbers . . . . . 5
1.3 Notation for Asymptotic Analysis . . . . . 6
1.4 Exercises . . . . . 7

**2 Homogeneous Linear Recurrence Relations** **11**
2.1 The Solution Space of (HL) . . . . . 12
2.2 The Matrix Form . . . . . 15
2.3 A Simpler Basis for the Solution Space . . . . . 17
2.3.1 Distinct eigenvalues . . . . . 19
2.3.2 Repeated Eigenvalues . . . . . 21
2.4 The Asymptotic Behavior of Solutions . . . . . 25
2.5 Exercises . . . . . 28

**3 Finite Difference Equations** **33**
3.1 Linear Difference Equations . . . . . 33
3.1.1 First-Order Equations . . . . . 34
3.1.2 Finding a particular solution via convolution . . . . 36

3.2 A Special Class of Linear Recurrences . . . . . 38
3.3 The Shift Operator on the Space of Sequences . . . . . 40
3.4 Formal Power Series . . . . . 43
3.4.1 Formal differentiation . . . . . 48
3.4.2 An application of Formal Power Series . . . . . 49
3.5 Exercises . . . . . 51

**4 Generating Functions** **59**
4.1 Counting strings with some restrictions . . . . . 59
4.2 An Overview of the Generating Function Technique . . . . 62
4.3 A Review of Partial Fractions . . . . . 69
4.4 The Generating Function Technique . . . . . 75
4.4.1 The Catalan numbers . . . . . 75
4.4.2 Stirling Numbers of the Second Kind . . . . . 77
4.5 Reversion of Generating Functions . . . . . 79
4.6 Exercises . . . . . 83

**5 Nonnegative Difference Equations** **91**
5.1 Nonnegative Polynomials . . . . . 92
5.1.1 The Dominant Root . . . . . 92
5.2 Homogeneous Nonnegative Recurrences . . . . . 97
5.2.1 Roundability or When are integer solutions rounded powers of an eigenvalue? . . . . . 99
5.2.2 Using the Rounding Theorem . . . . . 103
5.3 Nonhomogeneous Nonnegative Equations . . . . . 105
5.4 Estimating Roots . . . . . 110
5.4.1 Estimation of the dominant root . . . . . 110
5.4.2 Estimation of the second root . . . . . 111
5.5 Calculation of the Roots . . . . . 114
5.5.1 The rate of convergence in Newton's method . . . . 119
5.6 Exercises . . . . . 123

**6 Leslie's Population Matrix Model** **127**
6.1 Leslie's Model . . . . . 127
6.1.1 How To Tell If A Leslie Matrix Is Primitive . . . . . 131
6.2 Leslie's Convergence Theorem . . . . . 132
6.3 Nonprimitive Leslie Matrices . . . . . 134
6.3.1 A Simple Example . . . . . 134
6.3.2 Special Case – Only one positive fertility rate . . . . 135
6.3.3 Asympotically Periodic Leslie Matrices . . . . . 135
6.4 Companion Matrices . . . . . 137
6.4.1 Matrices with Repeated Eigenvalues . . . . . 145
6.5 Nonnegative Companion Matrices . . . . . 148
6.5.1 Periodic Nonnegative Companion Matrices . . . . . 150
6.6 Back to Leslie Matrices . . . . . 154

6.6.1 Periodic Leslie Matrices . . . . . 155
6.6.2 Averaging . . . . . 159
6.7 The Limiting Effect of $L$ on Nonnegative Vectors . . . . . 160
6.7.1 The Period of Total Population . . . . . 163
6.8 Exercises . . . . . 164

**7 Matrix Difference Equations** **169**
7.1 Homogeneous Matrix Equations . . . . . 169
7.2 Nonnegative Matrix Equations . . . . . 175
7.2.1 Perron-Frobenius Theory . . . . . 175
7.2.2 Applications to Markov Chains . . . . . 176
7.2.3 Graphs and Matrices . . . . . 178
7.2.4 Comments on Nonprimitivity . . . . . 183
7.3 Algorithms for Primitivity . . . . . 188
7.4 Matrix Difference Equations with Input . . . . . 195
7.4.1 Reduction to One Dimension . . . . . 197
7.4.2 Reduction to Homogeneous Form . . . . . 201
7.5 Exercises . . . . . 202

**8 Modular Recurrences** **207**
8.1 Periodicity . . . . . 208
8.1.1 Periodicity of linear modular recurrences . . . . . 211
8.2 Fast Modular Computations . . . . . 214
8.3 Periodicity of Linear Recurrences in Finite Fields . . . . . 216
8.4 The Period of First-Order Linear Recurrences . . . . . 218
8.4.1 First-order recurrences with maximal period . . . . . 221
8.5 Periodic Second-Order Recurrences . . . . . 222
8.5.1 The period of the modular Fibonacci sequences . . . 224
8.6 Applications . . . . . 228
8.6.1 Application 1: Pseudorandom number generation . . 228
8.6.2 Application 2: Integer Factorization . . . . . 232
8.7 Exercises . . . . . 236

**9 Computational Complexity** **243**
9.1 Analysis of Algorithms . . . . . 244
9.1.1 Measuring Running Time . . . . . 244
9.1.2 The Towers of Hanoi: An Example . . . . . 246
9.2 Computer Arithmetic . . . . . 251
9.2.1 Addition and subtraction . . . . . 252
9.2.2 Multiplication and division . . . . . 252
9.3 An Introduction to Divide-and-Conquer . . . . . 253
9.3.1 Example: Polynomial multiplication . . . . . 254
9.4 Simple Divide-and-Conquer Algorithms . . . . . 258
9.4.1 Example 1: A return to polynomial multiplication . 260
9.4.2 Example 2: Matrix multiplication . . . . . 260

9.4.3 Example 3: MERGESORT . . . . . 262
9.4.4 Example 4: Applications of Newton's method . . . . 262
9.5 The Fast Fourier Transform . . . . . 264
9.5.1 The general form of the Fast Fourier Transform . . . 265
9.5.2 The FFT when $n = 2^k$ . . . . . 267
9.5.3 Fast evaluation and fast interpolation . . . . . 270
9.5.4 The fast polynomial multiplication algorithm . . . . 271
9.6 Average Case Analysis . . . . . 274
9.6.1 The LARGETWO algorithm . . . . . 274
9.6.2 The QUICKSORT algorithm . . . . . 276
9.7 Exercises . . . . . 279

**10 Some Nonlinear Recurrences** **287**
10.1 Some Examples . . . . . 287
10.2 Nonlinear Systems . . . . . 289
10.2.1 Sarkovskii's Theorem . . . . . 292
10.3 Chaos . . . . . 293
10.3.1 A Simple Chaotic System . . . . . 293
10.4 Local Stability . . . . . 297
10.4.1 Local Stability of a Fixed Point . . . . . 297
10.4.2 Local Stability of a Cycle . . . . . 298
10.4.3 Local Stability in Two Dimensions . . . . . 300
10.5 Global Stability . . . . . 303
10.6 Linear Fractional Recurrences . . . . . 305
10.6.1 Asymptotic Behavior . . . . . 307
10.6.2 Rational Coefficients and Periodicity . . . . . 308
10.6.3 Chaotic-like Behavior . . . . . 312
10.6.4 Invariant Distributions . . . . . 313
10.6.5 Showing Global Stability . . . . . 318
10.6.6 Summary . . . . . 319
10.7 Conclusion . . . . . 320
10.8 Exercises . . . . . 321

**References** **325**

**A Worked Examples** **337**
A.1 Every eigenvalue is simple . . . . . 337
A.2 A multiple root . . . . . 341

**B Complex Numbers** **345**

**C Highlights of Linear Algebra** **351**
C.1 Vector Spaces and Subspaces . . . . . 351
C.2 Linear Independence and Basis . . . . . 352
C.3 Linear Transformations . . . . . 353

C.4 Eigenvectors . . . . . 354
C.5 Characteristic and Minimal Polynomials . . . . . 356

**D Roots in the Unit Circle** **357**
D.1 Marden's method . . . . . 358
D.2 Exercises . . . . . 364

**Index** **365**

# Preface

Some years ago we noticed that various seemingly disparate fields were using similar models and techniques to solve similar problems. From mathematics to computer science to engineering to biology, various forms of difference equations were appearing in research papers and in textbooks, but with no common background the same results were being independently derived over and over again. As we were noticing this, some mathematics curricula were being revised with discrete mathematics replacing calculus as the first college mathematics course. New discrete mathematics courses were created and several superb textbooks appeared. In some fields a year of discrete mathematics actually replaced a year of calculus, while in other fields students took both discrete mathematics and calculus.

With these changes, what happened to difference equations? Some texts in discrete mathematics ignored them. Others had a few examples of difference equations as applications of proof by induction. Still others devoted a chapter to difference equations, but only solved a few special cases and/or represented generating functions (also called Z-transforms) as the principal or only method for finding solutions. With this lack of common background, texts on algorithms, signal processing, and population biology were still forced to devote chapters to the difference equations used in their areas. Even students who took several of these courses had difficulty seeing that they were working with the same difference equations in different contexts. Many instructors have written notes to flesh out the coverage given in texts, but such notes are of necessity usually so terse that students are led to believe that difference equations are very complicated and hard to understand.

With these problems in mind, we set out to write a book on difference equations which is accessible to undergraduates. As a text, it is meant for undergraduate majors in one of the mathematical sciences, presumably in their junior or senior year. It is not necessary that a student be a star mathematics student in order to use it, but the instructor should be aware that a student with *neither* good computational skills *nor* an understanding of the structure of mathematical statements and proofs will likely end up frustrated.

We wanted our text to have an algorithmic spirit. In this book, each chapter leads to techniques which can be applied by hand to small examples and also can be programmed for larger examples. In many cases we give explicit algorithms. We decided to write them in pseudo-code rather than in a specific programming language for several reasons. First, it is easy to translate from pseudo-code into any reasonable programming language. Second, there are many programming languages available and translating from one language to another is often more difficult than translating from pseudo-code. Third, we are not sure that programming these algorithms is worth the effort because for almost all of our examples there are high-quality implementations readily available on the web. It probably makes more sense to use one of these codes rather than to cobble together a program that will only be used a few times and/or will be prone to problems when the data is not exactly in the form assumed by the programmer. A number of mathematically oriented computer packages are also available. For example, MATLAB, Maple and Mathematica all have packages which will solve difference equations and recurrence relations. In many cases these packages give numeric answers as well as symbolic solutions when possible. Using these packages is much simpler than programming from scratch.

In this book we start with the old story of Fibonacci's rabbits and progress through several generalizations, ending with some nonlinear difference equations. We deal with familar mathematical structures such as the real numbers, the complex numbers, the integers, or the integers modulo an integer. We were tempted to discuss more general structures in order to show, for example, how theories of computation could be represented as difference equations, but we soon discovered this would result in either a very large book or a very formal book which would be at variance with our goal of accessibility. After developing the theory and techniques for solving linear difference equations in Chapters 2 to 4, we specialize to equations with nonnegative coefficients in Chapters 5 and 6 and then consider a generalization to matrix difference equations in Chapter 7. Chapter 8 considers equations over other rings, including integers modulo m and finite fields. Chapter 9 considers some issues in computational complexity, including divide-and-conquer algorithms. We end with some nonlinear systems in Chapter 10. Along the way we use linear algebra, develop formal power series, solve some combinatorial problems, visit Perron-Froebinus theory,

use graph theory, discuss pseudorandom number generation and integer factorization, and use the FFT to multiply polynomials quickly.

There are four appendices serving different purposes. The first is a collection of worked examples, meant to supplement the early chapters of the book. Because the material in Appendices B and C is indispensible to an understanding of the book, we suggest working through them before beginning Chapter 2. Although many of the difference equations we consider have integer or real coefficients, it is often necessary to consider the coefficients as complex numbers. Appendix B gives the highlights of the complex analysis we need and no prior experience is necessary to understand this appendix. On the other hand, only the most exceptional student could learn new material at the rate with which linear algebra is presented in Appendix C. One of the aims of this book is to show students that linear algebra is a powerful and coherent subject whose ideas have diverse applications, and we hope Appendix C is a helpful review. Appendix D outlines a method of Morris Marden [92] which can be used to decide when the general solution of a difference equation converges to zero. This appendix is not needed for an understanding of the book.

## Notational Preliminaries

In this book, we use the following fairly standard notation:

$\mathbb{Z}$ is the set of **integers**.

$\mathbb{Z}_m$ is the set of **integers modulo m** .

$\mathbb{Z}^k$ is the set of all **k-tuples** with integer coordinates.

$\mathbb{Z}_m^k$ is the set of all **k-tuples** of integers modulo m.

$\mathbb{N}$ is the set of **natural numbers**, including 0; $\mathbb{N} = \{0, 1, 2, \dots\}$.

$\mathbb{N}^+$ is the set of **positive integers**.

$\mathbb{Q}$ is the set of **rational numbers**.

$\mathbb{R}$ is the set of **real numbers**.

$\mathbb{F}$ denotes a **finite field**.

$\mathbb{C}$ is the set of **complex numbers**.

$\mathbb{R}[x]$ is the set of **polynomials** with real coefficients.

$\mathbb{C}[x]$ is the set of **polynomials** with complex coefficients.

$\mathbb{Z}_m[x]$ is the set of **polynomials** whose coefficients are integers modulo m.

$\mathbb{F}[x]$ is the set of **polynomials** with coefficients from the finite field $\mathbb{F}$.

$\lfloor x \rfloor$ is the **floor** of $x \in \mathbb{R}$; the largest integer $n$ with $n \leq x$.

$\lceil x \rceil$ is the **ceiling** of $x \in \mathbb{R}$; the smallest integer $n$ with $n \geq x$.

# 1
# Fibonacci Numbers

This chapter is devoted to the Fibonacci numbers. We start with the familiar definition, move to some more sophisticated views, and then formulate some questions which are typical of those which can be addressed using the material of this book.

## 1.1 The Rabbit Problem

In the year 1202 the Italian mathematician Leonardo Pisano (which means Leonardo of Pisa) published the book *Liber Abbaci*,[1] a book of problems whose purpose was to illustrate the usefulness of Arabic numerals in arithmetic computations because at that time cumbersome Roman numerals were still being used in Italy. One of the problems discussed in Pisano's book considers pairs of breeding rabbits. Each pair of rabbits matures in two months. It produces one new pair each month thereafter, beginning with the last day of its second month. Starting with a single infant pair born at the beginning of Month 0, how many pairs will there be at the end of one year? We can can find our way to a solution by considering what happens in the first few months:

1. At the end of Month 0 there is only one pair and they are not yet breeding.

[1]The book was reprinted in 1857-1862 by Baldassarre Boncompagni [12]. A translation by L. Sigler [128] has recently been published by Springer-Verlag.

2. At the end of the Month 1 the first pair gives birth to a second pair.

3. At the end of Month 2 the second pair is one month old and not yet breeding. The first pair produces another pair, giving three pairs at the end of Month 2.

4. By the end of Month 3, the first pair has produced yet another pair and the second pair has given birth to its first progeny. Therefore there are five pairs at the end of Month 3.

5. For Month 4, the two new pairs born the previous month are not bearing young, but all three older pairs give birth to one new pair. Eight pairs are alive at the end of Month 4.

From this we observe that at the end of the $n^{\text{th}}$ month each pair that was alive at the $(n-2)^{\text{nd}}$ month has given birth to a new pair during the month. Therefore the total number of pairs alive at the end of the $n^{\text{th}}$ month is the number from the $(n-1)^{\text{st}}$ month *plus* one new pair for every pair which was alive at the end of the $(n-2)^{\text{nd}}$ month. In other words, the number of rabbit pairs at the end of $n$ months is the sum of the numbers at the end of the two previous months.

This most famous example of a recurrence relation is called the **Fibonacci numbers**, because Pisano is usually referred to as Fibonacci which means son of Bonaccio. The Fibonacci numbers are given by the sequence

$$1, 2, 3, 5, 8, \ldots$$

whose first two terms are $1, 2$ and each subsequent term is the sum of the preceding two terms. The Rabbit Problem asks for the thirteenth element of this sequence, which is the number 377. If we are asked for the number of rabbits after one thousand months, could we compute the $1001^{\text{st}}$ Fibonacci number without computing all 1000 numbers which came before? Examination of some slightly more sophisticated ways to view the Fibonacci numbers will provide an answer to this question.

## 1.2 The Fibonacci Sequence

In calculus we often think of a function such as $f(x) = x^2$ as a simpler object than an infinite sequence but a **sequence** $s_0, s_1, s_2, \ldots$ of complex numbers is really just a function

$$s : \mathbb{N} \to \mathbb{C},$$

where $s_n$ is the value of the function $s$ at the point $n$. In this book we will denote elements of sequences using subscripts and use $< s_n >$ as shorthand

for the entire sequence. We will also define a sequence by giving its "generic term," so $< n^2 >$ is the sequence $< s_n >$ with $s_n = n^2$ for all $n \in \mathbb{N}$.

From now on, we will slightly modify the Fibonacci sequence by defining the first two Fibonacci numbers to be 0 and 1 while each subsequent Fibonacci number is still the sum of its two immediate predecessors. Such a rule which defines the elements of a sequence by a formula involving some fixed number (in this case two) of preceding elements is called a **recurrence relation**. The Fibonacci sequence is defined by the linear recurrence

$$f_0 = 0, \quad f_1 = 1 \tag{1.1}$$

$$f_{n+2} = f_{n+1} + f_n \,. \tag{1.2}$$

The recurrence relation itself is (1.2) and the two values given in (1.1) are called the **initial conditions**. The system (1.2) and (1.1) is referred to as an **initial value problem**. The equation in (1.2) is called **linear** because the terms of the sequence are connected in a linear manner, using only linear combinations of previous elements of the sequence. (In general, scaling by real or complex constants is allowed.)

As we said above, it's often helpful to think of a sequence as a function $f$ defined on the set of natural numbers $\mathbb{N}$. The Fibonacci sequence is defined by $f(n) = f_n$ with

$$f(0) = 0, f(1) = 1 \tag{1.3}$$

$$f(n+2) = f(n+1) + f(n). \tag{1.4}$$

Here we begin the sequence at the position $f(0)$ because verifying and developing formulas is often easier when initializing with $f(0)$ rather than with $f(1)$.

When we define the Fibonacci sequence it's customary to think of the index $n$ in (1.2) and (1.4) as a natural number but there's no purely mathematical reason to limit ourselves to nonnegative $n$. From any two consecutive Fibonacci numbers the relation (1.2) allows us to compute the preceding element of the Fibonacci sequence using $f_n = f_{n+2} - f_{n+1}$. Therefore, we can proceed backwards as well as forwards, and any integer can be used as an index. Some "center" terms of the associated doubly infinite sequence are

$$\cdots, -8, 5, -3, 2, -1, 1, 0, 1, 1, 2, 3, 5, 8, \cdots .$$

(Exercise 1.4 asks you to verify this pattern.) Extending recurrences to include negative indices can have some advantages, sometimes giving a deeper insight into the theory. It's especially helpful for modeling population dynamics and other time-dependent phenomena.

### *1.2.1 Computing Fibonacci numbers*

The definition of the Fibonacci function given in (1.4) is called a **recursive definition**. It allows us to write a short recursive computer program to

compute (at least in theory) the value of the Fibonacci function at any given nonnegative $n$.

```
PROCEDURE  RECRFIB(n,f) (* Input n, Output f(n) *)
           IF n ≤ 1
              THEN  f := m
              ELSE RECRFIB(n − 1, g)
                   RECRFIB(n − 2, h)
                   f := g + h
```

Another style of program for computing $f(n)$ for nonnegative $n$ is given by

```
PROCEDURE FIB(n,f) (*Input n, Output f(n)*)
          Declare Integer Variables: t, j := 0
          IF n ≤ 1 THEN f := n
                   ELSE f := 1
                        FOR i:=2 TO n
                            t := f
                            f := f + j
                            j := t
                        ENDFOR
```

The first procedure, RECRFIB, computes by making calls to itself. Such a program is usually called **recursive**. Program FIB does not call on itself, but computes by repeatedly executing the statements in the FOR loop. Such a program is usually called **iterative**.

Once we know *some* program which computes a given function, it is natural to ask for a program which computes the function rapidly. This question is not as innocent as it sounds. Returning to the Fibonacci function, we might want to know how quickly we can produce a *single term* $f(n)$ in terms of the size of the input $n$, or we might want to know how quickly we can compute *all* of the first $n$ terms of the Fibonacci sequence. Or, we might want to answer these same questions in the context of constraints on the size of the memory of the computer used for the computation. These may be realistic or simply theoretical constraints. More effective procedures for computing Fibonacci numbers are given in [34].

### *1.2.2 A Formula for the Fibonacci Numbers*

Although the defining recurrence (1.2) *theoretically* allows us to compute $f(n)$ for any $n$, it's not an especially tidy mathematical expression for $f(n)$ since all previous terms are needed in order to calculate just one term of the Fibonacci sequence. For example, think about the question of quickly estimating the number of rabbit pairs after one thousand months of rabbit breeding. If the function $f$ had a nice formula, it would be easier to answer such a question. Fortunately, and perhaps surprisingly, there is such a formula for values of the Fibonacci function:

$$\boxed{f(n) = \frac{1}{\sqrt{5}}\left[\left(\frac{1+\sqrt{5}}{2}\right)^n - \left(\frac{1-\sqrt{5}}{2}\right)^n\right].} \tag{1.5}$$

You might recognize the number $\dfrac{1+\sqrt{5}}{2}$; it's often referred to as the **golden mean** or **golden section**. This formula (1.5) is known as **Binet's Formula** and can be derived using the techniques given in the next chapter (refer to Exercise 2.10). Although it is usually attributed to Jacques Phillipe Marie Binet (1786-1856), Donald Knuth [77, vol. 1, p. 82] says that Abraham de Moivre reported it in 1730 [40, pgs. 26-42] where he considered the general linear recurrence.

For now, we can appreciate the power of Binet's Formula by using it to estimate size of the $1000^{\text{th}}$ Fibonacci number. Since

$$\frac{1+\sqrt{5}}{2} \approx 1.618 \text{ and } \frac{1-\sqrt{5}}{2} \approx -.618\,.$$

and $(.618)^{1000}$ is very small, then

$$f(1000) \approx \frac{1}{\sqrt{5}}(1.618)^{1000}.$$

If we're interested in estimating the size of $f(1000)$ in terms of the number of its decimal digits we can use the base 10 logarithm. (Refer to Exercise 1.7.) Since

$$1000 \cdot \log(1.618) - .5\log(5) \approx 208.629,$$

we expect the number of decimal digits in $f(1000)$ to be about 209. In fact, the $1000^{\text{th}}$ Fibonacci number is the number

$$\begin{aligned} f(1000) = & 43466557686937456435688527675040625802564660517371 \\ & 78040248172908953655541794905189040387984007925516 \\ & 92959225930803226347752096896232398733224711616429 \\ & 96440906533187938298969649928516003704476137795166 \\ & 849228875, \end{aligned}$$

which does have the predicted 209 digits.

Letting $\lambda_0 = \frac{1+\sqrt{5}}{2}$, $\lambda_1 = \frac{1-\sqrt{5}}{2}$ and $\alpha = 1/\sqrt{5}$, Binet's Formula can be rewritten as

$$f(n) = \alpha(\lambda_0^n - \lambda_1^n)\,.$$

Since $|\lambda_1| < 1$ and $\alpha < 1/2$, then $|f_n - \alpha\lambda_0^n| = \alpha|\lambda_1|^n < 1/2$ and the fact that $f_n$ is an integer therefore gives the pleasant identity

$$f_n = Round\Big(\lambda_0^n/\sqrt{5}\Big) \text{ for all } n \geq 0, \tag{1.6}$$

where $Round(X)$ returns the integer nearest to $X$.[2] We analyze this notion of **roundability** in Chapter 5 (refer to Theorem 5.2.4).

## 1.3 Notation for Asymptotic Analysis

There are often many algorithms for the same problem. For instance, in the case of computing Fibonacci numbers we've already written two types of algorithms for computing a specific Fibonacci number, either using the definition directly or using Binet's Formula. Our usual method of assessing the efficiency of various algorithms for a problem will be to compare their running times, and usually this is done with an asymptotic analysis. For this there are three standard forms of notation, **Big-Oh**, **Big-Omega**, and **Big-Theta**. Each notation removes unimportant details so we can see the size of running time more clearly. This notation was codified for computer scientists by Donald Knuth [74, 75] in 1976 when he drew upon notation already used in analytic number theory. Big-Oh[3] was introduced by Bachmann in 1894, and something very similar to Big-Omega was used by Hardy and Littlewood in 1914. More information on the history of this notation (and that of Big-Theta from the 1960s) can be found in Knuth's article. Although our principal use of this notation is for positive real numbers, our definitions allow $T(n)$ to be a *complex-valued* sequence.

---

[2] Note that $Round(\frac{k}{2})$ is not defined when $k$ is an odd integer.

[3] The term Big-Omicron is used by Knuth but most other authors call this Big-Oh notation.

Let $T(n)$ and $f(n)$ be two complex-valued sequences. Then

$T(n) = O(f(n))$ (say $T(n)$ has order at most $f(n)$) means there exists a positive constant $c$ such that

$$|T(n)| \le c|f(n)| \quad \text{for all sufficiently large } n\,.$$

$T(n) = \Omega(f(n))$ (say $T(n)$ has order at least $f(n)$) means there exists a positive constant $c$ such that

$$|T(n)| \ge c|f(n)| \quad \text{for all sufficiently large } n\,.$$

$T(n) = \Theta(f(n))$ (say $T(n)$ has order exactly $f(n)$) means there exists positive constants $c_1, c_2$ such that

$$c_1|f(n)| \le |T(n)| \le c_2|f(n)| \quad \text{for all sufficiently large } n\,.$$

For instance, from (1.6) we know the asymptotic size of $f_n$ is $\Theta(\lambda_0^n)$ where $\lambda_0 = (1+\sqrt{5})/2$. To get some practice with this notation, you should verify that for any fixed pair of positive integers $i \le j$ each of the following holds

$$n^i = O(n^j) \quad , \quad n^j = \Omega(n^i) \quad , \quad \Theta(n^i) = \Theta(n^j) \text{ implies } i = j \quad .$$

## 1.4 Exercises

**Ex 1.1.** (Taken from *Liber Abbaci*, pp. 283 ff.) Translate the following:

‘‘Quot paria coniculorum in uno anno ex uno pario germinentur.’’
Qvidam posuit unum par cuniculorum in quodam loco, qui erat undique pariete circundatus, ut sciret, quot ex eo paria germinarentur in uno anno: cum natura eorum sit per singulum mensem aliud par germinare; et in secundo mense ab eorum natiuitate germinant. Quia suprascriptum par in primo mense germinat, duplicabis ipsum, erunt paria duo in uno mense.
Ex quibus unum, scilicet primum, in secundo mense germinat; et sic sunt in secundo mense paria 3; ex quibus in uno mense duo pregnantur; et germinatur in tercio mense paria 2 conciculorum; et sic sunt paria 5 in ipso mense;... Cum quibus etiam additis parijs 144, que germinatur in ultimo mense, erunt paria 377; et tot paria peperit suprascriptum par in prefato loco in capite unius anni. ...

**Ex 1.2.** Let $x \in \mathbb{R}$. Show that $\lfloor x \rfloor = \lceil x \rceil$ if and only if $x \in \mathbb{Z}$.

**Ex 1.3.** Suppose $< s_n >$ is any sequence which satisfies the Fibonacci recurrence (1.2) but possibly has different integer initial values. Let $n_1, n_2 \in \mathbb{N}$ be any fixed indices. Show that any term $s_n$ can be written as a linear combination of $s_{n_1}, s_{n_2}$ with *integer* coefficients.

**Ex 1.4.** Use (1.2) to show that for the Fibonacci function we have

$$f(-n) = (-1)^{n+1} f(n) \text{ for all } n \geq 0.$$

**Ex 1.5.** Verify Binet's Formula (1.5) for $f(2), f(3), f(4)$.

**Ex 1.6.** Use mathematical induction to prove that every element of the Fibonacci sequence satisfies Binet's Formula.

**Ex 1.7.** Show that the base 10 logarithm of each of the integers $1, 2, \ldots, 9$ lies in the interval $[0, 1)$, and that the logarithm of any two-digit integer lies in the interval $[1, 2)$. In general, show that the number of decimal digits of an integer $n$ is the ceiling of its logarithm to the base 10.

**Ex 1.8.** If the base 10 logarithm of an integer begins with 3.53, how many digits does the integer have and what's its first digit?

**Ex 1.9 (Lucas Numbers).** [4] Consider the sequence $< L_n >$ generated by

$$L_0 = 2, \quad L_1 = 1$$
$$L_{n+2} = L_{n+1} + L_n \, .$$

(a) Calculate $L_2, L_3, L_4$.
(b) Find constants $A, B \in \mathbb{R}$ such that for each of $n = 0, 1$ it is true that

$$L_n = A \left( \frac{1+\sqrt{5}}{2} \right)^n - B \left( \frac{1-\sqrt{5}}{2} \right)^n . \tag{1.7}$$

(c) Use mathematical induction to prove that your formula in the previous part holds for all $n \geq 0$.

**Ex 1.10.** For this exercise, write a short program which computes the Fibonacci and the Lucas sequences.
(a) Use your program to calculate $f_{30}$ and $L_{30}$. Check your answer using the closed forms in (1.5) and (1.7). Compute the ratio $\dfrac{f_{30}}{L_{30}}$.
(b) The ratio $\dfrac{f_n}{L_n}$ has a limiting value. Use your program to calculate this value to ten decimal places.

[4] The French mathematician Édouard Lucas (1842-1891) studied the properties of this and other sequences in the first volume of the *American Journal of Mathematics* (specifically page 186) which appeared in 1878.

**Ex 1.11.** For this exercise, consider the sequence $< s_n >$ defined by

$$s_0 = 0, \quad s_1 = 1$$
$$s_{n+2} = 2s_{n+1} + 2s_n \, .$$

(a) Calculate the first five terms of the sequence $< s_n >$.
(b) Check that the following formula correctly calculates $s_2, s_3, s_4$:

$$s_n = \frac{1}{\sqrt{12}} \left[ (1+\sqrt{3})^n - (1-\sqrt{3})^n \right] .$$

(c) Show that the general $n^{\text{th}}$ term of the sequence $< s_n >$ satisfies the formula in the previous part.

**Ex 1.12.** Show that the variable $t$ in the procedure FIB can be eliminated by using $f := f + j$, $j := f - j$.

**Ex 1.13.** Show that every natural number $n$ can be expressed in the "binary" form $n = \sum_{i=2}^{K} b_i f_i$ where each $b_i$ is either 0 or 1 and $K$ depends on the natural number $n$. Show further that this binary Fibonacci representation is NOT unique. When the additional stipulation that no two consecutive $b_i$ are both 1 is imposed, show that the binary Fibonacci representation is unique.

**Ex 1.14.** Define $\log_{FIB}(n)$ to equal the least index $i$ for which $f_i \geq n$.
(a) Calculate some values of $\log_{FIB}(n)$.
(b) For each $i \geq 1$, define $\#(i)$ to be the number of natural numbers $n$ for which $\log_{FIB}(n) = i$. Show that $\#(i) = f_i$ for all $i \geq 3$.

**Ex 1.15.** Show that any solution to

$$s_n = s_{n-1} + \lfloor \sqrt{n} \rfloor$$

with nonnegative initial conditions satisfies $s_n = \Theta(n^{3/2})$.

# 2

# Homogeneous Linear Recurrence Relations

The simplest type of linear recurrence relation is the homogeneous linear recurrence with constant coefficients, one in which $s_{n+k}$ is given as a *linear function* of $s_n, \ldots, s_{n+k-1}$. In other words, for all $n \geq 0$

$$\text{(HL)} \qquad s_{n+k} = c_1 s_{n+k-1} + c_2 s_{n+k-2} + \cdots + c_k s_n \,,$$

where $c_1, \ldots, c_k$ are complex constants and $c_k \neq 0$. This is called a **$\mathbf{k^{th}}$-order homogeneous linear recurrence relation** with constant coefficients. The purpose of this chapter is to analyze these recurrence relations using tools from linear algebra. Appendix C contains a review of the basic linear algebra which we assume here. Even if we were only interested in *integer* recurrence relations, we would still need to consider recurrences whose coefficients are real and complex numbers because this is more or less forced on us by the linear algebra. In addition, more compact formulas can often be obtained by using more general number systems.

For any set of $k$ initial values, the recurrence relation (HL) can be applied to compute the infinite sequence $< s_n >$ which satisfies this recurrence: From $s_0, \ldots, s_{k-1}$ the $s_k$ term is specified by $s_k = c_1 s_{k-1} + \cdots + c_k s_0$; the $s_{k+1}$ term by $s_{k+1} = c_1 s_k + \cdots + c_k s_1$; and so on. Equation (HL) with given values $s_0, \cdots, s_{k-1}$ is called an **initial value problem**. What we've just informally shown is that every such initial value problem has a solution and that the solution is unique. The first $k$ terms determine the rest of the sequence. In Exercise 2.1 we ask you to supply a formal verification of this fact, which is fundamental for the remainder of this chapter.

**Arithmetic Operations on Sequences**

We can add and scale sequences in exactly the same way as functions. That means $< s_n > + < t_n > = < s_n + t_n >$. Viewed as functions, this is the rule $(s+t)(n) = s(n) + t(n)$. A sequence is multiplied by scalar $\lambda \in \mathbb{C}$ using the rule $\lambda(s_n) = (\lambda s_n)$. The set of all functions from $\mathbb{N}$ into $\mathbb{C}$ (which is the same as the set of all complex sequences) is a complex vector space under these operations of addition and scalar multiplication.

## 2.1 The Solution Space of (HL)

We begin by analyzing the second-order initial value problem

$$s_{n+2} = 3s_{n+1} - 2s_n\,, \quad s_0 = 2\,,\ s_1 = 3, \tag{2.1}$$

which generates the sequence

$$2, 3, 5, 9, 17, 33, 65, 129, 257, 513, \ldots$$

and has the closed formula $s_n = 2^n + 1$. We might ask what the sequence looks like under other initial conditions. Here are two such sequences:

$$2, 5, 11, 23, 47, 95, 191, 383, 767, 1535, \ldots$$
$$3, 1, -3, -11, -27, -59, -123, -251, -507, -1019, \ldots$$

It's not quite as easy to guess a formula for the $n^{\text{th}}$ term of the sequence starting with $2, 5$. However, you might notice that

$$23 = 24 - 1\,,$$
$$47 = 48 - 1\,,$$
$$95 = 96 - 1\,,$$

and arrive at the formula $s_n = 3 \cdot 2^n - 1$. By thinking negatively, we can find the formula $s_n = 5 - 2^{n+1}$ for the third set of initial conditions $s_0 = 3, s_1 = 1$. (In Exercise 2.2 you're asked to prove these formulas by an inductive argument.) Since we used the same recurrence (2.1) to generate all three sequences, it's not surprising that there are similarities among the formulas. For instance, all three formulas involve powers of 2. One of our goals is to understand how and why the sequences corresponding to different initial values are similar. To put this another way, we're interested in understanding the structure of the space of all solutions to a fixed $k^{\text{th}}$-order homogeneous linear recurrence relation when the initial conditions range over all $k$–tuples of complex numbers.

Placing this in a more general setting, we've observed already that the set of all complex sequences forms a vector space over $\mathbb{C}$. In Exercise 2.5 you're asked to prove that the set of solutions to (HL) is a *subspace* of the

space of all complex sequences which means that every linear combination of solutions is also a solution. We'll call this the **solution space of (HL)** and will denote it by $\mathcal{X}$.

We've seen that any sequence in $\mathcal{X}$ is completely determined by its first $k$ terms. This can be neatly expressed by defining a map $\pi$ which picks out these terms and writes them as a vector in $\mathbb{C}^k$. Accordingly, we define

$$\pi : \mathcal{X} \to \mathbb{C}^k \text{ by } \pi(< s_n >) = (s_{k-1}, \ldots, s_1, s_0)^T. \tag{2.2}$$

Because every sequence in $\mathcal{X}$ has a unique string of $k$ initial elements, this map is a well-defined function. The fact that any choice of values $s_0, \ldots, s_{k-1}$ can be extended to a infinite sequence which is in $\mathcal{X}$ means that $\pi$ is an **onto function**. The companion fact that every initial value problem has a *unique* solution translates to the statement that $\pi$ is a **one-to-one function** (that is, if $\pi(x) = \pi(y)$ for $x, y \in \mathcal{X}$ then $x = y$). Since $\pi$ is one-to-one and onto, we call it a **bijection** or a **bijective function** and know that $\pi$ has a (two-sided) inverse. This inverse is the map which assigns to any vector $\boldsymbol{\alpha} = (\alpha_{k-1}, \ldots, \alpha_0)^T \in \mathbb{C}^k$ the unique solution in $\mathcal{X}$ which begins with the initial values $s_0 = \alpha_0$, $s_1 = \alpha_1$, ...,$s_{k-1} = \alpha_{k-1}$.

For any two solutions $< x_n >, < y_n > \in \mathcal{X}$ and complex scalars $\alpha$ and $\beta$, we have[1]

$$\begin{aligned}
\alpha * \pi(< x_n >) + \beta * \pi(< y_n >) &= \alpha * \begin{pmatrix} x_{k-1} \\ \vdots \\ x_1 \\ x_0 \end{pmatrix} + \beta * \begin{pmatrix} y_{k-1} \\ \vdots \\ y_1 \\ y_0 \end{pmatrix} \\
&= \begin{pmatrix} \alpha * x_{k-1} + \beta * y_{k-1} \\ \vdots \\ \alpha * x_1 + \beta * y_1 \\ \alpha * x_0 + \beta * y_0 \end{pmatrix} \\
&= \pi(\alpha * < x_n >) + \beta * < y_n >),
\end{aligned}$$

which shows that $\pi : \mathcal{X} \to \mathbb{C}^k$ is a linear transformation. Since we've already shown it is a bijection, $\pi$ is an invertible transformation which is often called a **vector space isomorphism**. From linear algebra we know that the inverse of an invertible linear transformation is also linear and we obtain the following

[1] Good notation often leads to seductively simple formulas which can obscure some important details, so it is good to form the habit of making a mental note of mulitiple uses of the same symbol. In this regard, observe that the symbol "+" has three different meanings in this equation: first as addition of vectors in $\mathbb{C}^k$, second as addition of complex numbers, and finally as addition of sequences. Check that you can also find the three different uses of the symbol "$*$".

**Theorem 2.1.1.** *Let $\mathcal{X}$ be the solution space of a $k^{th}$-order homogeneous linear recurrence relation. Then the map $\pi$ defined in (2.2) is an isomorphism between the vector spaces $\mathcal{X}$ and $\mathbb{C}^k$. In particular, $\mathcal{X}$ is a $k$–dimensional complex vector space.*

Since any element of a vector space is a (unique) linear combination of basis vectors, Theorem 2.1.1 can be interpreted as giving a method for constructing solutions $< s_n > \in \mathcal{X}$ using the $k$ sequences generated by the basic initial conditions. Let $\mathbf{e}_i$ be the $i^{\text{th}}$ standard basis vector in $\mathbb{C}^k$, the column vector with 1 in the $i^{\text{th}}$ position and zeros elsewhere. Then $\pi^{-1}(\mathbf{e}_i) =< s_n >$ is the solution whose only non-zero initial condition is $s_{k-i} = 1$. Since the set $\{\mathbf{e}_1, \ldots, \mathbf{e}_k\}$ is a basis for $\mathbb{C}^k$ and $\pi^{-1}$ is an isomorphism, the set $\{\pi^{-1}(\mathbf{e}_1), \ldots, \pi^{-1}(\mathbf{e}_k)\}$ is a basis for $\mathcal{X}$. Moreover, for given initial conditions

$$s_0 = \alpha_0\,, s_1 = \alpha_1\,, \ldots, s_{k-1} = \alpha_{k-1}\,,$$

the corresponding solution sequence is

$$\alpha_0\pi^{-1}(\mathbf{e}_k) + \alpha_1\pi^{-1}(\mathbf{e}_2) + \cdots + \alpha_{k-1}\pi^{-1}(\mathbf{e}_1)\,.$$

We illustrate this with two examples, the Fibonacci sequence and the recurrence in (2.1) above.

For the Fibonacci recurrence, the basic sequences are the two sequences

$$\pi^{-1}(\mathbf{e}_1) = (0, 1, 1, 2, 3, 5, 8, 13, 21, \ldots)$$

and

$$\pi^{-1}(\mathbf{e}_2) = (1, 0, 1, 1, 2, 3, 5, 8, 13, 21, \ldots)\,.$$

If we want to find the sequence generated by the recurrence

$$s_{n+2} = s_{n+1} + s_n\,, \quad s_0 = -1\,, s_1 = 3\,,$$

we would add 3 times $\pi^{-1}(\mathbf{e}_1)$ to $-1$ times the sequence $\pi^{-1}(\mathbf{e}_2)$, and this gives

The Fibonacci numbers enjoy many special properties which are not necesarily shared by all recurrences. For instance, the sequence $\pi^{-1}(\mathbf{e}_1)$ is simply the shift of the sequence $\pi^{-1}(\mathbf{e}_2)$ one term to the left. It follows that any sequence $< s_n >$ satisfying the Fibonacci recurrence has general term

$$s_n = af_n + bf_{n-1}\,,$$

for $a = s_1\,, \quad b = s_0\,.$

$-1, 3, 2, 5, 7, 12$

For recurrence (2.1), the basic sequences are

$$\pi^{-1}(\mathbf{e}_1) = (0, 1, 3, 7, 15, 31, 63, 127, 255, 511, \ldots)$$

and

$$\pi^{-1}(\mathbf{e}_2) = (1, 0, -2, -6, -14, -30, -62, -126, -254, -510, \ldots)\,.$$

Therefore, the ninth term of the solution to (2.1) with $s_0 = 4$, $s_1 = 11$ is

$$s_8 = 4(-254) + 11(255) = 1789\,.$$

To summarize: Knowledge of the basic sequences of a homogeneous linear recurrence allows us to express the $n^{\text{th}}$ term of any solution as a linear combination of simpler quantities, namely, the $n^{\text{th}}$ terms of the basic sequences. Because this procedure uses $k$ basic sequences to find the one sequence of interest, it isn't very useful unless formulas for the basic sequences can be easily obtained. We'd like a basis for $\mathcal{X}$ consisting of sequences which are guaranteed to have an easy-to-find and simple formula for its $n^{\text{th}}$ term. Obtaining such a basis is the goal of the next sections of this chapter.

## 2.2 The Matrix Form

Since each term of a $k^{\text{th}}$-order recurrence relation is determined by the $k$ preceding terms, it is useful to think of it as a function on $k$-tuples of consecutive terms and a *linear* recurrence can be represented as a matrix function. For example, the recurrence (2.1) can be written as

$$\begin{pmatrix} s_{n+2} \\ s_{n+1} \end{pmatrix} = \begin{bmatrix} 3 & -2 \\ 1 & 0 \end{bmatrix} \begin{pmatrix} s_{n+1} \\ s_n \end{pmatrix}, \quad \begin{pmatrix} s_1 \\ s_0 \end{pmatrix} = \begin{pmatrix} 3 \\ 2 \end{pmatrix}, \tag{2.3}$$

while the usual Fibonacci numbers are given by the matrix equation

$$\begin{pmatrix} f_{n+2} \\ f_{n+1} \end{pmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} \begin{pmatrix} f_{n+1} \\ f_n \end{pmatrix}, \quad \begin{pmatrix} f_1 \\ f_0 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}. \tag{2.4}$$

In each case the matrix takes us from one pair of consecutive terms to the next.

In general, we express (HL) in matrix form by introducing the vectors

$$S_n = \begin{pmatrix} s_{n+k-1} \\ \vdots \\ s_{n+1} \\ s_n \end{pmatrix} \text{ for all } n \geq 0, \tag{2.5}$$

and finding a matrix $A$ such that $S_{n+1} = AS_n$. Since $s_{n+k} = c_1 s_{n+k-1} + c_2 s_{n+k-2} + \cdots + c_k s_n$, this is accomplished using the matrix

$$A = \begin{bmatrix} c_1 & c_2 & \dots & c_{k-1} & c_k \\ 1 & 0 & \dots & 0 & 0 \\ \dots & & & & \\ 0 & 0 & \dots & 1 & 0 \end{bmatrix}, \tag{2.6}$$

which when applied to $S_n$ shifts the components down one entry and puts the next term, $s_{n+k}$, in the first component. The matrix $A$ is called the **companion matrix of the recurrence**. It is also referred to as the **companion matrix of the polynomial** $x^k - c_1 x^{k-1} - \cdots - c_k$.

When the matrix form is used in an initial value problem,

$$S_1 = AS_0\ ,\ S_2 = AS_1 = A^2S_0\ ,\ S_3 = AS_2 = A^3S_0 \text{ and so on,}$$

and this inductively gives the **matrix form**

$$S_n = A^nS_0\,. \tag{2.7}$$

For our examples the matrix form of the usual Fibonacci recurrence is

$$\begin{pmatrix} f_{n+1} \\ f_n \end{pmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}^n \begin{pmatrix} 1 \\ 0 \end{pmatrix}\ ,$$

and the matrix form of the recurrence (2.1) with initial conditions $s_0 = 2$ and $s_1 = 3$ is given by

$$S_n = \begin{bmatrix} 3 & -2 \\ 1 & 0 \end{bmatrix}^n \begin{pmatrix} 3 \\ 2 \end{pmatrix}. \tag{2.8}$$

For formula (2.7) to be useful, we must be able to quickly compute powers of the companion matrix. Matrix powers can be easily computed for diagonal matrices, and the next easiest are matrices which are **diagonalizable**, namely, square matrices which have a basis of eigenvectors. When $A$ is not diagonalizable, a modification involving Jordan matrices can be used. (Refer to Appendix C.)

Returning to the second example above, we set

$$A = \begin{bmatrix} 3 & -2 \\ 1 & 0 \end{bmatrix},$$

which has characteristic polynomial

$$ch_A(x) = \det\left(\begin{bmatrix} 3-x & -2 \\ 1 & -x \end{bmatrix}\right) = x^2 - 3x + 2 = (x-2)(x-1)\,,$$

with *distinct* eigenvalues, $\lambda_1 = 2$ and $\lambda_2 = 1$, and so $A$ is diagonalizable. To find the eigenvectors, solve $(A - 2I)\mathbf{v}_1 = 0$ and $(A - 1I)\mathbf{v}_2 = 0$ to obtain

$$\mathbf{v}_1 = \begin{pmatrix} 2 \\ 1 \end{pmatrix} \text{ and } \mathbf{v}_2 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}.$$

Therefore, for

$$D = \begin{bmatrix} 2 & 0 \\ 0 & 1 \end{bmatrix} \text{ and } P = \begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix},$$

powers can be computed using

$$\begin{aligned} A^n &= P\,D^n\,P^{-1} \\ &= \begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 2 & 0 \\ 0 & 1 \end{bmatrix}^n \begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix}^{-1} \\ &= \begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 2^n & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ -1 & 2 \end{bmatrix} \\ &= \begin{bmatrix} 2^{n+1}-1 & -2^{n+1}+2 \\ 2^n-1 & -2^n+2 \end{bmatrix}. \end{aligned}$$

Equation (2.8) therefore becomes

$$S_n = \begin{bmatrix} 3 & -2 \\ 1 & 0 \end{bmatrix}^n \begin{pmatrix} 3 \\ 2 \end{pmatrix} = \begin{bmatrix} 2^{n+1}-1 & -2^{n+1}+2 \\ 2^n-1 & -2^n+2 \end{bmatrix} \begin{pmatrix} 3 \\ 2 \end{pmatrix} = \begin{pmatrix} 2^{n+1}+1 \\ 2^n+1 \end{pmatrix},$$

from which we derive $s_n = 2^n+1$ for all $n \geq 0$. A change of initial conditions is equivalent to changing the vector $S_0 = (3,2)^T$ and the same procedure would yield the general term of the sequence generated from those new initial conditions. You apply this method to the Fibonacci sequence in Exercise 2.10.

Our general procedure for the case when $A$ is diagonalizable can be summarized as follows:

*Let $s_{n+k} = c_1 s_{n+k-1} + c_2 s_{n+k-2} + \cdots + c_k s_n$ and $S_0 = \boldsymbol{\alpha}$. Suppose the companion matrix $A$ of this recurrence is diagonalizable with eigenvalues $\lambda_1, \ldots, \lambda_k$.*

1. *Find a basis for $\mathbb{C}^k$ consisting of eigenvectors of $A$.*
2. *Use the basis elements as columns to form the matrix $P$.*
3. *Form the diagonal matrix $D$ with the eigenvalues on the diagonal, written in the order corresponding to the columns of $P$.*
4. *Compute $PD^nP^{-1} = A^n$ and the vector $A^n\boldsymbol{\alpha}$.*

*Then the solution is the sequence of first components of the vectors $A^n\boldsymbol{\alpha}$.*

## 2.3 A Simpler Basis for the Solution Space

In Section 2.1 we saw that the set of solutions to any homogeneous linear recurrence relation is a vector space, and in the last section the eigenvalues of the companion matrix played a key role in solving recurrences. In this section we explore more properties of these eigenvalues, with the object of

obtaining a simple basis for the solution space. We will refer to

$$ch(x) = x^k - c_1 x^{k-1} - \cdots - c_{k-1}x - c_k$$

as the **characteristic polynomial of the recurrence relation (HL)** and its roots (the eigenvalues of the companion matrix) will be called the **eigenvalues of the recurrence**.

We can obtain an eigenvector associated with an eigenvalue $\lambda$ by reminding ourselves that pre-multiplying a vector $\mathbf{v} = (v_1, \ldots, v_k)^T$ by the companion matrix $A$ shifts its first $k-1$ components down one position and inserts $c_1 v_1 + c_2 v_2 + \cdots + c_k v_k$ into the first position. Since $\lambda$ is an eigenvalue, $c_1\lambda^{k-1} + c_2\lambda^{k-1} + \cdots + c_k = \lambda^k$ and

$$A(\lambda^{k-1}, \lambda^{k-2}, \ldots, \lambda, 1)^T = (\lambda^k, \lambda^{k-1}, \ldots, \lambda)^T = \lambda(\lambda^{k-1}, \lambda^{k-2}, \ldots, \lambda, 1)^T;$$

that is,

$$\mathbf{v}_\lambda = (\lambda^{k-1}, \lambda^{k-2}, \ldots, \lambda, 1)^T$$

is an eigenvector associated with $\lambda$.

**Theorem 2.3.1.** *Consider the homogeneous recurrence*

$$s_n = c_1 s_{n-1} - \cdots - c_{k-1}s_{n-k+1} - c_k s_{n-k}\,.$$

*(a) If $\lambda$ is an eigenvalue of the recurrence then $\pi^{-1}(\boldsymbol{v}_\lambda) =< \lambda^n >$, where $\pi$ is the vector space isomorphism in (2.2). Consequently, $< \lambda^n > \in \mathcal{X}$.*

*(b) If the recurrence has $k$ distinct eigenvalues $\lambda_1, \ldots, \lambda_k$, then the $k$ sequences $< \lambda_1^n >, \ldots, < \lambda_k^n >$ form a basis for $\mathcal{X}$ and every solution $< s_n >$ has the form*

$$s_n = a_1\lambda_1^n + a_2\lambda_2^n + \cdots + a_k\lambda_k^n \tag{2.9}$$

*for some $a_1, \ldots, a_k \in \mathbb{C}$.*

*Proof.* In part (a), for all $n \geq 0$ we define $s_n = \lambda^n$ and the associated vectors $S_n$ as in (2.5). Then $S_n = \lambda^n \mathbf{v}_\lambda$ and

$$AS_n = \lambda^n A\mathbf{v}_\lambda = \lambda^{n+1}\mathbf{v}_\lambda = S_{n+1}\,,$$

implying $< s_n > \in \mathcal{X}$ with initial vector $S_0 = \mathbf{v}_\lambda$. When the recurrence has $k$ distinct eigenvalues, the eigenvectors $\mathbf{v}_{\lambda_i}$ form a basis for $\mathbb{C}^k$, and the $k$ sequences $< \lambda_i^n > = \pi^{-1}(\mathbf{v}_{\lambda_i})$ therefore form a basis for $\mathcal{X}$. Since every element of a vector space can be uniquely written as a linear combination of basis vectors, this implies that (2.9) holds. □

### *2.3.1 Distinct eigenvalues*

For a specific vector $S_0 = \boldsymbol{\alpha}$ of initial conditions, how do we solve the initial value problem, that is, how do we find the $a_i$ in (2.9)? Using the inherent linear algebra and the isomorphism $\pi$, this becomes the equivalent problem of solving

$$a_1 \mathbf{v}_{\lambda_1} + \cdots + a_k \mathbf{v}_{\lambda_k} = \boldsymbol{\alpha}$$

for the coefficients $a_1, \cdots, a_k$. Translating this to a matrix equation, we want to find (the unique) $a_1, \cdots, a_k \in \mathbb{C}$ such that

$$\begin{bmatrix} 1 & 1 & 1 & \dots & 1 \\ \lambda_1 & \lambda_2 & \lambda_3 & \dots & \lambda_k \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \lambda_1^{k-1} & \lambda_2^{k-1} & \lambda_3^{k-1} & \dots & \lambda_k^{k-1} \end{bmatrix} \begin{pmatrix} a_1 \\ a_2 \\ \vdots \\ a_k \end{pmatrix} = \boldsymbol{\alpha}. \tag{2.10}$$

The coefficient matrix in (2.10) is called the **Vandermonde matrix** associated with $\lambda_1, \ldots, \lambda_k$. It is named after Alexandre Vandermonde (1735-1796), the developer of the modern theory of determinants, and has applications in almost every area of mathematics. For an interesting survey of its many applications, consult [71].

For $c_1^2 \neq -4c_2$, we consider the recurrence

$$s_{n+2} = c_1 s_{n+1} + c_2 s_n, \quad S_0 = \boldsymbol{\alpha}\,,$$

whose characteristic polynomial $ch(x) = x^2 - c_1 x - c_2$ has distinct eigenvalues since its discriminant $D = c_1^2 + 4c_2 \neq 0$. (In Exercise 2.18 you answer the question of what happens when $D$ is zero.) The eigenvalues of the recurrence are

$$\lambda_1 = (c_1 + \sqrt{D})/2 \text{ and } \lambda_2 = (c_1 - \sqrt{D})/2\,,$$

with associated Vandermonde matrix

$$V = \begin{bmatrix} 1 & 1 \\ \lambda_1 & \lambda_2 \end{bmatrix}.$$

The determinant of this matrix is $\lambda_2 - \lambda_1 = -\sqrt{D} \neq 0$ and its inverse is

$$V^{-1} = \frac{1}{\sqrt{D}} \begin{bmatrix} -\lambda_2 & 1 \\ \lambda_1 & -1 \end{bmatrix}.$$

You're asked to complete this calculation in Exercise 2.11.

Because each vector of initial conditions specifies an element of $\mathcal{X}$, the system of equations in (2.10) has a unique solution for each $\boldsymbol{\alpha}$, which means that the Vandermonde matrix must always be invertible. Turning to the

question of calculating the inverse, for all $i = 1, \ldots, k$ we define the auxiliary polynomial

$$P_i(x) = \prod_{\substack{j=1 \\ j \neq i}}^{k} (x - \lambda_j) = b_{i,1} + b_{i,2}x + \ldots + b_{i,k}x^{k-1} .$$

Since the $\lambda_i$ are distinct, for all $i$ $P_i(\lambda_i) \neq 0$, and for $j \neq i$, $P_i(\lambda_j) = 0$.

**Theorem 2.3.2.** *Let $B$ be the $k \times k$ matrix whose $i$-th row records the coefficients of the polynomial $P_i(x)$ (written according to increasing powers of $x$). Then $V^{-1} = DB$, where $D$ is the diagonal matrix which has the (non-zero) diagonal entries $1/P_1(\lambda_1), \ldots, 1/P_k(\lambda_k)$.*

*Proof.* This proof is inspired by a mathematical note by F.D.Parker [108]. The diagonal entries in the matrix product $BV$ have the form

$$(b_{i,1}, b_{i,2}, \ldots, b_{i,k})(1, \lambda_i, \ldots, \lambda_i^{k-1})^T = P_i(\lambda_i) \neq 0 ,$$

and the nondiagonal entries are

$$(b_{i,1}, b_{i,2}, \ldots, b_{i,k})(1, \lambda_j, \ldots, \lambda_j^{k-1})^T = P_i(\lambda_j) = 0.$$

Therefore, $DBV$ is the identity and $V^{-1} = DB$, as claimed. □

We use this result to find an explicit closed form for the general term of

$$s_{n+2} = s_{n+1} + 2s_n \, , \; S_0 = (1, 5)^T.$$

Since the characteristic polynomial $ch(x) = x^2 - x - 2 = (x-2)(x+1)$ has roots $\lambda_1 = 2, \lambda_2 = -1$; $P_1(x) = 1 + x$ , $P_2(x) = -2 + x$ give

$$B = \begin{bmatrix} 1 & 1 \\ -2 & 1 \end{bmatrix} ,$$

and from $P_1(2) = 3, P_2(-1) = -3$ we have

$$V^{-1} = \frac{1}{3} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} B = \frac{1}{3} \begin{bmatrix} 1 & 1 \\ 2 & -1 \end{bmatrix} .$$

Therefore,

$$\begin{pmatrix} a_1 \\ a_2 \end{pmatrix} = V^{-1} \begin{pmatrix} 1 \\ 5 \end{pmatrix} = \frac{1}{3} \begin{bmatrix} 1 & 1 \\ 2 & -1 \end{bmatrix} \begin{pmatrix} 1 \\ 5 \end{pmatrix} = \begin{pmatrix} 2 \\ -1 \end{pmatrix} ,$$

and (2.9) gives $s_n = 2\lambda_1^n - \lambda_2^n = 2^{n+1} + (-1)^{n+1}$.

### *2.3.2 Repeated Eigenvalues*

Beginning with any vector $\mathbf{v} \in \mathbb{C}^k$, we can consider the **orbit** of $\mathbf{v}$ under $A$, which is defined to be the set

$$\{\mathbf{v}, A\mathbf{v}, A^2\mathbf{v}, \dots\}.$$

When $\lambda$ is an eigenvalue, the orbit of $\mathbf{v}_\lambda$ is particularly nice and yields the appealing sequence $< \lambda^n >$ in $\mathcal{X}$. When some eigenvalues are repeated, the vectors $\mathbf{v}_{\lambda_i}$ still form a linearly independent set but the set has fewer than $k$ elements. Although we will do considerably more work to identify enough additional vectors with good orbits to form a basis, the actual result (given in Theorem 2.3.6 below) is only slightly more complicated.

If $\lambda$ is an eigenvalue of multiplicity $m$, there exists a polynomial $f(x)$ with complex coefficients such that

$$ch(x) = (x-\lambda)^m f(x) \text{ where } f(\lambda) \neq 0.$$

Denoting the differentiation operator on the space of polynomials by $D$, this implies that the value of $D^j(ch(x))$ at $x = \lambda$ equals zero for all $1 \leq j < m$ and is non-zero for $j = m$. Using the fact that

$$D^j(x^i) = \begin{cases} \dfrac{i!}{(i-j)!}x^{i-j} & \text{if } j \leq i \\ 0 & \text{if } j > i \end{cases},$$

in Exercise 2.13 you prove the polynomial identity

$$D^j(x^i) = xD^j(x^{i-1}) + jD^{j-1}(x^{i-1}), \tag{2.11}$$

where $D^0$ is the identity operator and the second summand on the right side should be interpreted as zero when $j = 0$.

We now construct what are called the cyclic subspaces of $\mathbb{C}^k$ associated with the eigenvalues of $A$. This study is motivated by another concept in linear algebra, the **Rational Canonical Form**, a topic which is more advanced than our review in Appendix C. Because our description is explicit, what follows can be viewed as an illustration of Rational Canonical Form and you can consult [67, Sections 7.1-7.2] if you're interested in more information.

**Proposition 2.3.3.** *If $\lambda$ is an eigenvalue of multiplicity $m$ then*

$$A\boldsymbol{v}_\lambda^{(j)} = \lambda\boldsymbol{v}_\lambda^{(j)} + j\boldsymbol{v}_\lambda^{(j-1)} \text{ for all } 0 \leq j < m, \tag{2.12}$$

*where $\boldsymbol{v}_\lambda^{(-1)} = 0$, $\boldsymbol{v}_\lambda^{(0)} = \boldsymbol{v}_\lambda$, and for all $j \geq 1$, $\boldsymbol{v}_\lambda^{(j)}$ denotes the element of $\mathbb{C}^k$ obtained by applying $D^j$ to each component of $(x^{k-1}, \dots, x, 1)^T$ and then evaluating each component at $x = \lambda$.*

*Proof.* Setting $\mathbf{x} = (x^{k-1}, \ldots, x, 1)^T$, we first will prove

$$AD^j(\mathbf{x}) = xD^j(\mathbf{x}) + jD^{j-1}(\mathbf{x}) \text{ for all } j < m\,. \tag{2.13}$$

From the linearity of the operator $D$ we obtain

$$AD^j(\mathbf{x}) = D^j(A\mathbf{x}).$$

Recalling that for any vector $\mathbf{v}$, the last $k-1$ components of $A\mathbf{v}$ are obtained by shifting down the first $k-1$ components of $\mathbf{v}$, for any $i \neq 1$ the $i^{\text{th}}$ component of $AD^j(\mathbf{x})$ is $D^j(x^{k-i+1})$ which from (2.11) equals $xD^j(x^{k-i}) + jD^{j-1}(x^{k-i})$. This proves the equality of the last $k-1$ components in the vector equation (2.13).

The first component of $A\mathbf{x}$ is

$$c_1x^{k-1} + \ldots c_{k-1}x + c_k = x^k - ch(x),$$

which implies that the first component of $D^j(A\mathbf{x})$ is

$$\begin{aligned} D^j(x^k - ch(x)) &= D^j(x^k) - D^j(ch(x)) \\ &= xD^j(x^{k-1}) + jD^{j-1}(x^{k-1}) - D^j(ch(x)), \end{aligned}$$

again from (2.11). Equality of the first components in (2.12) is obtained from this and the fact that $D^j(ch(x))$ has value 0 at $x = \lambda$. □

For the **cyclic subspace corresponding to** $\lambda$

$$X_\lambda = \text{Span}\{\mathbf{v}_\lambda^{(0)}, \ldots, \mathbf{v}_\lambda^{(m-1)}\}\,, \tag{2.14}$$

this proposition implies that $A$ maps $X_\lambda$ into $X_\lambda$, which means that $A$ is an *operator* on the subspace $X_\lambda$ and this allows us to apply the theory of linear algebra to the vector space $X_\lambda$.

**Theorem 2.3.4.** *If $\lambda$ is an eigenvalue of multiplicity $m$ then $(x-\lambda)^m$ is the minimal polynomial of the operator $A$ on $X_\lambda$, and $S = \{\boldsymbol{v}_\lambda^{(0)}, \ldots, \boldsymbol{v}_\lambda^{(m-1)}\}$ is a basis for $X_\lambda$.*

*Proof.* From the definition of $X_\lambda$ in (2.14), $\dim(X_\lambda) \leq m$ and $S$ is a generating set for $X_\lambda$. It suffices to prove $\dim(X_\lambda) \geq m$. We do this by showing that the degree of $min$, the minimal polynomial of $A$ restricted to the subspace $X_\lambda$, equals $m$. In fact we prove that $min(x) = (x-\lambda)^m$. To prove this we show for all $0 \leq j < m$,

$$(A - \lambda I)^j \mathbf{v}_\lambda^{(j)} \neq \mathbf{0} \text{ and } (A - \lambda I)^m \mathbf{v}_\lambda^{(j)} = \mathbf{0}. \tag{2.15}$$

Fix $j < m$. Then (2.12) can be rewritten in the form

$$(A - \lambda I)\mathbf{v}_\lambda^{(j)} = j\mathbf{v}_\lambda^{(j-1)}\,.$$

Repeated use of this gives

$$\begin{aligned}(A-\lambda I)^j \mathbf{v}_\lambda^{(j)} &= (A-\lambda I)^{j-1}((A-\lambda I)\mathbf{v}_\lambda^{(j)}) \\ &= j(A-\lambda I)^{j-1}\mathbf{v}_\lambda^{(j-1)} \\ &= j(A-\lambda I)^{j-2}((A-\lambda I)\mathbf{v}_\lambda^{(j-1)}) \\ &= j(j-1)\cdot(A-\lambda I)^{j-2}\mathbf{v}_\lambda^{(j-2)} \\ &\;\;\vdots \\ &= j!\mathbf{v}_\lambda^{(0)}\,,\end{aligned}$$

which means

$$(A-\lambda I)^j \mathbf{v}_\lambda^{(j)} = j!\mathbf{v}_\lambda^{(0)} \text{ for all } j < m. \tag{2.16}$$

Since $\mathbf{v}_\lambda^{(0)}$ is an eigenvector, it is non-zero and therefore $(A-\lambda I)^j$ is not the zero transformation on $X_\lambda$, which proves that $min(x) \neq (x-\lambda)^j$ for all $j < m$. On the other hand, from (2.16) we also obtain for all $j = 0, 1, \ldots, m-1$

$$\begin{aligned}(A-\lambda I)^m \mathbf{v}_\lambda^{(j)} &= j!\cdot(A-\lambda I)^{m-j}\mathbf{v}_\lambda^{(0)} \\ &= j!\cdot(A-\lambda I)^{m-j-1}((A-\lambda I)\mathbf{v}_\lambda^{(0)}) \\ &= (A-\lambda I)^{m-j-1}\mathbf{0} = \mathbf{0}\,,\end{aligned}$$

since $\mathbf{v}_\lambda^{(0)} = \mathbf{v}_\lambda$ is an eigenvector corresponding to $\lambda$. Therefore, $(A-\lambda I)^m$ *is* the zero transformation on the basis $\{\mathbf{v}_\lambda^{(0)} \ldots, \mathbf{v}_\lambda^{(m-1)}\}$, and by linearity, $(A-\lambda I)^m$ must be zero on all of $X_\lambda$. This proves $min(x) = (x-\lambda)^m$. □

**Corollary 2.3.5.** *Let $\lambda_1, \ldots, \lambda_t$ be the different eigenvalues of the recurrence, with corresponding multiplicities $m_1, \ldots, m_t$. Then*

$$\{\boldsymbol{v}_{\lambda_1}^{(0)}, \ldots, \boldsymbol{v}_{\lambda_1}^{(m_1-1)}\,,\; \ldots\;,\; \boldsymbol{v}_{\lambda_t}^{(0)}, \ldots, \boldsymbol{v}_{\lambda_t}^{(m_t-1)}\} \tag{2.17}$$

*is a basis for $\mathbb{C}^k$.*

*Proof.* For each $i$, let $X_i = X_{\lambda_i}$ and let $min_i(x)$ denote the minimal polynomial of $A$ restricted to $X_i$. For $i \neq j$, $X_i \cap X_j$ is an $A$-invariant subspace whose minimal polynomial divides each of $min_i(x), min_j(x)$ and so also divides their greatest common denominator. But $\lambda_i \neq \lambda_j$ gives $\gcd((x-\lambda_i)^m, (x-\lambda_j)^m) = 1$, implying $min(x) = 1$ and $X_i \cap X_j = \{0\}$. Therefore,

$$\text{Span}(X_1 \cup X_2 \cup \ldots \cup X_t) = \text{Span}(X_1) \cup \ldots \cup \text{Span}(X_t) = X_1 \cup X_2 \cup \ldots \cup X_t\,,$$

giving

$$\dim(X_1 \cup X_2 \cup \ldots \cup X_t) = m_1 + \ldots + m_t,$$

which does equal $k = \deg(ch)$ since the characteristic polynomial is

$$ch(x) = (x - \lambda_1)^{m_1} \dots (x - \lambda_t)^{m_t}.$$

□

This basis is the simpler one we promised earlier, and gives the following result for the solution of the general homogeneous linear recurrence.

**Theorem 2.3.6.** *Let $\lambda_1, \dots, \lambda_t \in \mathbb{C}$ be distinct. Then $< s_n >$ is a solution to the recurrence (HL) with $ch(x) = (x - \lambda_1)^{m_1} \dots (x - \lambda_t)^{m_t}$ if and only if the $n^{th}$ term of $< s_n >$ has the form*

$$s_n = a_1(n)\lambda_1^n + \dots + a_t(n)\lambda_t^n \tag{2.18}$$

*where each $a_i(x)$ is a polynomial whose degree is less than $m_i$.*

*Proof.* In Exercise 2.15 you check that each sequence whose $n^{\text{th}}$ term has the form given in (2.18) satisfies (HL). If $\{\mathbf{b}_1, \dots, \mathbf{b}_t\}$ is the basis in (2.17) the fact that $\pi$ is an isomorphism from $\mathcal{X}$ to $\mathbb{C}^k$ implies that every solution to (HL) has the form

$$< s_n >= \beta_1 \pi^{-1}(\mathbf{b}_1) + \dots + \beta_t \pi^{-1}(\mathbf{b}_t)\,, \tag{2.19}$$

for some complex constants $\beta_1, \dots, \beta_t$. Each $\mathbf{b}_j$ equals $\mathbf{v}_\lambda^{(i)}$ for some eigenvalue $\lambda$ and some $i$ less than the multiplicity of $\lambda$, and the $n^{\text{th}}$ term of $\pi^{-1}(\mathbf{v}_\lambda^{(i)})$ is $D^i(x^n)$ evaluated at $x = \lambda$, that is, equals

$$\begin{cases} \dfrac{n!}{(n-i)!}\lambda^{n-i} & \text{if } j \le i \\ 0 & \text{if } j > i \end{cases},$$

where the coefficient $n!/(n-i)!$ is a polynomial in $n$ whose degree equals $i$. Therefore, (2.19) yields the result. □

As an illustration of the method consider the following initial value problem.

$$s_{n+4} = s_{n+3} + 3s_{n+2} - 5s_{n+1} + 2s_n \text{ with } s_0 = 1, s_1 = -1, s_2 = 0, s_3 = 1\,.$$

We have $ch(x) = x^4 - x^3 - 3x^2 + 5x - 2$ and the companion matrix is

$$A = \begin{bmatrix} 1 & 3 & -5 & 2 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

with eigenvalues $\lambda_1 = 1, \lambda_2 = -2$ and respective multiplicities $m_1 = 3$, $m_2 = 1$. Therefore, there exist polynomials $a_1(x), a_2(x)$ such that any solution $< s_n >$ to the recurrence has general term

$$s_n = a_1(n)(1)^n + a_2(n)(-2)^n .$$

Further, $\deg(a_1) < 3$ and $\deg(a_2) < 1$ imply there exist constants $\beta_i \in \mathbb{C}$ such that

$$a_1(x) = \beta_0 + \beta_1 x + \beta_2 x^2 \, ; \, a_2(x) = \beta_3$$

which gives

$$s_n = (\beta_0 + \beta_1 n + \beta_2 n^2) + \beta_3(-2)^n .$$

Using the initial values, this reduces to solving a system of four equations in the unknowns $\beta_0, \beta_1, \beta_2, \beta_3$ with augmented matrix

$$\begin{bmatrix} 1 & 0 & 0 & 1 & 1 \\ 1 & 1 & 1 & -2 & -1 \\ 1 & 2 & 4 & 4 & 0 \\ 1 & 3 & 9 & -8 & 1 \end{bmatrix} .$$

Gaussian elimination can be used to obtain the row reduced form

$$\begin{bmatrix} 1 & 0 & 0 & 0 & 8/9 \\ 0 & 1 & 0 & 0 & -8/3 \\ 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 1/9 \end{bmatrix} ,$$

and from this $a_1(x) = \frac{8}{9} - \frac{8}{3}x + x^2$ and $a_2(x) = \frac{1}{9}$, giving

$$s_n = \frac{8}{9} - \frac{8}{3}n + n^2 + \frac{1}{9}(-2)^n .$$

The $k \times k$ matrix $V$ whose columns consist of the vectors $\mathbf{v}_\lambda^{(i)}$ (written as columns according to the order given in (2.17)) can be called the **generalized Vandermonde matrix** associated with the recurrence. In order to find the polynomials $a_1(x), \ldots, a_t(x)$ corresponding to the initial conditions $S_0 = \boldsymbol{\alpha}$, we solve the matrix equation $V\mathbf{x} = \boldsymbol{\alpha}$. As with the standard Vandermonde matrix, the fact that every initial value problem for (HL) has a unique solution translates to the invertibility of the generalized Vandermonde matrix. The explicit form of $V^{-1}$ is found in Exercise 2.17.

## 2.4 The Asymptotic Behavior of Solutions

We will now use the matrix form $S_n = A^n S_0$ to estimate the asymptotic size of solutions to (HL) without computing $A^n$. The basic idea is to write

the initial condition $S_0$ as a linear combination of $k$ linearly independent vectors $C_1, \ldots, C_k$ for which the asymptotic size of $A^n C_i$ is known.

We consider the principal case when $A$ is a diagonalizable matrix which means there exists a basis $\{C_1, \ldots, C_k\}$ of eigenvectors for $\mathbb{C}^k$, and write $S_0 = \sum_{i=1}^{k} a_i C_i$ for some $a_i \in \mathbb{C}$. If this basis happens to be an *orthogonal* set (in which case the matrix $A$ is said to be **orthogonally diagonalizable**) then $C_j^T C_i = 0$ when $i \neq j$ and so

$$C_j^T S_0 = C_j^T \sum_{i=1}^{k} a_i C_i = \sum_{i=1}^{k} a_i C_j^T C_i = a_j \|C_j\|^2 ,$$

giving $a_j = C_j^T X_0 / \|C_j\|^2$ where $\|C\|$ is the Euclidean length of the vector $C$. Orthogonality therefore gives an easy expansion of $S_0$ in terms of the column eigenvectors of $A$. Although the orthogonality of the eigenvectors can't be guaranteed, we will see that a helpful property which we call biorthogonality always holds when $A$ is diagonalizable.

If $\lambda$ is an eigenvalue each of the two homogeneous systems $(A - \lambda I)X = 0$ and $X(A - \lambda I) = 0$ has a non-zero solution. For our purposes, we'll call a non-zero solution to $(A - \lambda I)X = 0$ a **column eigenvector** while a non-zero solution to the second system will be referred to as a **row eigenvector**. Letting $Z$ be the $k \times k$ matrix whose columns are the column eigenvectors $C_1, \ldots, C_k$ then $AZ = ZD$ where $D$ is the diagonal matrix with diagonal entries $\lambda_1, \ldots, \lambda_k$. From the linear independence of its columns, $Z$ is an invertible matrix and $Z^{-1}A = DZ^{-1}$, which means the $i^{\text{th}}$ row $R_i$ of $Z^{-1}$ is a row eigenvector corresponding to $\lambda_i$. From $Z^{-1}Z = I$ we also know that $R_i C_j = 0$ when $i \neq j$, and we call the two sets of vectors $\{R_1, \ldots, R_k\}$ and $\{C_1, \ldots, C_k\}$ a pair of **biorthogonal sets**. (Also note that $R_i C_i = 1$ for all $i$.) Mimicking the orthogonal case, from $S_0 = a_1 C_1 + \ldots + a_k C_k$ we have

$$R_i S_0 = R_i(a_1 C_1 + \ldots + a_k C_k) = a_i R_i C_i = a_i .$$

Therefore, $S_0 = \sum_{i=1}^{k} (R_i S_0) C_i$, and

$$S_n = A^n S_0 = \sum_{i=1}^{k} (R_i S_0)(A^n C_i) , \tag{2.20}$$

which can be recorded as

**Lemma 2.4.1.** *Let $A$ be any diagonalizable $k \times k$ matrix. Let $Z$ be the matrix whose columns are the linearly independent eigenvectors $C_1, \ldots, C_k$, and let $R_1, \ldots, R_k$ be the rows of $Z^{-1}$. Then any solution to $S_{n+1} = AS_n$ can be written in the form*

$$S_n = \sum_{i=1}^{k} (R_i S_0) \lambda_i^n C_i .$$

When $A$ has fewer than $k$ linearly independent eigenvectors, let $Z$ be the matrix whose columns are the generalized eigenvectors. Since they form a basis for $\mathbb{C}^k$, the above argument still yields (2.20) where $R_1, \ldots, R_k$ are the rows of the matrix $Z^{-1}$. Even in this more general case when $\lambda_1$ is a *simple* eigenvector, then

$$(2.21)\quad S_n = A^n S_0 = \sum_{i=1}^{k} (R_i S_0)(A^n C_i) = (R_1 S_0)\lambda_1^n C_1 + \sum_{i=2}^{k} (R_i S_0) A^n C_i \,,$$

does hold, where each $C_i$ is a generalized eigenvector corresponding to some eigenvalue $\lambda \neq \lambda_1$.

Ordering the *different* eigenvalues according to decreasing complex modulus:

$$(2.22)\qquad |\lambda_1| \geq |\lambda_2| \geq \ldots \geq |\lambda_t| \text{ (where } t \leq k) \quad ,$$

$\lambda_1$ is called a **dominant eigenvalue of the recurrence**, and if $|\lambda_1| > |\lambda_2|$ holds then $|\lambda_1|$ is called the **strictly dominant eigenvalue**. (Note that strict dominance not prevent $\lambda_1$ from being a multiple eigenvalue.)

For the case when $\lambda_1$ is a *simple strictly dominant* eigenvalue, (2.21) gives

$$\frac{S_n}{\lambda_1^n} = (R_1 S_0) C_1 + Y_n \,,$$

where $Y_n = \sum_{i=2}^{k} (R_i S_0) A^n C_i / \lambda_1^n$. Since each coordinate $|Y_n|_i$ of $|Y_n|$ is bounded above by $|\lambda_2|^n$ times a polynomial in $n$ (refer to Theorem 2.3.6), the limit $\lim_{n\to\infty} |Y_n|_i / \lambda_1^n$ is zero. Therefore,

$$\lim_{n\to\infty} \frac{S_n}{\lambda_1^n} = (R_1 S_0) C_1,$$

where $C_1$ is a column eigenvector of $\lambda_1$ and $R_1$ is the specific row eigenvector of $\lambda_1$ chosen above. For any other row eigenvector $R$ corresponding to $\lambda_1$, $(RS_0)/(RC_1) = R_1 S_0$ holds and we obtain

**Theorem 2.4.2.** *If $\lambda_1$ is a simple strictly dominant eigenvalue of the $k \times k$ matrix $A$, then*

$$\lim_{n\to\infty} \frac{S_n}{\lambda_1^n} = aC$$

*where $C$ is any column eigenvector corresponding to $\lambda_1$, $R$ is any row eigenvector corresponding to $\lambda_1$, and $a = (RS_0)/(RC)$.*

The upshot of this result is that for many reasonable homogeneous recurrences and most initial conditions, the asymptotic behavior of a solution can be considered to be one-dimensional. Full details of a solution would require a somewhat complicated formula, but as long as $RS_0 \neq 0$ the solution converges in a ratio sense to a single vector. When the dot product

$RS_0$ is zero, the theorem only says that the solution grows more slowly than $|\lambda_1|^n$ and it gives no more information about the solution. The other main point is that the value of $a$ can be computed without computing the rest of the coefficients in the full expansion of the solution.

## 2.5 Exercises

**Ex 2.1.** Show that every initial value problem has a unique solution.

**Ex 2.2.** Use induction to verify the closed formulas for the recurrence relation (2.1) under each of the following initial conditions: $S_0 = (3,2)^T$ ; $S_0 = (1,3)^T$.

**Ex 2.3.** Suppose that the sequence $< s_n >$ satisfies the Fibonacci recurrence and $s_3 = 5, s_6 = -2$. What is $s_4$? What is $s_{10}$? What are the initial values $s_0$ and $s_1$?

**Ex 2.4.** Construct an argument which concludes there is a sequence $< s_n >$ which satisfies the Fibonacci recurrence for which $s_{20} = 23$ and $s_{1000} = 56$.

**Ex 2.5.** Show directly that a linear combination of solutions to (HL) is itself a solution.

**Ex 2.6.** For a fixed polynomial $Q$ in $k \geq 2$ variables consider the set $V$ of all sequences $< s_n >$ which are solutions to the $k^{\text{th}}$-order recurrence relation

$$s_{n+k} = Q(s_n, s_{n+1}, \ldots, s_{n+k-1}) .$$

If $V$ is a vector space, show that $Q$ must be a linear with zero constant term.

We know that the set of all sequences of complex numbers forms a complex vector space under the operations of coordinatewise addition and scaling. Applying the definition of linear independence (refer to Appendix C) to this vector space, we see that sequences $\phi_1(n), \ldots, \phi_k(n)$ are **linearly independent** if

$$b_1\phi_1(n) + \ldots + b_k\phi_k(n) = 0 \text{ for all } n \geq 0 \implies b_1 = b_2 = \ldots = b_k = 0 .$$

**Ex 2.7.** Show that every $k^{\text{th}}$-order linear difference equation has $k$ linearly independent solutions.

**Ex 2.8.** Let $\phi_1(n), \ldots, \phi_k(n)$ be $k$ linearly independent solutions to a $k^{\text{th}}$-order homogeneous linear difference equation. Show that the solution to any initial value problem for this difference equation can be written in the form $\sum_{i=1}^{k} b_i\phi_i(n)$.

**Ex 2.9.** Let $\phi_1(n), \ldots, \phi_k(n)$ be solutions to a $k^{\text{th}}$-order linear difference equation. Show that the following three statements are equivalent:

1. $\sum_{i=1}^{k} b_i \phi_i(n)$ is the zero sequence.
2. For all $0 \leq n < k$, $\sum_{i=1}^{k} b_i \phi_i(n) = 0$.
3. $\sum_{i=1}^{k} b_i \phi_i(n) = 0$ for $k$ consecutive values of $n$.

**Ex 2.10.** Use diagonalization to obtain Binet's Formula (1.5), the closed form for the Fibonacci numbers.

**Ex 2.11.** Let $s_{n+2} = c_1 s_{n+1} + c_2 s_{n+2}$ be a second-order recurrence relation with eigenvalues $\lambda_1 = (c_1 + \sqrt{D})/2$ and $\lambda_2 = (c_1 - \sqrt{D})/2$ where $D = c_1^2 - 4c_2$. If $D \neq 0$ show that the sequence whose $n^{\text{th}}$ term is

$$\frac{1}{\sqrt{D}}\Big(-\alpha_0 \lambda_1 \lambda_2 (\lambda_1^{n-1} - \lambda_2^{n-1}) + \alpha_1 (\lambda_1^n - \lambda_2^n)\Big)$$

is the unique solution with initial conditions $S_0 = \boldsymbol{\alpha}$.

**Ex 2.12.** Show that the Vandermonde matrix associated with $\lambda_1, \ldots, \lambda_k$ is invertible if and only if the $\lambda_i$ are distinct.
Hint: Pre-multiply the matrix by a row vector where the product for each entry is interpreted as the evaluation of a polynomial at $\lambda_i$.

**Ex 2.13.** Use the product rule for differentiation and induction to substantiate identity (2.11).

**Ex 2.14.** Solve the initial value problem

$$s_{n+3} = 2s_{n+2} + 5s_{n+1} - 6s_n \; ; \;\; s_0 = 9 \, , \, s_1 = -18 \, , \, s_2 = 66.$$

**Ex 2.15.** Check that each sequence whose $n^{\text{th}}$ term has the form given in (2.18) satisfies the recurrence (HL).

**Ex 2.16.** Use Theorem 2.3.6 to find the solution to

$$s_{n+4} = 8s_{n+2} - 16s_n \, , \, S_0 = (-1, 8, 4, 16)^T.$$

**Ex 2.17.** Let $\lambda_1, \cdots, \lambda_t$ be distinct complex numbers and let $m_1, \ldots, m_t$ be any positive integers which sum to $k$. Let $V$ be the $k \times k$ **generalized Vandermonde matrix** associated with this data; that is, the matrix $V$ whose columns are the vectors given in (2.17).

(a) Construct an argument which proves that $V$ is invertible.

(b) If $A$ is the companion matrix for $P(x) = (x-\lambda_1)^{m_1}(x-\lambda_2)^{m_2} \cdots (x-\lambda_t)^{m_t}$, compute $V^{-1}AV$.

(c) For $1 \leq i \leq t$, $1 \leq j \leq m_i$, define $Q(x) = \dfrac{P(x)}{(x-\lambda_i)^j}$ and expand it as a polynomial of degree $k-1$ in $x$:

$$Q(x) = b_0 + b_1 x + \cdots + b_{k-j} x^{k-j} + \cdots + b_{k-l} x^{k-1} \, .$$

Show that $b_{k-j+1} = \ldots = b_{k-1} = 0$; that is, $b_{k-j}$ is the last non-zero coefficient in $Q(x)$.

(d) Show that

$$(b_0, b_1, \cdots, b_{k-l}) \cdot v_{\lambda_i}^{(t)} \begin{cases} \neq 0 & \text{if } j = t \\ = 0 & \text{if } j < t\,. \end{cases}$$

(e) Compute the precise value of the inner product in (d).

(f) Use the information from the previous parts to construct the inverse of $V$.

**Ex 2.18.** Let $s_{n+2} = c_1 s_{n+1} + c_2 s_{n+2}$ be a second-order recurrence relation such that $D = c_1^2 + 4c_2$ is zero. Using the techniques of Section 2.3.2, show that the sequence with general term

$$s_n = \alpha_0 \left(\frac{c_1}{2}\right)^n + n\left(\alpha_1 - \frac{\alpha_0 c_1}{2}\right)\left(\frac{c_1}{2}\right)^{n-1}$$

is the unique solution to (HL) with $S_0 = \boldsymbol{\alpha}$.

**Ex 2.19.** Write a general procedure for solving (HL) given knowledge of the roots of the characteristic polynomials and their multiplicities.

**Ex 2.20.** Show that

$$\lim_{n\to\infty} \frac{a(n+1)}{a(n)} = 1$$

holds for any polynomial $a(x) \in \mathbb{C}[x]$.

In Exercises 2.21 - 2.23, use the ordering of eigenvalues given in (2.22).

**Ex 2.21.** Show that if $|\lambda_1| < 1$, then $\lim_{n\to\infty} s_n = 0$.

**Ex 2.22.** Show that if the coefficients in (HL) are *integers* and $c_k \neq 0$, then $|\lambda_1| \geq 1$.

**Ex 2.23.** Let $s_{n+2} = c_1 s_{n+1} + c_2 s_n$ be a second order recurrence with $c_1 \neq 0$ and two real eigenvalues.

(a) Show that $\lambda_1$ is strictly dominant.

(b) Prove that for any $< s_n > \in \mathcal{X}$ the limit $\lim_{n\to\infty} \frac{s_n}{\lambda_1^n}$ exists. Moreover, when $s_1 \neq s_0\lambda_2$ show that this limit must be non-zero.

**Ex 2.24.** For each $n \geq 1$ let $D_n$ be the determinant of the $n \times n$ tridiagonal matrix

$$\begin{bmatrix} \alpha & 1 & 0 & 0 & \ldots & 0 & 0 & 0 \\ 1 & \alpha & 1 & 0 & \ldots & 0 & 0 & 0 \\ 0 & 1 & \alpha & 1 & \ldots & 0 & 0 & 0 \\ \ldots & & & & & & & \\ 0 & 0 & 0 & 0 & \ldots & 1 & \alpha & 1 \end{bmatrix},$$

where $\alpha$ is some fixed real number. Show that the sequence $< D_n >$ satisfies the difference equation $D_n = \alpha D_{n-1} - D_{n-2}$ with initial conditions $D_0 = 1$,

$D_1 = \alpha$. Find the values of $\alpha$ for which $\lim_{n\to\infty} |D_n| = \infty$. For what values of $\alpha$ is $D_n = O(n)$? What is the asymptotic size of $D_n$ for other values of $\alpha$? Find the values of $\alpha$ for which $D_n$ is periodic, and decide what periods are possible.

# 3
# Finite Difference Equations

## 3.1 Linear Difference Equations

A difference equation is the discrete analog of a differential equation. Although differential equations are typically studied earlier in a mathematical curriculum, there are many respects in which the theory of difference equations is simpler. A **finite difference equation** has the general form

$$s_n = \Phi(s_{n-1}, s_{n-2}, \ldots, s_{n-k}, n), \tag{3.1}$$

where $\Phi$ is a fixed complex-valued function and the integer $k$ is called the **order** of the recurrence. The **initial value problem** for (3.1) is the problem of finding a sequence $< s_n >$ which satisfies (3.1) for a given function $\Phi$ and a fixed initial value vector $S_0 = (s_{k-1}, \ldots, s_1, s_0)$. An initial value problem has only one solution because (as in Chapter 2) for any fixed $k$ initial values $s_0, \ldots, s_{k-1}$, the $k^{\text{th}}$-term is specified by $s_k = \Phi(s_{k-1}, s_{k-2}, \ldots, s_0, k)$, and all successive values are similarly specified. From this we obtain

**Theorem 3.1.1 (Existence and uniqueness theorem).** *Every initial value problem for a finite difference equation has a unique solution.*

The essential ingredient in the proof of this result was the assumption that $\Phi$ is a *function*, that is, that $\Phi$ returns exactly one value for a given input.

The function $\Phi$ is called a **linear function** in the variables $s_{n-1}, \ldots, s_{n-k}$ if there exist $k+1$ complex-valued functions $g_1, \ldots, g_k, \Psi$ defined on the

natural numbers such that for all $n \geq k$

$$\Phi(s_{n-1}, s_{n-2}, \ldots, s_{n-k}, n) = g_1(n)s_{n-1} + \ldots + g_k(n)s_{n-k} + \Psi(n)$$

where $\Psi$ is called the **input** or **forcing function**. When $\Phi$ is linear then (3.1) is called a **linear difference equation**. For the special case when the coefficients $g_1, \ldots, g_k$ are constants, (3.1) becomes

$$s_n = c_1 s_{n-1} + \ldots + c_k s_{n-k} + \Psi(n) \quad \text{for} \quad c_1, \ldots, c_k \in \mathbb{C},$$

and is called a **constant coefficient difference equation**. The solution of an initial value problem for a linear constant coefficient difference equation is similar to that for a linear constant coefficient differential equation, where a sum plays the role of the integral. The theory for difference equations is much simpler. For instance, we've already seen that uniqueness is immediate in the case of difference equations, whereas for differential equations uniqueness is not guaranteed for all functions.

In this chapter we're interested in linear constant coefficient difference equations, and we simply refer to them as difference equations or recurrence relations. Every such relation can be written in the form

$$\text{(L)} \qquad s_n - c_1 s_{n-1} - c_2 s_{n-2} - \cdots - c_k s_{n-k} = \psi(n) \text{ for } n \geq k\,,$$

for some complex-valued function $\psi$ and constants $c_1, \ldots, c_k$ where $c_k$ is non-zero. The *homogeneous* linear constant coefficient equation (HL) studied in Chapter 2 is the special case when the input is zero.

### *3.1.1 First-Order Equations*

A **first-order recurrence** has the form

$$\text{(L1)} \qquad s_n = \lambda s_{n-1} + \psi(n) \text{ for all } n \geq 1\,,$$

where the function $\psi$ is defined for all positive natural numbers and $\lambda \neq 0$. When we consider the associated initial value problem for fixed $s_0 = \alpha_0$, we have

$$\begin{aligned}
s_0 &= \alpha_0 \\
s_1 &= \lambda s_0 + \psi(1) = \lambda\alpha_0 + \psi(1)\,; \\
s_2 &= \lambda s_1 + \psi(2) = \lambda^2\alpha_0 + \lambda\psi(1) + \psi(2)\,; \\
s_3 &= \lambda s_2 + \psi(3) = \lambda^3\alpha_0 + \lambda^2\psi(1) + \lambda\psi(2) + \psi(3) \\
&\vdots
\end{aligned}$$

and this gives the following form for the solution.

**Theorem 3.1.2.** *The initial value problem*

$$s_n = \lambda s_{n-1} + \psi(n), \quad s_0 = \alpha_0 \tag{3.2}$$

*always has the unique solution* $< s_n >$ *with general term*

$$s_n = \alpha_0 \lambda^n + \sum_{i=1}^{n} \psi(i) \lambda^{n-i} \tag{3.3}$$

*(where the sum is defined to be zero when* $n = 0$*.)*

*Proof.* Since we already know that (3.2) has a unique solution, we will show by induction on $n$ that the solution has the form given in (3.3).

For $n = 0$, observe that (3.3) reduces to $s_0 = \alpha_0$, which is the initial condition. For the induction step, assume that (3.3) does satisfy (3.2) for all $n = 0, \ldots, K-1$ and then use the defining equation (3.2) to find $s_K$ from $s_{K-1}$ and $\psi(K)$

$$\begin{aligned} s_K &= \lambda s_{K-1} + \psi(K) \\ &= \lambda \left( \alpha_0 \lambda^{K-1} + \sum_{i=1}^{K-1} \psi(i) \lambda^{K-1-i} \right) + \psi(K) \\ &= \alpha_0 \lambda^K + \sum_{i=1}^{K} \psi(i) \lambda^{K-i}, \end{aligned}$$

as required. □

This proof is remarkably simple when compared to the corresponding theorem for differential equations. To solve a first-order initial value problem in differential equations,

$$cs(t) = D(s(t)) + \psi(t), \quad s(0) = \alpha_0,$$

it's necessary to multiply through by an "integrating factor" $e^{ct}$ so that the equation becomes

$$D\left(e^{ct} s(t)\right) = e^{ct} \psi(t),$$

which has the solution

$$s(t) = e^{-ct} \left( \int_0^t e^{ct} \psi(t)\, dt \right) + \alpha_0$$

provided the integral exists. From this we see several ways in which differential equations are more difficult: the existence of a solution is not guaranteed, finding a solution is harder, and some theory is needed to deal with uniqueness.

Let's look at the simplest example of a nonhomogeneous first-order recurrence, when the input $\psi(n) = c$ is a constant,

$$s_n = \lambda s_{n-1} + c, \quad s_0 = \alpha_0 \,. \tag{3.4}$$

From (3.3) we have

$$s_n = \alpha_0 \lambda^n + c(\lambda^{n-1} + \lambda^{n-2} + \cdots + 1),$$

which can be computed to be

$$s_n = \begin{cases} \alpha_0 \lambda^n + c\dfrac{1-\lambda^n}{1-\lambda} & \text{if } \lambda \neq 1 \\ \alpha_0 + cn & \text{if } \lambda = 1 \end{cases} . \tag{3.5}$$

### *3.1.2 Finding a particular solution via convolution*

We have just proved that the sequence $< v_n >$ defined by

$$v_0 = 0\,;\; v_n = \sum_{i=1}^{n} \psi(i)\lambda^{n-i} = (\psi(1), \ldots, \psi(n)) \cdot (\lambda^{n-1}, \ldots, \lambda, 1)$$

(where the last expression should be read as a convolution, or dot product) is a particular solution to the first-order equation

$$s_n = \lambda s_{n-1} + \psi(n)$$

and the solution can be rewritten as the dot product

$$v_n = (\psi(1), \ldots, \psi(n)) \cdot (t_{n-1}, \ldots, t_0) \,.$$

where $t_n = \lambda^n$ is the solution to the homogeneous initial value problem

$$s_n = \lambda s_n \quad \text{with} \quad s_0 = 1 \,.$$

We can generalize this formula to general $k^{\text{th}}$-order difference equations.

**Theorem 3.1.3.** *Let $< t_n >$ be the solution to the $k^{th}$-order homogeneous initial value problem*

$$s_n = c_1 s_{n-1} + c_2 s_{n-2} + \ldots + c_k s_{n-k} \quad \textit{with} \quad (s_{k-1}, \ldots, s_0) = (1, 0, \ldots, 0) \,.$$

*Then for any complex sequence $\psi(n)$, the sequence $< v_n >$ given by*

$$v_0 = \ldots = v_{k-1} = 0 \quad , \quad v_n = (\psi(k), \ldots, \psi(n)) \cdot (t_{n-1}, \ldots, t_{k-1})$$

*is a particular solution to the difference equation*

$$s_n = c_1 s_{n-1} + c_2 s_{n-2} + \ldots + c_k s_{n-k} + \psi(n) \,. \tag{3.6}$$

*Proof.* Since $v_0 = \ldots = v_{k-1} = 0$, the value $v_k = \psi(k)t_{k-1} = \psi(k)$ does satisfy (3.6)for $n = k$. For $n > k$, since $t_0 = \ldots = t_{k-2} = 0$ then $c_1 v_{n-1} + c_2 v_{n-2} + \ldots + c_k v_{n-k}$ equals

$$\begin{aligned} &c_1(t_{n-2}\psi(k) + \ldots + \ t_k\ \psi(n-2)\ + t_{k-1}\psi(n-1)) \\ &c_2(t_{n-3}\psi(k) + \ldots + t_{k-1}\psi(n-2) + \ t_{k-2}\psi(n-1)) \\ &\qquad\qquad + \ldots + \\ &c_k\,(t_{n-1-k}\psi(k) + \ldots + \ t_1\ \psi(n-2)\ + \ \ t_0\,\psi(n-1)) \end{aligned}$$

Adding down the "columns", this becomes

$$\begin{aligned} &\Big(\sum_{i=1}^{k} c_i t_{n-1-i}\Big)\psi(k) + \ldots + \Big(\sum_{i=1}^{k} c_i t_{k-i}\Big)\psi(n-1) \\ &\quad = t_{n-1}\psi(k) + \ldots + t_k\psi(n-1) \\ &\quad = (t_{n-1}, \ldots, t_k, 1)\cdot(\psi(k), \ldots, \psi(n-1), \psi(n)) - \psi(n) \\ &\quad = v_n - \psi(n) \end{aligned}$$

by definition of $v_n$. By induction, $< v_n >$ *is* a solution to

$$v_n = c_1 v_{n-1} + \ldots + c_k v_{n-k} + \psi(n)\,.$$

□

Let's find a particular solution to the second-order equation

$$s_n = 2s_{n-1} - s_{n-2} + 2^n\,. \tag{3.7}$$

As in the theorem, letting $< t_n >$ be the solution to the associated homogeneous initial value problem

$$s_n = 2s_{n-1} - s_{n-2} \quad ; \quad (s_1, s_0) = (1, 0)\,,$$

its first few terms are $t_0 = 0, t_1 = 1, t_2 = 2, t_3 = 3, \ldots$ and $t_n = n$ can be proved in general by induction. Then

$$v_n = (\psi(2), \ldots, \psi(n))\cdot(n-1, \ldots, 1) = (2^2, \ldots, 2^{n-1}, 2^n)(n-1, \ldots, 2, 1)$$

is a particular solution of (3.7). In this case it's relatively easy to get a closed form for $< v_n >$ because the sequence satisfies a first-order equation:

$$\begin{aligned} v_n &= (2^2, \ldots, 2^n)\cdot(1, \ldots, 1) + (2^2, \ldots, 2^{n-1}, 2^n)\cdot(n-2, \ldots, 1, 0) \\ &= 2^2(1 + \ldots + 2^{n-2}) + v_{n-1} = 2^2(2^{n-1} - 1) + v_{n-1}\,. \end{aligned}$$

Therefore, using $\lambda = 1$ and $\Psi(n) = 4(2^{n-1} - 1)$, $< v_n >$ is a solution to $v_n = \lambda v_{n-1} + \Psi(n)$ and from the result for first-order equations

$$\begin{aligned} v_n &= (\Psi(1), \ldots, \Psi(n))\cdot(\lambda^{n-1}, \ldots, \lambda) \\ &= 4\sum_{i=0}^{n-1}(2^i - 1) = 4(2^n - 1) - 4n = 4(2^n - 1 - n) \end{aligned}$$

is a particular solution to (3.7). Notice that determining this closed form relied on being able to compute $t_n$ easily as well as on seeing a linear relationship between consecutive elements of the constructed particular solution $< v_n >$.

## 3.2 A Special Class of Linear Recurrences

We now generalize the last section and develop an efficient method for obtaining a particular solution for a frequently occurring class of linear recurrences, the ones whose forcing function is $\psi(n) = \lambda^n p(n)$ where $\lambda \in \mathbb{C}$ and $p(x)$ is a polynomial.

**Theorem 3.2.1.** *For any polynomial $p(x)$ and any constant $\lambda$, consider the linear recurrence*

$$s_n = c_1 s_{n-1} + \ldots + c_k s_{n-k} + \lambda^n p(n)\,.$$

*Let $\lambda_1, \ldots, \lambda_t$ be the eigenvalues of the recurrence with respective multiplicities $m_1, \ldots, m_t$ and define $\delta$ by*

$$\delta = \begin{cases} 0 & \text{if } \lambda \notin \{\lambda_1, \ldots, \lambda_t\} \\ m_i & \text{if } \lambda = \lambda_i \end{cases}\,.$$

*If $\deg(p) = d$ then there exists a polynomial $q(x)$ with $\deg(q) \leq d$ such that $v_n = \lambda^n n^\delta q(n)$ is a particular solution of the recurrence. Moreover, the polynomial $q(x)$ can be calculated in $O(d^2)$ operations.*

*Proof.* Our proof actually constructs the polynomial $q$. Since the equation is homogeneous when $\lambda$ is zero, we may assume that $\lambda \neq 0$. We note that $\lambda^n n^\delta q(n)$ is a solution iff for all integers $n \geq 0$,

$$\begin{aligned}&\lambda^n n^\delta q(n) - c_1 \lambda^{n-1}(n-1)^\delta q(n-1) \\ &\quad - c_2 \lambda^{n-2}(n-2)^\delta q(n-2) - \ldots - c_k \lambda^{n-k}(n-k)^\delta q(n-k) = \lambda^n p(n)\,.\end{aligned}$$

Dividing both sides by the non-zero $\lambda^{n-k}$, the equation becomes a polynomial equation in $n$ which holds on the (infinite) set of natural numbers and so in fact the equation is the polynomial identity

$$\lambda^k x^\delta q(x) - c_1 \lambda^{k-1}(x-1)^\delta q(x-1) - \ldots - c_k (x-k)^\delta q(x-k) = \lambda^k p(x)\,.$$

Remember that $p(x) = p_d x^d + \ldots + p_1 x + p_0$ is a fixed polynomial and our goal is to construct coefficients $q_d, \ldots, q_1, q_0$ such that $q(x) = q_d x^d + \ldots +$

$q_1x + q_0$ satisfies this identity. Computing a few coefficients, the coefficient of $x^{d+\delta}$ on the left side is

$$\lambda^k q_d - c_1\lambda^{k-1}q_d - \ldots - c_k q_d$$

which is $ch(\lambda)q_d$, and the coefficient of $x^{\delta+d-1}$ is

$$\lambda^k q_{d-1} - c_1\lambda^{k-1}\left((-1)\binom{d+\delta}{1}q_d + q_{d-1}\right)$$
$$- c_2\lambda^{k-2}\left((-2)\binom{d+\delta}{1}q_d + q_{d-1}\right) - \ldots - c_k\left((-k)\binom{d+\delta}{1}q_d + q_{d-1}\right),$$

which is $-t_1\binom{d+\delta}{1}q_d + t_0 q_{d-1}$ for $t_0 = ch(\lambda)$ and $t_1 = \sum_{j=1}^{k} c_j(-j)\lambda^{k-j}$. In general, defining $t_i = \sum_{j=1}^{k} c_j(-j)^i\lambda^{k-j}$ for all $i = 0, 1, \ldots, d$, it can be shown that the coefficient of $x^{\delta+d-j}$ on the left side is

$$-t_j\binom{\delta+d}{j}q_d - \ldots - t_1\binom{\delta+d-j-1}{1}q_{d-j-1} + t_0 q_{d-j}\,.$$

First considering the case when $\lambda$ is not an eigenvalue, then $t_0 = ch(\lambda)$ is non-zero, $\delta = 0$, and the requirement becomes the dot product

$$\left(-t_j\binom{d}{j}, \ldots, -t_1\binom{d-j-1}{1}, t_0\right)\cdot(q_d, \ldots, q_{d-j-1}, q_{d-j}) = \lambda^k p_j\,,$$

for all $j = 0, \ldots, d$, a system of $d+1$ equations in the unknowns $q_d, \ldots, q_1, q_0$. The coefficient matrix is a lower triangular matrix in which each diagonal entry is the non-zero $t_0$, and this means the system has a unique solution which gives a unique polynomial $q(x)$ whose degree is at most $d$.

If $\lambda$ is a simple eigenvalue, then $\delta = 1$ and $t_0 = 0$ which means that the coefficient of $x^{d+\delta}$ on the left side is $q_d t_0 = 0$ and $q_d = 1$ can be used. The remaining restrictions are given by the dot product

$$\left(-t_j\binom{1+d}{j}, \ldots, -t_1\binom{1+d-j-1}{1}\right)\cdot(q_d, \ldots, q_{d-j-1}) = \lambda^k p_j$$

for all $j = 1, \ldots, d$, again a lower triangular system in which each diagonal entry is now $t_1$. Fortunately,

$$k\,ch(x) - xch'(x) = \sum_{j=1}^{k} c_j(-j)x^{k-j}$$

which means that

$$k\,ch(\lambda) - \lambda ch'(\lambda) = t_1\,.$$

Since $\lambda$ is a *simple* eigenvalue, then $ch(\lambda) = 0$ but $ch'(\lambda) \neq 0$, and so $t_1 \neq 0$ as required.

What about the remaining cases, when the multiplicity of $\lambda$ is $m > 1$? Then $ch(\lambda) = 0$ and for each $j < m$ the value of $D^{(j)}(ch)$ at $x = \lambda$ is zero, which means that the first $\delta = m$ coefficients, $q_d, q_{d-1}, \ldots, q_{d-m+1}$ can be chosen to be zero. But $t_m$ can be written as a linear combination of $ch(\lambda), D(ch)(\lambda), \ldots, D^{(m)}(ch)(\lambda)$ (refer to Exercise 3.6), and $t_m \neq 0$ which means that the upper triangular system yields unique values for the coefficients $q_{d-m}, \ldots, q_0$.

Now for the operation count. Since $k$ is a constant, each of the $O(d^2)$ entries in the coefficient matrix can be computed with a constant number of multiplications. Further, because the matrix is in triangular form, the associated system of equations can be solved by **back substitution**. That is, $q_d$ can be computed in one division; $q_{d-1}$ is computed using this known value of $q_d$ and one division; in general, each $q_{d-j}$ is computed from the values $q_d, q_{d-1}, \ldots, q_{d-j-1}$ using $O(j)$ operations. This means that, once the coefficient matrix has been computed, the coefficient set $q_d, \ldots, q_0$ can be found using $O(d^2)$ operations. □

For the second-order recurrence considered earlier,

$$s_n = 2s_{n-1} - s_{n-2} + 2^n\,,$$

$\lambda = 2$ is not an eigenvalue of the recurrence, which means $\delta = 0$ and $\deg(q) = 0$ follows from $p(x) = 1$. This gives the particular solution $v_n = 2^n c$ where $c$ is a constant. To solve for $c$,

$$v_n - 2v_{n-1} + v_{n-2} = 2^n c - 2^n c + 2^{n-2} c = 2^{n-2} c\,,$$

giving $c = 4$ and $v_n = 2^{n+2}$, which is a different solution from the one we found earlier. Appendix A contains more worked problems which use this technique.

## 3.3 The Shift Operator on the Space of Sequences

In contrast with the homogeneous recurrences in Chapter 2, when $\psi$ is not the zero function the set of solutions to (L) is not a vector space, but the problem of solving these nonhomogeneous recurrences can be put into a linear algebra context by looking at things slightly differently. For the moment we consider the infinite-dimensional complex space $\mathcal{S}$ of *doubly infinite* sequences

$$\ldots, s_{-2}, s_{-1}, s_0, s_1, s_2, \ldots\,,$$

and define the **shift operator** $\sigma$ on $\mathcal{S}$ to be the function which shifts a sequence $s \in \mathcal{S}$ one step to the right,

$$\sigma(< s_n >) =< s_{n-1} > \ .$$

This operator is defined on the vector space $\mathcal{S}$. We'd like a right-shift operator on $\mathcal{S}^+$, the space of singly infinite sequences, but how is it defined? That is, when we shift a *singly* infinite sequence one step to the right, what fills the vacated $0^{\text{th}}$ term? In Exercise 3.1 you show that there's only one assignment which results in a *linear* operator: the $0^{\text{th}}$ term must always equal 0. Accordingly, the linear **shift operator** $\sigma$ on $\mathcal{S}^+$ is defined as

$$\sigma(< s_0, s_1, s_2, s_3, \cdots >) =< 0, s_0, s_1, s_2, s_3, \cdots > . \tag{3.8}$$

(Here we're using the same symbol for the shift operator on both $S$ and $S^+$.) In Chapter 2 powers of the differentiation operator $D$ denoted repeated differentiation (repeated composition of the operator) and powers of the differentiation operator are linear. In general, integer powers of a linear operator are linear, and in fact any polynomial in a linear operator is a linear operator. For example, because the operator $\sigma^k$ shifts a sequence $k$ steps to the right, applying the operator $2 - 3\sigma + 4\sigma^3$ to any sequence $< s_n >$ results in a sequence whose $n^{\text{th}}$ term (for $n \geq 3$) is $2s_n - 3s_{n-1} + 4s_{n-3}$. Since the left side of (L) is the $n^{\text{th}}$ term of the sequence $s - c_1\sigma(s) - c_2\sigma^2(s) - \cdots - c_k\sigma^k(s)$, it is precisely the $n^{\text{th}}$ term of the sequence $L(s)$, where $L$ is the linear operator $L = I - c_1\sigma - c_2\sigma^2 - \cdots - c_k\sigma^k$. This representation allows us to see another analogy between difference equations and differential equations. The general $k^{\text{th}}$-order linear constant coefficient differential equation has the form

$$x(t) - c_1 D(x(t)) - c_2 D^2(x(t)) - \cdots - c_k D^k(x(t)) = \beta(t)\,, \tag{3.9}$$

where $D$ is the differentiation operator on the vector space of infinitely differentiable functions on $\mathbb{R}$ or $\mathbb{C}$ (or some other domain of definition). The left-hand side of (3.9) can be written as $L_D(x)$ for

$$L_D = I - c_1 D - c_2 D^2 - \cdots - c_k D^k$$

and (3.9) becomes the functional equation $L_D(x) = \beta$. Likewise, the difference equation (L) can be written as $L(s) = \boldsymbol{\psi}$, where

$$\boldsymbol{\psi} =< 0, 0, \ldots, 0, \psi(k), \psi(k+1), \cdots >$$

is an element of the vector space $\mathcal{S}^+$. We see that $s \in \mathcal{S}^+$ is a solution to (L) if and only if the sequence $L(s) - \boldsymbol{\psi}$ has zeros from the $k^{\text{th}}$ term onwards. We define the $k$-dimensional subspace (refer to Exercise 3.3)

$$V_k = \{ s \in S^+ \,:\, s_n = 0 \text{ for all } n \geq k \},$$

and obtain

$$s \in S^+ \text{ is a solution to (L)} \iff L(s) - \boldsymbol{\psi} \in V_k. \tag{3.10}$$

**Theorem 3.3.1.** *For any choice of constants $c_1, c_2, \ldots, c_k \in \mathbb{C}$, the operator $L = I - c_1\sigma - c_2\sigma^2 - \cdots - c_k\sigma^k$ is invertible on $\mathcal{S}^+$ and the set of preimages $L^{-1}(V_k)$ is a $k$-dimensional subspace of $\mathcal{S}^+$.*

*Proof.* To prove the invertibility of $L$ we show any choice of sequence $y \in \mathcal{S}^+$ has a unique preimage $s \in \mathcal{S}^+$. For this, we define the complex-valued function $\psi$ on $\mathbb{N}$ by $\psi(n) = y_n$, the $n^{\text{th}}$ term of the given sequence $y$, and also let $\boldsymbol{\alpha} = (y_{k-1}, \ldots, y_0)$. Then $L(s) = y$ encodes the initial value problem

$$s_n - c_1 s_{n-1} - c_2 s_{n-2} - \cdots - c_k s_{n-k} = \psi(n)\,;\; S_0^T = \boldsymbol{\alpha},$$

which we know always has a unique solution $s \in \mathcal{S}^+$. This proves there exists a unique $s \in \mathcal{S}^+$ with $L(s) = y$. Since the restriction of $L^{-1}$ to the $k$-dimensional subspace $V_k$ must be invertible, $L^{-1}(V_k)$ is a $k$-dimensional subspace of $\mathcal{S}^+$. □

Therefore, (3.10) becomes

$$s \in S^+ \text{ is a solution to (L)} \iff s \in L^{-1}(\boldsymbol{\psi}) + L^{-1}(V_k). \tag{3.11}$$

Geometrically this means that the set of solutions to (L) forms a $k$-dimensional **hyperplane**, a translate of a $k$-dimensional subspace by a single vector. Algebraically this means that we can find the general solution to (L) by first finding a particular solution to $L(s) = \boldsymbol{\psi}$ and then adding the general solution to the homogeneous equation $L(s) = 0$. Once the general solution is known, any initial value problem can be solved. In general there's no algorithm which gives a particular solution in closed form, and in fact some classes of linear recurrences don't have a "closed form" solution [39, 116].

Let's solve the following initial value problem

$$s_0 = 0\,,\; s_1 = 1\,,\; s_n = 4s_{n-1} - 4s_{n-2} + 3^n(n-1)\,.$$

Then $\psi(n)$ has the form $\psi(n) = \lambda^n p(n)$, where $\lambda = 3$ and $p(x) = x - 1$ is a polynomial of degree $d = 1$. Using Theorem 3.2.1, since $\lambda_1 = 2$ is the only root of $ch(x) = (x-2)^2$ and $\lambda \neq \lambda_1$, then $\delta = 0$, and there is a particular solution of the form

$$v_n = 3^n q(n)\,,$$

where $\deg(q) \leq \deg(p) = 1$. Also, $< v_n >$ can be constructed by the method in the proof of the theorem. For this, $ch(3) = 1$ and $ch'(3) = 2$, which gives

$$t_0 = ch(3) = 1 \text{ and } t_1 = 2ch(3) - 3ch'(3) = -4\,.$$

Since $p_1 = 1$, $p_0 = -1$, the coefficients of $q(x)$ satisfy the associated system of equations

$$\begin{bmatrix} 1 & 0 \\ 4 & 1 \end{bmatrix} \begin{pmatrix} q_1 \\ q_0 \end{pmatrix} = \lambda^2 \begin{pmatrix} 1 \\ -1 \end{pmatrix} = \begin{pmatrix} 9 \\ -9 \end{pmatrix}\,,$$

giving $q_1 = 9$ and $q_0 = -4q_1 - 9 = -45$, which means $v_n = 3^n(9n - 45)$ is a particular solution to the recurrence. Although $< v_n >$ doesn't satisfy the initial conditions, every solution to the recurrence has the form $s_n = h_n + v_n$ where $h_n = (\alpha n + \beta)2^n$ is the general solution to the homogeneous recurrence. From the initial conditions,

$$0 = s_0 = (\alpha \cdot 0 + \beta)2^0 + (9 \cdot 0 - 45)3^0$$
$$1 = s_1 = (\alpha \cdot 1 + \beta)2^1 + (9 \cdot 1 - 45)3^1 ,$$

which together yield $\beta = 45, \alpha = 19/2$ and $s_n = (\frac{19}{2}n + 45)2^n + 3^n(9n - 45)$.

## 3.4 Formal Power Series

We've used linear operators which are polynomials in $\sigma$ and we now go one step further and discuss formal power series in $\sigma$. The adjective "formal" is used because we aren't concerned with convergence of the power series but rather treat it as a purely formal object. The interested reader should consult Ivan Niven's paper [105], a good introduction to the application of formal power series to number theory and combinatorics. In the remainder of this chapter we concentrate on the theory of formal power series which will form the foundation for the next chapter on generating functions.

**Definition 3.4.1.** A **formal power series** in the variable $x$ is an infinite sum

$$\gamma = \sum_{i \geq 0} a_i x^i$$

with coefficients $a_i$ in $\mathbb{C}$. Two formal power series are equal if and only if all corresponding coefficients are equal.

Note that a polynomial in $x$ is a power series whose coefficients are zero after some point, and a formal power series can be thought of as an "infinite" polynomial. This is exactly the point of view taken for the addition and multiplication of formal power series: For $\gamma_1 = \sum_{i \geq 0} a_i x^i$ and $\gamma_2 = \sum_{i \geq 0} b_i x^i$, their sum is

$$\gamma_1 + \gamma_2 = \sum_{i \geq 0} (a_i + b_i) x^i .$$

For multiplication of power series, we also mimic what happens with polynomial multiplication. The product $\gamma_1 \cdot \gamma_2$ has constant term $a_0 b_0$; the coefficient of $x$ in the product

The product of power series defined here is called a **convolution**. Exercises 3.12 to 3.17 is a series of exercises on convolution.

is $a_0b_1 + a_1b_0$; the coefficient of $x^2$ is $a_0b_2 + a_1b_1 + a_2b_0$, and so on. This leads to the general formula for the product of two power series

$$\gamma_1 \cdot \gamma_2 = \Big(\sum_{i\geq 0} a_i x^i\Big)\Big(\sum_{j\geq 0} b_j x^j\Big) = \sum_{n\geq 0}\Big(\sum_{i+j=n} a_i b_j\Big)x^n\,. \tag{3.12}$$

An important point to note is that since there are only $n+1$ pairs of indices $i, j$ which sum to $n$, each coefficient on the right side is a *finite* sum of products $a_ib_j$ in $\mathbb{C}$, and as such is a complex number. Using these definitions for addition and multiplication, formal power series behave as polynomials in many respects and Theorem 3.4.1 below summarizes the algebraic properties of the **integral domain** of formal power series.

Here's a proof technique that we call the Finiteness Argument which explains a good deal about the way formal power series behave algebraically. Given any formal power series $\gamma = \sum a_i x^i$, we define its **partial sums** just as in calculus: for any nonnegative integer $d$, the $d^{\text{th}}$ partial sum is the polynomial $\gamma_d = \sum_{i=0}^{d} a_i x^i$. As an example, we will use partial sums to investigate the computation of coefficients of the power series $\gamma^2$. From the definition of multiplication, the $0^{\text{th}}$ coefficient of $\gamma^2$ is $a_0^2$; the first coefficient is $2a_0a_1$; the second is $a_1^2 + 2a_0a_2$ and so on. Compare this to the sequence of squares of the partial sums:

$$\begin{aligned}
\gamma_0^2 &= a_0^2\\
\gamma_1^2 &= a_0^2 + 2a_0a_1x + a_1^2x^2\\
\gamma_2^2 &= a_0^2 + 2a_0a_1x + (a_1^2 + 2a_0a_2)x^2 + 2a_1a_2x^3 + a_2^2x^4\\
&\vdots
\end{aligned}$$

What happens is that the $0^{\text{th}}$ coefficient of $\gamma^2$ is the same as that of $\gamma_d^2$ for all $d$, the $1^{\text{st}}$ coefficient of $\gamma^2$ is the same as the first coefficient of $\gamma_d^2$ for $d \geq 1$, the $2^{\text{nd}}$ coefficient of $\gamma^2$ is the same as that of $\gamma_d^2$ for $d \geq 2$. This pattern continues, as we will prove below. This is what we refer to as the **Finiteness Argument**. It allows us to reduce any question about finitely many coefficients of a formal power series to a question about the coefficients of one of its partial sums.

For example, the statement

"The $17^{\text{th}}$ coefficient of $\gamma^2$ is positive."

can be checked by testing the $17^{\text{th}}$ coefficient of any $\gamma_d^2$ with $d \geq 17$ and the statement can *not* be checked by looking at the $17^{\text{th}}$ coefficient of $\gamma_{16}^2$. In a similar way, consider a statement such as

"All the coefficients of $\gamma^2$ are positive".

If we can show that for every $d \geq 0$ the first $d$ coefficients of $\gamma_d^2$ are positive, then we could prove this statement as well.

The same ideas apply to more complicated expressions in more than one formal power series. For example, for power series $\gamma, \delta, \epsilon$, the $n^{\text{th}}$ coefficient of $3\delta - 4\gamma\epsilon^4 + \epsilon$ coincides with the $n^{\text{th}}$ coefficient of the polynomial $3\delta_d - 4\gamma_d\epsilon_d^4 + \epsilon_d$ for any $d \geq n$. Writing this more succinctly, for $\Psi(x, y, z) = 3y - 4xz^4 + z$, the $n^{\text{th}}$ coefficient of $\Psi(\gamma, \delta, \epsilon)$ equals the $n^{\text{th}}$ coefficient of $\Psi(\gamma_d, \delta_d, \epsilon_d)$ for all $d \geq n$.

| The Finiteness Argument |
|---|
| *Let $\Psi(x_1, \ldots, x_t)$ be a polynomial in the variables $x_1, \ldots, x_t$.*<br>*Let $\gamma_1, \ldots, \gamma_t$ be formal power series in $x$.*<br>*For each $i$, $(\gamma_i)_d$ will denote the $d^{th}$ partial sum of $\gamma_i$.*<br>*Then for each $n$, the $n^{th}$ coefficient of $\Psi(\gamma_1, \ldots, \gamma_t)$ coincides with the $n^{th}$ coefficient of the polynomial $\Psi((\gamma_1)_d, \ldots, (\gamma_t)_d)$ for all $d \geq n$.* |

The idea behind the proof of the Finiteness Argument is the following: If we first consider the most basic polynomials

$$\Psi_1(x_1, x_2) = x_1 + x_2 \quad \text{and} \quad \Psi_2(x_1, x_2) = x_1 x_2$$

and two power series $\gamma_1, \gamma_2$, the conclusion of the Finiteness Argument for each of $\Psi_1(\gamma_1, \gamma_2)$, $\Psi_2(\gamma_1, \gamma_2)$ follows from the respective definitions of addition and multiplication for power series. Since the arithmetic operations inherent in the polynomial $\Psi$ form a *finite* sequence of additions and multiplications of pairs of formal power series, the result follows by induction on the number of operations used to build up to $\Psi(\gamma_1, \ldots, \gamma_t)$.

The Finiteness Argument is a useful way to see that algebraic properties of polynomials transfer to the same properties for formal power series.

**Theorem 3.4.1.** *The set of formal power series in $x$ (with coefficients in $\mathbb{R}$ or $\mathbb{C}$) forms an* **integral domain** *under the operations of addition and multiplication. This means that the following properties hold:*

*(a) The operations of addition and multiplication satisfy the associative and commutative laws, and also satisfy the usual distributive law of multiplication over addition.*
*(b) The constant polynomial 0 is the additive identity.*
*(c) Every power series has an additive inverse, obtained by negating all of its coefficients.*
*(d) The constant polynomial 1 is the multiplicative identity.*
*(e) Whenever the product $\gamma_1 \cdot \gamma_2$ equals the zero power series at least one of the factors $\gamma_i$ must equal zero.*

*Proof.* We'll prove that the distributive law holds and then just say that all other properties can be proved in the same manner. Let $\gamma$, $\delta$, and $\epsilon$ be three arbitrary formal power series in $x$. The distributive law says that

$\gamma(\delta+\epsilon) = \gamma\delta + \gamma\epsilon$. This is an infinite string of statements asserting an equality between the $n^{\text{th}}$ coefficient of $\gamma(\delta+\epsilon)$ and the $n^{\text{th}}$ coefficient of $\gamma\delta + \gamma\epsilon$. Fix any $n \geq 0$. Since polynomial multiplication distributes over polynomial addition, for every $d \geq n$ we have the partial sum arithmetic $\gamma_d(\delta_d + \epsilon_d) = \gamma_d\delta_d + \gamma_d\epsilon_d$. Therefore, the Finiteness Argument implies that $\gamma(\delta+\epsilon) = \gamma\delta + \gamma\epsilon$, proving the distributive law does hold for power series. □

We use the modifier *finiteness* because the argument reduces a question about the infinitely many coefficients in a formal power series to the analogous question for its partial sums which are polynomials or finite power series. There's a more sophisticated finiteness argument which involves the substitution of polynomials into power series. For example, if we substitute the polynomial $x^2$ into the power series

$$(1 + 3x + 5x^2 + \cdots) = \sum_{i\geq 0}(2i+1)x^i\,,$$

we would expect to get the power series

$$(1 + 3x^2 + 5x^4 + \cdots) = \sum_{i\geq 1}(2i+1)x^{2i}\,.$$

Substituting the polynomial $x + x^2$ into the same power series should yield

$$1 + 3(x+x^2) + 5(x+x^2)^2 + 7(x+x^2)^3 + \cdots \\ = 1 + 3x + 8x^2 + 17x^3 + 35x^4 + \cdots\,.$$

This type of substitution works because the coefficient of the $x^n$ term in the new power series is a *finite* sum of monomials.

More precisely, consider any power series $\gamma = \sum_{i\geq 0} a_i x^i$ and a polynomial $P(x)$ with *zero* constant term. Because the constant term of $P(x)$ is zero, the exponent of every term in $P(x)^i$ is at least $i$. Thus, in the expansion of

$$\gamma(P(x)) = a_0 + a_1P(x) + a_2P(x)^2 + \cdots\,,$$

no summand $a_iP(x)^i$ with $i > n$ can contribute to the $x^n$ term. This shows two things: that the $x^n$ term in $\gamma(P(x))$ is a finite sum of monomials (proving that $\gamma(P(x))$ is a well defined formal power series) and that the $n^{\text{th}}$ term of $\gamma(P(x))$ is the same as the $n^{\text{th}}$ term of the polynomial $\gamma_d(P(x))$ for any $d \geq n$. This allows the following extension of the Finiteness Argument:

The Extended Finiteness Argument

*Let $\Psi(x_1, \ldots, x_t)$ be a polynomial in the variables $x_1, \ldots, x_t$.*
*Let $\gamma_1, \ldots, \gamma_t$ be formal power series in $x$.*
*Let $P_1(x), \ldots, P_t(x)$ be polynomials with zero constant term.*
*Then*

$$\Psi(\gamma_1(P_1), \gamma_2(P_2), \cdots, \gamma_t(P_t))$$

*is a well defined power series whose $n^{th}$ term coincides with the $n^{th}$ term of the polynomial*

$$\Psi((\gamma_1)_d(P_1), (\gamma_2)_d(P_2), \cdots, (\gamma_t)_d(P_t)), \text{ for all } d \geq n.$$

In Theorem 3.4.1 the set of formal powers series was shown to be an integral domain, so called because it has the same algebraic properties as the integers. Unlike what happens in $\mathbb{Z}$, the next result shows there are many power series which have multiplicative inverses.

**Theorem 3.4.2.** *Let $P(x)$ be a polynomial in $x$ with zero constant term. Then $\Gamma = \sum_{i \geq 0} P(x)^i$ is a well defined formal power series and $\Gamma$ is the multiplicative inverse of $1 - P(x)$ in the integral domain of formal power series.*

*Proof.* We set $\Psi(x_1, x_2) = x_1 x_2$ and

$$\gamma_1 = 1 - x \quad ; \quad \gamma_2 = \sum_{i \geq 0} x^i \quad ; \quad P_1(x) = P_2(x) = P(x)\,.$$

Then for each $d \geq 1$, the $d^{\text{th}}$ partial sums of $\gamma_1$ and $\gamma_2$ are $(\gamma_1)_d = 1 - x$ and $(\gamma_2)_d = 1 + x + \ldots + x^d$. Polynomial multiplication gives

$$(\gamma_1)_d \cdot (\gamma_2)_d = 1 - x^{d+1}$$

and

$$\Psi((\gamma_1)_d(P), (\gamma_2)_d(P)) = 1 - P^{d+1}\,.$$

Since the $0^{\text{th}}$ and $d^{\text{th}}$ terms of $\Psi((\gamma_1)_d(P), (\gamma_2)_d(P))$ are therefore 1 and 0 respectively, from the Extended Finiteness Argument we have

$$1 = \Psi(\gamma_1(P), \gamma_2(P)) = (1 - P(x)) \sum_{i \geq 0} P(x)^i,$$

as required. □

Using $P(x) = \lambda x$ in the last result we have

**Theorem 3.4.3 (The Geometric Power Series).** *For non-zero $\lambda \in \mathbb{C}$, the inverse of $1 - \lambda x$ is*

$$\frac{1}{1-\lambda x} = \sum_{n\geq 0} \lambda^n x^n \,. \tag{3.13}$$

When $\lambda$ is restricted to be a real number and $x$ satisfies $|\lambda x| < 1$, Theorem 3.4.3 becomes the formula for the geometric series from calculus. In calculus it's a statement about a function defined within its radius of convergence $|x| < |\lambda|^{-1}$, and (3.13) has no meaning outside its radius of convergence. For us $\sum_{n\geq 0} \lambda^n x^n$ is a purely formal object.

### *3.4.1 Formal differentiation*

We next use the process of formal differentiation to develop the power series for rational functions of the form $(1-\lambda x)^{-m}$. They will be used in the next chapter on generating functions.

The **formal derivative** of a power series $\gamma = \sum_{i\geq 0} a_i x^i$ is defined to be the formal power series

$$\mathcal{D}(\gamma) = \sum_{i\geq 1} i a_i x^{i-1}. \tag{3.14}$$

The usual rules for differentiation, such as the sum rule and the product rule, can be verified for formal derivatives. Using the definition of convolution,

$$\begin{aligned}\Big(\frac{1}{1-x}\Big)^2 = \Big(\sum_{k\geq 0} x^k\Big)^2 &= \sum_{n\geq 0} \Big(\sum_{i+j=n} 1^2\, x^n\Big) \\ &= \sum_{n\geq 0} (n+1) x^n = \mathcal{D}\Big(\sum_{n\geq 0} x^n\Big) = \mathcal{D}\Big(\frac{1}{1-x}\Big)\,,\end{aligned}$$

and induction gives (refer to Exercise 3.27)

$$\mathcal{D}^j\Big(\sum_{n\geq 0} x^n\Big) = \frac{j!}{(1-x)^{j+1}} \tag{3.15}$$

where $0! = 1$ and $\mathcal{D}^0$ is the identity operator. Letting $\gamma = \sum_{n\geq 0} x^n$,

$$\frac{(m-1)!}{(1-x)^m} = \mathcal{D}^{m-1}(\gamma) = (m-1)! \sum_{n\geq 0} \binom{n+m-1}{n} x^n \,,$$

and we have

$$\frac{1}{(1-\lambda x)^m} = \sum_{n\geq 0} \binom{n+m-1}{n} (\lambda x)^n \,, \tag{3.16}$$

the power series for $(1-\lambda x)^{-m}$ when $m$ is a positive integer. This is a type of Binomial Theorem, because if we define a **generalized binomial coefficient** $\binom{r}{n}$ for any rational $r$ and natural number $n$ by

$$\binom{r}{n} = \frac{r(r-1)\cdots(r-n+1)}{n!} \,,$$

then for negative integer $r = -m$ we have

$$\binom{-m}{n} = (-1)^n \binom{m+n-1}{n}$$

and (3.16) becomes

$$(1+x)^{-m} = \sum_{n\geq 0} \binom{-m}{n} (-1)^n (-x)^n = \sum_{n\geq 0} \binom{-m}{n} x^n \,.$$

In [105], the **Generalized Binomial Theorem**

$$(1+x)^r = \sum_{n\geq 0} \binom{r}{n} x^n \tag{3.17}$$

is proved for all rational $r$.

As often happens in mathematics, a good idea like formal power series can be used in a wide variety of areas. Formal power series with real coefficients are widely used in combinatorics (for example, refer to [134, 135]), and formal power series with complex coefficients have applications in physics and statistics. For these applications as well as for technical reasons, the definition of convolution for power series with complex coefficients can be defined slightly differently. For other applications, such as the theory of automata and formal languages (refer to [124]), formal power series are further generalized to allow the coefficents to lie in a ring or semi-ring rather than in a field.

### *3.4.2 An application of Formal Power Series*

We now return to the question of inverting the operator $L = I - c_1\sigma - c_2\sigma^2 - \cdots - c_k\sigma^k$ where $\sigma$ is the shift operator $\sigma(< s_n >) =< 0, s_0, s_1, \ldots >$ on $\mathcal{S}^+$, the space of infinite sequences of complex numbers. In order to make sense of this, for any choice of $a_i \in \mathbb{C}$ we need to define the formal power series $\Gamma = \sum a_i\sigma^i$ in such a way that it is a linear operator on $\mathcal{S}^+$. For

$s \in \mathcal{S}^+$, it would be natural to interpret $\Gamma(s)$ as an infinite formal sum of the sequences

$$a_0 s + a_1 \sigma(s) + a_2 \sigma^2(s) + \cdots .$$

Writing these summand sequences as rows of the infinite array

$$\begin{array}{ccccccccc} a_0 s_0 & , & a_0 s_1 & , & a_0 s_2 & , & a_0 s_3 & , & \cdots \\ 0 & , & a_1 s_0 & , & a_1 s_1 & , & a_1 s_2 & , & \cdots \\ 0 & , & 0 & , & a_2 s_0 & , & a_2 s_1 & , & \cdots \\ \vdots & & \vdots & & \vdots & & \vdots & & \cdots \end{array} ,$$

because each column has only finitely many non-zero terms the sum of each column is a complex number. This is the motivation for defining $\Gamma(s)$ to be the sequence whose $n^{\text{th}}$ term is

$$a_0 s_n + a_1 s_{n-1} + \cdots + a_n s_0 = (a_0, a_1, \cdots, a_n) \cdot (s_n, s_{n-1}, \cdots, s_0) .$$

In Exercise 3.26 you prove that the function $\Gamma$ defined in this way is a linear operator on $\mathcal{S}^+$.

Next suppose the operators $\Gamma_1 = \sum a_i \sigma^i$, $\Gamma_2 = \sum b_i \sigma^i$ are equal on $\mathcal{S}^+$, that is, $\Gamma_1(s) = \Gamma_2(s)$ for every $s \in \mathcal{S}^+$. This is equivalent to requiring $\Gamma_1 - \Gamma_2$ to be the zero operator on $\mathcal{S}^+$, and in particular this must be true for the sequence $s^*$ whose only non-zero term is its $n^{\text{th}}$ term, with $s^*_n = 1$. Since the $n^{\text{th}}$ term of $(\Gamma_1 - \Gamma_2)(s^*)$ is $a_n - b_n = 0$, then $\Gamma_1$ and $\Gamma_2$ must be equal as power series. This means that algebraic identities involving formal power series are also valid as identities for the linear operators they represent. For example, if $\Gamma_1 \Gamma_2 = 1$ holds in the ring of formal power series then $\Gamma_1 \Gamma_2 = I$ holds as linear operators, and $\Gamma_2$ is the multiplicative inverse of $\Gamma_1$.

As an example, let's use the shift operator to solve the first-order recurrence

$$s_{n+1} = \lambda s_n + n^2 \text{ with } s_0 = \alpha_0 . \tag{3.18}$$

We first shift indices to get $s_n - \lambda s_{n-1} = (n-1)^2$ and rewrite the recurrence as $L(s) = \boldsymbol{\psi}$ where $\boldsymbol{\psi} =< \alpha_0, 0, 1, 4, 9, \cdots >$ and $L = I - \lambda\sigma$. Applying Theorem 3.4.2 with $P(x) = \lambda x$,

$$L^{-1} = I + \lambda\sigma + \lambda^2\sigma^2 + \lambda^3\sigma^3 + \cdots = \sum_{i \geq 0} \lambda^i \sigma^i$$

and the solution is $s = L^{-1}(\boldsymbol{\psi})$ whose $n^{\text{th}}$ term is

$$s_n = (\alpha_0, 0, 1, 4, \cdots, (n-1)^2) \cdot (\lambda^n, \lambda^{n-1}, \cdots, \lambda, 1) .$$

Therefore, the solution to (3.18) is

$$s_n = \alpha_0 \lambda^n + \sum_{i=0}^{n-2} (n-1-i)^2 \lambda^i .$$

(Compare this with Theorem 3.1.2 and our analysis in Section 3.1.2.)

In this example, finding the inverse of $L$ was relatively easy. For higher order recurrences the situation is more difficult. For instance, in the second-order case we must consider the linear operator

$$L = I - c_1\sigma - c_2\sigma^2$$

and Theorem 3.4.2 gives

$$L^{-1} = \sum_{i\geq 0}(c_1\sigma + c_2\sigma^2)^i = I + (c_1\sigma + c_2\sigma^2) + (c_1\sigma + c_2\sigma^2)^2 + \cdots .$$

Setting $L^{-1} = \sum_{i\geq 0} a_i\sigma^i$ we find

$$\begin{aligned} a_0 &= 1 \\ a_1 &= c_1 \\ a_2 &= c_1^2 + c_2 \\ a_3 &= c_1^3 + 2c_1c_2 \\ a_4 &= c_1^4 + 3c_1^2c_2 + c_2^2 \\ a_5 &= c_1^5 + 4c_1^3c_2 + 4c_1c_2^2 \\ &\vdots \end{aligned}$$

The pattern among the $a_i$ seems to be $a_0 = 1,\ a_1 = c_1$ and $a_{i+2} = c_1a_{i+1} + c_2a_i$. The techniques in the next chapter will show that the coefficients of $L^{-1}$ *do* always satisfy the original recurrence. (Refer to Exercise 4.9.)

## 3.5 Exercises

**Ex 3.1.** Let $\gamma : \mathcal{S}^+ \to \mathcal{S}^+$ be defined by

$$\gamma(s_0, s_1, s_2, \ldots) = (a, s_0, s_1, s_2 \ldots) \text{ for some } a \in \mathbb{C}.$$

Show $\gamma$ is linear iff $a = 0$.

**Ex 3.2.** Consider the (right) shift operator on $\mathcal{S}$, $\sigma(< s_n >) =< 0, s_0, s_1, \ldots >$.

(a) Verify that $\sigma$ is a linear transformation on $\mathcal{S}$.
(b) Prove that the left shift operator is an inverse for $\sigma$, and therefore $\sigma$ is an invertible operator on $\mathcal{S}$.

**Ex 3.3.** Show that the space of singly infinite sequences, $\mathcal{S}^+$, is infinite-dimensional by finding $n$ linearly independent vectors for every $n \geq 1$. Verify that for each $k$, $V_k = \{s \in S^+ : s_n = 0 \text{ for all } n \geq k\}$ is a $k$-dimensional subspace of $\mathcal{S}^+$.

**Ex 3.4.** Show that the Fibonacci sequence $< f_n >$ and the shifted sequence $< f_{n-1} >$ are linearly independent. This means that every solution to the Fibonacci recurrence $s_n = s_{n-1} + s_{n-2}$ can be written as $\alpha f_{n-1} + \beta f_n$ where the scalars $\alpha$ and $\beta$ depend on the initial conditions. In particular, find a formula for the **Lucas numbers** (which satisfy the Fibonacci recurrence with $L_0 = 2$, $L_1 = 1$) in the form $L_n = \alpha f_{n-1} + \beta f_n$. Are there integer sequence solutions to $s_n = s_{n-1} + s_{n-2}$ for which $\alpha$ and $\beta$ are not integers?

**Ex 3.5.** For this exercise, define a **full solution** to a $k^{\text{th}}$-order homogeneous difference equation to be any solution which is *not* the solution to any lower order difference equation.

(a) Show that if $s$ is a full solution to a $k^{\text{th}}$-order homogeneous difference equation then the sequences $s, \sigma(s), \ldots, \sigma^{k-1}(s)$ are linearly independent.

(b) Show that $< 2^n >$ is not a full solution to $s_n = 4s_{n-1} - 4s_{n-2}$, but $< n2^n >$ is a full solution to the recurrence.

(c) Show that every solution to $s_n = 4s_{n-1} - 4s_{n-2}$ can be written as

$$s_n = a_1 n 2^n + a_2 (n-1) 2^{n-1}$$

for some constants $a_1, a_2$.

**Ex 3.6.** For the polynomial $ch(x) = x^k - c_1 x^{k-1} - \ldots - c_k$ and any $i = 1, \ldots, k$ show that $t_i = \sum_{j=1}^{k} c_j (-j)^i \lambda^{k-j}$ can be written as a linear combination of $ch(x), D(ch)(x), \ldots, D^{(j)}(ch)(x)$.

**Ex 3.7.** Solve

$$s_0 = 2\,,\ s_1 = -3\,,\ s_{n+2} = -3s_{n+1} - 2s_{n-2} + (-1)^n\,,\ \textit{for } n \geq 2.$$

**Ex 3.8.** Use the Finiteness Argument to show that the ring of formal power series has no zero divisors; that is, if $\gamma_1, \gamma_2$ are two power series whose product is the zero polynomial, then at least one of $\gamma_1$, $\gamma_2$ is the zero polynomial.

**Ex 3.9.** Let $\gamma_1$ and $\gamma_2$ be two formal power series in $x$. Show that $\gamma_1(\gamma_2(x))$ makes sense as a formal power series in $x$, provided the constant term of $\gamma_2$ is zero. Check that the substitution of $\gamma_2 = 1 + x$ (which has non-zero constant term) doesn't work.

**Ex 3.10.** Let $\Psi(\gamma_1, \ldots, \gamma_t)$ be a polynomial in finitely many power series $\gamma_1, \ldots, \gamma_t$ and let $P$ be a polynomial in $x$ with zero constant term. Use the Extended Finiteness Argument to show that first expanding $\Psi$ and then substituting $P$ for $x$ results in the same power series as first substituting $P$ for $x$ and then expanding $\psi$.

**Ex 3.11 (The Quadratic Formula for Power Series).** If $\alpha = \sum_{i \geq 1} a_i x^i$ where each $a_i$ is real, show there exists a unique $\beta = \sum_{i \geq 1} b_i x^i$ such that $\beta$

satisfies $(1+\beta)^2 = 1+\alpha$ (and we can denote $1+\beta = \sqrt{1+\alpha}$). Use this to prove that whenever $\gamma_1 = \sum_{i\geq 0} a_i x^i$ and $\gamma_2 = \sum_{i\geq 0} b_i x^i$ are power series with real coefficients and $a_0^2 - 4b_0 \geq 0$ then the equation

$$x^2 + \gamma_1 x + \gamma_0 = 0$$

has the two solutions in power series given by

$$\frac{-\gamma_1 \pm \sqrt{\gamma_1^2 - 4\gamma_2}}{2}\,.$$

The process of **convolution** appeared when we defined multiplication of power series. It occurs in many other contexts in mathematics. This next sequence of exercises explains convolution and gives some examples. The definitions of a group and a ring are needed to work most of these exercises.

**Ex 3.12.** Show that the product of two formal power series can be defined as

$$\left(\sum_{i\geq 0} a_i x^i\right)\left(\sum_{j\geq 0} b_j x^j\right) = \sum_{n\geq 0}\left(\sum_{l\geq 0} a_l b_{n-l}\right) x^n$$

where $b_{n-l}$ is considered to be zero for negative subscripts.

**Ex 3.13.** Let $G$ be an additive abelian group and let $\phi, \psi : G \to \mathbb{R}$ be functions which are zero for all but finitely many $g \in G$. Define the convolution $\phi * \psi$ by

$$(\phi * \psi)(g) = \sum_{h\in G} \phi(h)\psi(g-h)\,.$$

Show that

(a) $\phi * \psi$ is well defined (meaning that the sum above is finite);
(b) $(\phi * \psi)(g) = 0$ for all but finitely many $g \in G$;
(c) $\phi$ commutes with $\psi$; that is, $\phi * \psi = \psi * \phi$.

**Ex 3.14.** Let $G$ be as in Exercise 3.13 and consider the set $R$ of all functions $G \to \mathbb{R}$ for which are zero for all but finitely many elements of $G$.

(a) Using function addition for the addition and using convolution for the multiplication, show $R$ is a commutative ring.
(b) Show that the function which is 1 at the additive identity of $G$ and is zero elsewhere is a multiplicative identity for this ring.
(c) Either show that $R$ is guaranteed to have no zero divisors or find a group $G$ for which the associated $R$ does have zero divisors.

**Ex 3.15.** As usual, $\mathbb{R}[x]$ denotes the set of polynomials in the variable $x$ with real coefficients. For any polynomial in $\mathbb{R}[x]$, define a corresponding function on $\mathbb{N}$ whose value at $i \in \mathbb{N}$ is the $i^{\text{th}}$ coefficient of the polynomial. Use this to show that $\mathbb{R}[x]$ is the ring defined in Exercise 3.14 with $G = \mathbb{N}$.

**Ex 3.16.** Consider the set $R = \{\phi : \mathbb{Z} \to \mathbb{R} \mid \phi(n) = 0 \text{ for all } n < 0\}$. Define the convolution of two such functions by

$$(\phi * \psi)(n) = \sum_{i \in \mathbb{N}} \phi(i)\psi(n-i)\,.$$

Show that $R$ forms a commutative ring under the operations of usual addition of functions and convolution. Identify an identity element for convolution.
Note: This ring $R$ is the ring of formal power series with real coefficients.

**Ex 3.17.** Let $f, g : \mathbb{R} \to \mathbb{R}$ be two integrable functions. Define the convolution of $f$ and $g$ by

$$f * g(t) = \int_{-\infty}^{+\infty} f(u)g(t-u)du\,.$$

Without worrying about the convergence of the integral or the integrability of the resulting function, explain how this is a continuous analog of the convolutions defined above. Show that this serves as multiplication, making the set of integrable functions into a ring. If you know enough, worry about the convergence of the integrals. If you know what the Dirac Delta function is, show that it's an identity for convolution.

The point of the next string of exercises (Exercises 3.18 to 3.26) is to give another perspective on the finiteness arguments used to verify the algebraic properties of power series and, more generally, on the definition of formal power series. These should only be seriously attempted by a student who has had a course in the foundations of real analysis. Other students are encouraged to read through these exercises.

**Ex 3.18.** Let $\mathbb{R}[x]$ be the set of polynomials in $x$ with real coefficients. Given a non-zero polynomial in $\mathbb{R}[x]$, define its **order** $\operatorname{ord}(f)$ to be the degree of the lowest term with a non-zero coefficient. Thus, the order of a polynomial is 0 unless its constant term is zero, and if the constant term is zero, the order is 1 unless there is no linear term, and so on. Show that the order of $f$ is $m \geq 1$ if and only if $x = 0$ is a root of $f$ with multiplicity $m$.

**Ex 3.19.** Pick your favorite constant $c > 1$ and define a function $|\cdot| : \mathbb{R}[x] \to \mathbb{N}$ by $|0| = 0$ and $|f| = c^{-\operatorname{ord}(f)}$ for non-zero $f$. Show that this **absolute value** satisfies the following familiar properties:

(a) $|f| \geq 0$ and $|f| = 0$ if and only if $f = 0$. ($|\cdot|$ is positive definite.)
(b) $|f + g| \leq |f| + |g|$. (The triangle inequality.)
(c) $|f - g| = |g - f|$. (Symmetry.)
(d) $|fg| = |f||g|$. (Multiplicativity.)

**Ex 3.20.** Review the construction of the real numbers $\mathbb{R}$ from the rational numbers $\mathbb{Q}$ using **Cauchy sequences**, paying careful attention to the

properties of the absolute value which are used to make this construction work. Now define a Cauchy sequence of polynomials in $\mathbb{R}[x]$ to be a sequence $< f_n >$ of polynomials such that, for every $\epsilon > 0$, there is some $M$ such that $|f_n - f_m| < \epsilon$ whenever $m, n \geq M$ (where $|\cdot|$ is defined in the previous exercise). Next, define two Cauchy sequences $< f_n >$ and $< g_n >$ of polynomials to be equivalent if for every $\epsilon > 0$ there is an $M$ such that $|f_n - g_m| < \epsilon$ for $m, n \geq M$. Show this is an equivalence relation on the set of Cauchy sequences of polynomials. To do this, you do not need to have any idea of what a Cauchy sequence looks like—the only thing you need to use are the properties of the absolute value.

**Ex 3.21.** Show that every Cauchy sequence of polynomials is equivalent to a unique sequence of $< g_d >$ of polynomials such that $\deg g_d = d$ and $g_d = g_{d-1} + a_d x^d$ for all $d \geq 1$. In other words, a sequence such that $g_d$ is the partial sum of a formal power series in $x$. Conclude that equivalence classes of Cauchy sequences of polynomials are the same objects as formal power series.

**Ex 3.22.** Mimicking the construction of $\mathbb{R}$ from $\mathbb{Q}$, show how to define arithmetic operations on equivalence classes of Cauchy sequences of polynomials and prove that these are well defined (independent of which equivalent sequences are used.) Show that these operations obey the same algebraic laws as they do when applied to polynomials. The last exercise says that you are really defining operations on formal power series, but you should do this exercise blind to this and only use properties of $|\cdot|$.

**Ex 3.23.** Show that the arithmetic operations defined above are the same ones we defined for formal power series.

**Ex 3.24.** This is a generalization of Exercise 3.19 from polynomials to power series. Define an absolute value on $\mathcal{S}^+$ as follows. Given a non-zero sequence $< x_n >$, define the order of $\mathrm{ord}(x_n)$ to be the first $n$ for which $x_n \neq 0$. Choosing your favorite $c > 1$ again, let

$$\| < x_n > \| = c^{-\mathrm{ord}(x_n)}.$$

Show that this absolute value is positive definite, symmetric and satisfies the triangle inequality.

**Ex 3.25.** A space with an absolute value such as the ones above is called a **metric space** and, important for us, is called a **complete metric space** if every Cauchy sequence of elements from the space has a limit in the space. Show that $\mathcal{S}^+$ is complete by showing that, if we are given a sequence $< x^i >$ *of sequences* $< x^i_n >$ which is Cauchy in the sense that for every $\epsilon > 0$ there is an $I$ with the property that $\|x^i - x^j\| < \epsilon$ for $i, j \geq I$, then there is a sequence $y$ so that $\lim_{i \to \infty} x_i = y$. (This limit involves *another* epsilon-$I$ definition, this time with $\|\cdot\|$.)

**Ex 3.26.** (This exercise requires the preceding exercises.)

(a) Show that if $\lim_{n\to\infty} f_n = \Gamma$, where the $f_n$ are polynomials in $\sigma$ and $\Gamma$ is a power series in $\sigma$, then for any sequence $s \in \mathcal{S}^+$, we have $\lim_{n\to\infty} f_n(s)$ exists and is equal to the sequence $\Gamma(s)$ as defined in Section 3.4.2.
(b) Show that any formal power series in $\sigma$ is a linear operator on $\mathcal{S}^+$.

**Ex 3.27.** Use induction to show that for all $j \geq 0$,

$$\mathcal{D}^j\Big(\sum_{n\geq 0} x^n\Big) = \frac{j!}{(1-x)^{j+1}},$$

where $0! = 1$ and $\mathcal{D}^0$ is the identity operator.

**Ex 3.28.** Let $D^j$ denote the ordinary $j$-th derivative with respect to $x$. For any formal power series $S(x) = \sum_{k\geq j} s_k x^k$ show that for all $j \geq 1$,

$$\mathcal{D}^j\Big(S(x)\Big) = \sum_{k\geq j} s_k D^j(x^k) = j! \sum_{k\geq 0} \binom{k+j}{j} s_{k+j} x^k.$$

**Ex 3.29.** For rationals $r, s$, use the fact that $(1+x)^{r+s} = (1+x)^r(1+x)^s$ to show that the binomial coefficients satisfy

$$\binom{r+s}{k} = \sum_{i+j=k} \binom{r}{i}\binom{s}{j} \text{ for all } k \geq 0\,.$$

Also use $(1-x^2)^r = (1-x)^r(1+x)^r$ to get

$$\binom{r}{n/2} = (-1)^{n/2} \sum_{k=0}^{n} \binom{r}{k}\binom{r}{n-k} \text{ for even } n\,.$$

**Ex 3.30.** For $L = 1-2\sigma+\sigma^2$, show $L^{-1} = \sum_{i\geq 0}(i+1)\sigma^i$. Show the inverse of $1-\sigma-\sigma^2$ equals $\sum_{i\geq 0} f_{i+1}\sigma^i$ where $f_i$ is the $i^{\text{th}}$ Fibonacci number. From this calculation derive a representation of the Fibonacci numbers as sums of binomial coefficients.

**Ex 3.31.** Show that $\mathcal{D}(\gamma) = 0$ implies that the power series $\gamma$ is a constant. Use that to prove that $\mathcal{D}(\gamma_1) = \mathcal{D}(\gamma_2)$ implies that $\gamma_1 = \gamma_2 + c$ for some constant $c$. Further show that the general solution to $\mathcal{D}(\gamma) = \sum_{n\geq 0} a_n x^n$ is $\gamma = c + \sum_{n\geq 1} \frac{a_{n-1}}{n} x^n$.

**Ex 3.32 (Exponential power series).** Show that $\mathcal{D}(\gamma) = \gamma$ implies $\gamma = \sum_{n\geq 0} \frac{1}{n!} x^n$, the Taylor series for $e^x$. Use this fact to show that $\gamma = \sum_{n\geq 0} \frac{1}{n!}(-x)^n$ is the multiplicative inverse of $\gamma = \sum_{n\geq 0} \frac{1}{n!} x^n$. (Hint: Multiply by $e^{-x}$.)

**Ex 3.33.** (a) Use the product rule for formal differentiation to show that for any formal power series $\gamma$, $e^{-x}\mathcal{D}(\gamma) - \gamma e^{-x} = \mathcal{D}(e^{-x}\gamma)$.

(b) Solve the formal equation $\mathcal{D}(\gamma) - \gamma = \sum_{n\geq 0} \frac{1}{n!}x^n$ subject to the condition that $\gamma = 1 + \sum_{n\geq 1} a_n x^n$. (HINT: Multiply both sides by $e^{-x}$.)

# 4
# Generating Functions

This chapter is devoted to the study of generating functions, a technique for solving difference equations. After giving some motivation for the term "generating function" in the first section, the remainder of the chapter concentrates on using generating functions to solve recurrences. In this study we obtain new proofs of some earlier results as well as an efficient method for solving a general class of initial value problems.

The term "generating function" is said to have been coined by Pierre-Simon Laplace (1749-1827), but they formed the basis for a technique used earlier by Abraham DeMoivre in 1730 [40] to solve linear recurrences.

## 4.1 Counting strings with some restrictions

To introduce generating functions we begin with a combinatorial problem about strings whose solution uses generating polynomials. For any $n \geq 0$, an ordered list of $n$ symbols from an alphabet is called an **n-string**. The only 0-string is the string with no elements, named the **nullstring**.

For the alphabet $\mathbf{A} = \{x, y, z\}$, there are three 1-strings, the symbols in the alphabet $\mathbf{A}$. One way to obtain a listing of all 2-strings is to use the formal algebraic sum obtained from multiplying $(x + y + z) \cdot (x + y + z)$ according to the distributive law for multiplication:

$$(x + y + z) \cdot (x + y + z) = (x + y + z)x + (x + y + z)y + (x + y + z)z;$$

$$(4.2)\quad (x+y+z)\cdot(x+y+z) = xx+yx+zx+xy+yy+zy+xz+yz+zz.$$

Since each 2-string appears exactly once as a summand on the right side of (4.2), the polynomial $(x+y+z)\cdot(x+y+z)$ is said to generate the 2-strings. Using the commutative law and the laws of exponents, (4.2) can be compressed to

$$(4.3)\qquad (x+y+z)^2 = 1x^2+2xy+2xz+2yz+1y^2+1z^2\,.$$

Although the compressed polynomial in (4.3) is no longer a complete listing of 2-strings, it does encode some important information about the strings. For instance, the term $2xy$ indicates there are two 2-strings with one $x$ and one $y$; $1z^2$ says that there is one 2-string which uses only the symbol $z$. In general, for all $n \geq 0$ the polynomial $P_n(x,y,z) = (x+y+z)^n$ generates all $n$-strings formed using the alphabet $\mathbf{A} = \{x,y,z\}$, and the coefficient of $x^i y^j z^k$ in its compressed representation counts the number of $n$-strings which has $i$ $x$'s, $j$ $y$'s, and $k$ $z$'s.

When $x=1, y=1, z=1$ is substituted into $P_n(x,y,z)$, each summand in (4.2) is 1 and accordingly $P_n(1,1,1) = 3^n$, the total number of $n$-strings. The value of $P_n(x,y,z)$ for other choices of the variables $x,y,z$ can be used to count the number of strings with certain characteristics. For instance, suppose we want to calculate the four quantities $s_{00}^{(n)}, s_{01}^{(n)}, s_{10}^{(n)}, s_{11}^{(n)}$, where for any $i,j \in \{0,1\}$, $s_{ij}^{(n)}$ equals the number of $n$-strings in which the number of $y$'s is $i \pmod 2$ and the number of $z$'s is $j \pmod 2$. For example, for $n=2$ from (4.2) we obtain

$$s_{00}^{(2)} = |\{xx, yy, zz\}| = 3\ ;\ s_{01}^{(2)} = |\{zx, xz\}| = 2\ ;$$
$$s_{10}^{(2)} = |\{yx, xy\}| = 2\ ;\ s_{11}^{(2)} = |\{zy, yz\}| = 2.$$

Since each $n$-string is counted once in the numbers $s_{00}^{(n)}, s_{01}^{(n)}, s_{10}^{(n)}, s_{11}^{(n)}$, then

$$(4.4)\qquad 3^n = (1+1+1)^n = P_n(1,1,1) = s_{00}^{(n)} + s_{01}^{(n)} + s_{10}^{(n)} + s_{11}^{(n)}$$

holds for all $n \geq 0$. When we substitute $x=1, y=-1, z=1$ into $P_n(x,y,z)$ each string with an odd number of $y$'s is counted as negative. Since strings with an odd number of $y$'s are counted in $s_{10}^{(n)}$ and $s_{11}^{(n)}$ then

$$(4.5)\qquad 1 = (1-1+1)^n = P_n(1,-1,1) = s_{00}^{(n)} + s_{01}^{(n)} - s_{10}^{(n)} - s_{11}^{(n)}.$$

Similarly,

$$(4.6)\qquad 1 = (1+1-1)^n = P_n(1,1,-1) = s_{00}^{(n)} - s_{01}^{(n)} + s_{10}^{(n)} - s_{11}^{(n)}$$

and

$$(4.7)\quad (-1)^n = (1-1-1)^n = P_n(1,-1,-1) = s_{00}^{(n)} - s_{01}^{(n)} - s_{10}^{(n)} + s_{11}^{(n)}.$$

The equations in (4.4) through (4.7) form a nonsingular system of four equations in four unknowns and can be solved to obtain all of $s_{00}^{(n)}$, $s_{01}^{(n)}$, $s_{10}^{(n)}$, $s_{11}^{(n)}$.

For example, consider the question of finding a formula for $s_{00}^{(n)}$ which would calculate the number of $n$-strings which have an even number of $y$'s and an even number of $z$'s. Obtaining a complete list of all $n$-strings for large $n$ would be a daunting task. On the other hand, summing the four equations (4.4) through (4.7) results in

$$3^n + 2 + (-1)^n = 4s_{00}^{(n)} ,$$

and we have the exact formula

$$s_{00}^{(n)} = \frac{3^n + 2 + (-1)^n}{4}, \tag{4.9}$$

which has been obtained without listing all permissible strings. Checking this formula for $n = 2$, $xx, yy, zz$ is a complete listing of permissible strings and $(3^2+2+(-1)^2)/4$ does equal 3. The values of $P_n(1, y, z)$ for $y, z \in \{\pm 1\}$ allowed us to do this computation.

We close this section with an indication of how recurrence techniques from earlier chapters can also be used to calculate $s_{00}^{(n)}$. To get a recurrence relation which relates $s_{00}^{(n)}$ to some $s_{jk}^{(i)}$ for earlier $i < n$, we decompose each $n$-string into two substrings, its first character and the remaining string of length $n-1$. Any permissible $n$-string has one of the following three forms: Its first character is $x$ and the rest is a permissible $(n-1)$-string; its first character is $y$ and the remaining $(n-1)$-string has an odd number of $y$'s and an even number of $z$'s; its first character is $z$ and the rest has an even number of $y$'s and an odd number of $z$'s. Among the $n$-strings counted in $s_{00}^{(n)}$ there are $s_{00}^{(n-1)}$ strings of the first type, $s_{10}^{(n-1)}$ of the second type and $s_{01}^{(n-1)}$ of the third type. Therefore,

$$s_{00}^{(n)} = s_{00}^{(n-1)} + s_{10}^{(n-1)} + s_{01}^{(n-1)}. \tag{4.10}$$

Similar arguments give

$$s_{01}^{(n)} = s_{01}^{(n-1)} + s_{11}^{(n-1)} + s_{00}^{(n-1)}; \tag{4.11}$$

$$s_{10}^{(n)} = s_{10}^{(n-1)} + s_{00}^{(n-1)} + s_{11}^{(n-1)}; \tag{4.12}$$

$$s_{11}^{(n)} = s_{11}^{(n-1)} + s_{01}^{(n-1)} + s_{10}^{(n-1)}. \tag{4.13}$$

This approach to the problem is completed in Exercises 4.2 and 4.3.

## 4.2 An Overview of the Generating Function Technique

We've just shown how the polynomial $P_n(x, y, z) = (x+y+z)^n$ can be used to generate all $n$-strings from the alphabet $\mathbf{A} = \{x, y, z\}$. The remainder of this chapter will consider generating functions in *one* variable, which form the basis of a technique for solving linear recurrences. For a more comprehensive discussion of generating functions and their applications the reader is referred to [148]. Also, in [133], Richard Stanley gives a survey of the use of generating functions in combinatorics.

We first consider the simplest case, *polynomial* generating functions in one variable. For this, recall that a polynomial in the variable $x$ with complex coefficients can be written in the form $p(x) = s_0 + s_1x + \ldots + s_rx^r$ for some complex constants $s_0, s_1, \ldots, s_r$. Complete knowledge of the polynomial is equivalent to knowledge of its coefficients, the sequence $< s_0, s_1, \ldots, s_r >$. In this way $p(x)$ can be considered to be the generating polynomial for the finite sequence $< s_0, s_1, \ldots, s_r >$, and conversely $< s_0, s_1, \ldots, s_r >$ is the sequence generated by the polynomial $p(x)$. For example, the **Binomial Theorem** says

$$(1+x)^n = \sum_{k=0}^{n} \binom{n}{k} x^k, \text{ for any positive integer } n$$

where the **binomial coefficient** $\binom{n}{k} = \dfrac{n!}{k!(n-k)!}$ is the number of combinations of $n$ things taken $k$ at a time, and the polynominal $p_n(x) = (1+x)^n$ records the sequence $\binom{n}{0}, \binom{n}{1}, \ldots, \binom{n}{n}$. Because of this, $p_n(x)$ can be called the generating function for the $n^{\text{th}}$ row of Pascal's triangle. Here the generating function is a polynomial because each row of Pascal's triangle contains only finitely many entries. Because the solution of an initial value problem is an infinite sequence, its generating function is an infinite series rather than a polynomial unless the solution sequence is eventually zero.

### The Generating Function of a Sequence

*The* **generating function** *for an infinite sequence $< s_n >$ is the formal power series*

$$S(x) = \sum_{n \geq 0} s_n x^n.$$

*When the sequence is finite, say $< s_0, s_1, \ldots, s_r >$, its generating function is the polynomial*

$$s_0 + s_1x + \ldots + s_rx^r.$$

As in Chapter 3, a generating function is a power series, an algebraic object which can be formally manipulated using algebraic operations, and the sum and product of the generating functions $S(x) = \sum_{n\geq 0} s_n x^n$ and $T(x) = \sum_{n\geq 0} t_n x^n$ are respectively

$$S(x) + T(x) = \sum_{n\geq 0} (s_n + t_n)x^n \text{ and } S(x) \cdot T(x) = \sum_{n\geq 0} \Big(\sum_{j=0}^{n} s_j t_{n-j}\Big) x^n.$$

Remember that a formal power series is never evaluated at a specific value of $x$ unless it can be shown it's more than purely formal by proving it converges on an open interval containing $x$.

The following examples demonstrate the generating function method for solving linear recurrences.

*Example* 4.2.1. **(The general first-order homogeneous recurrence)** For $\lambda \in \mathbb{C}$, the generating function $S(x)$ for solution to the recurrence $s_{n+1} = \lambda s_n$ satisfies

$$S(x) = s_0 + x \sum_{n\geq 0} s_{n+1} x^n = s_0 + x \sum_{n\geq 0} \lambda s_n x^n = s_0 + \lambda x S(x),$$

which can be solved for $S(x)$ to get

$$S(x) = \frac{s_0}{1 - \lambda x}. \tag{4.14}$$

By Theorem 3.4.3,

$$S(x) = \frac{s_0}{1 - \lambda x} = \sum_{n\geq 0} s_0 \lambda^n x^n,$$

and equality of formal power series gives $s_n = s_0 \lambda^n$, as predicted.

To summarize, we used the recurrence to find a rational form for the generating function and then "expanded" this rational function into another power series. The solution was obtained by equating the coefficients of the two power series.

*Example* 4.2.2. **(The Fibonacci sequence)** The Fibonacci generating function is the formal power series $F(x) = \sum_{n\geq 0} f_n x^n$ which satisfies

$$F(x) = f_0 + f_1 x + \sum_{n\geq 2} f_n x^n = x + \sum_{n\geq 2} (f_{n-1} + f_{n-2}) x^n.$$

At this point a creative re-indexing can be performed to get

$$F(x) = x + \sum_{n\geq 1} f_n x^{n+1} + \sum_{n\geq 0} f_n x^{n+2} = x + x \sum_{n\geq 0} f_n x^n + x^2 \sum_{n\geq 0} f_n x^n$$

which can be rewritten as

$$F(x) = x + xF(x) + x^2F(x).$$

Collecting the terms containing $F(x)$ and solving we obtain

$$F(x) = \frac{x}{1 - x - x^2}, \tag{4.15}$$

which is more complicated than the same stage (4.14) of the last example. The technique of partial fractions from calculus can be used to decompose $F(x)$ into the sum of two generating functions, each of which is a geometric power series. (The general technique of partial fractions is developed in Section 4.3.)

In order to decompose the rational function on the right side of (4.15) into partial fractions, we find the roots of $1 - x - x^2 = 0$ to be $\lambda_1^{-1} = \dfrac{-1 + \sqrt{5}}{2}$, $\lambda_2^{-1} = \dfrac{-1 - \sqrt{5}}{2}$. Therefore, from $\lambda_1^{-1}\lambda_2^{-1} = -1$ we have

$$1 - x - x^2 = -(\lambda_1^{-1} - x)(\lambda_2^{-1} - x) = (1 - \lambda_1 x)(1 - \lambda_2 x).$$

By partial fractions there exist constants $A, B$ such that

$$F(x) = \frac{x}{1 - x - x^2} = \frac{A}{1 - \lambda_1 x} + \frac{B}{1 - \lambda_2 x}. \tag{4.16}$$

Before finding the constants $A$ and $B$, we note that Theorem 3.4.3 allows us to write the right-hand side of (4.16) as the formal power series

$$F(x) = \sum_{n \geq 0} \left( A\lambda_1^n + B\lambda_2^n \right) x^n$$

and the uniqueness of formal power series implies

$$f_n = A{\lambda_1}^n + B{\lambda_2}^n, \quad \text{for all } \ n \geq 0\,. \tag{4.18}$$

To find $A$ and $B$ return to (4.16). Since

$$(1 - \lambda_1 x)(1 - \lambda_2 x) = 1 - x - x^2,$$

we have

$$\frac{x}{1 - x - x^2} = \frac{A(1 - \lambda_2 x) + B(1 - \lambda_1 x)}{1 - x - x^2}.$$

Because these two rational expressions for $F(x)$ have the same denominator, the numerators must also be equal, giving the polynomial equation

$$x = A(1 - \lambda_2 x) + B(1 - \lambda_1 x)\,.$$

As polynomials this implies

$$0 = A + B \text{ and } 1 = -A\lambda_2 - B\lambda_1 ,$$

and (4.18) becomes **Binet's Formula** (which we already mentioned in Chapter 1)

$$f_n = \frac{1}{\sqrt{5}}\Big(\Big(\frac{1+\sqrt{5}}{2}\Big)^n - \Big(\frac{1-\sqrt{5}}{2}\Big)^n\Big) .$$

*Example* 4.2.3. Consider the recurrence

$$s_0 = 2 \ , \ s_1 = 3 \ , \ s_{n+2} = 6s_{n+1} - 9s_n \text{ for all } \ n \geq 0. \tag{4.21}$$

Its generating function is

$$\begin{aligned} S(x) = \sum_{n\geq 0} s_n x^n &= s_0 + s_1 x + \sum_{n\geq 0} s_{n+2} x^{n+2} \\ &= 2 + 3x + \sum_{n\geq 0} (6s_{n+1} - 9s_n) x^{n+2} \\ &= 2 + 3x + 6x \sum_{n\geq 0} s_{n+1} x^{n+1} - 9x^2 \sum_{n\geq 0} s_n x^n \\ &= 2 + 3x + 6x \sum_{n\geq 1} s_n x^n - 9x^2 \sum_{n\geq 0} s_n x^n \\ &= 2 + 3x + 6x(S(x) - 2) - 9x^2 S(x) \end{aligned}$$

and

$$(1 - 6x + 9x^2)S(x) = 2 - 9x . \tag{4.22}$$

Therefore,

$$S(x) = \frac{2-9x}{1-6x+9x^2} = \frac{2-9x}{(1-3x)^2} = \frac{A}{1-3x} + \frac{B}{(1-3x)^2},$$

and we can solve to get $A = 3$, $B = -1$. Using this and (3.16)

$$\begin{aligned} S(x) &= \frac{3}{1-3x} - \frac{1}{(1-3x)^2} \\ &= 3\sum_{n\geq 0} 3^n x^n - \sum_{n\geq 0} (n+1)(3x)^n ; \end{aligned}$$

$$s_n = 3^{n+1} - (n+1)3^n = (2-n)3^n .$$

As noted earlier, the first step in each of these three solutions was to represent the generating function as a rational function in which the denominator bears a strong resemblance to the characteristic polynomial of

the recurrence. Namely, in each case the generating function of the homogeneous equation

$$s_{n+k} = c_1 s_{n+k-1} + \ldots + c_k s_n$$

is represented as a rational function whose denominator is the **reciprocal polynomial**

$$ch^R(x) = 1 - c_1 x - \ldots - c_k x^k$$

of the characteristic polynomial

$$ch(x) = x^k - c_1 x^{k-1} - c_2 x^{k-2} - \ldots - c_k \,.$$

Before proving the general result, we consider one more example, this time a *nonhomogeneous* equation

$$(4.24) \quad s_0 = 1 \,,\ s_1 = 4 \,,\ s_{n+2} = 6s_{n+1} - 9s_n + 4(n-1) \text{ for all } \ n \geq 0 \,.$$

As in the last example, we have

$$\begin{aligned} S(x) &= 1 + 4x + \sum_{n\geq 0} s_{n+2} x^{n+2} \\ &= 1 + 4x + 6x \sum_{n\geq 1} s_n x^n - 9x^2 \sum_{n\geq 0} s_n x^n + 4x^3 \sum_{n\geq 0} n x^{n-1} - 4x^2 \sum_{n\geq 0} x^n \,. \end{aligned}$$

At this point we want the rational representation of $\sum_{n\geq 0} n x^{n-1}$, which is a special case of the following formula proved in Chapter 3.

$$(4.25) \qquad \frac{1}{(1-\lambda x)^m} = \sum_{n\geq 0} \binom{n+m-1}{n} (\lambda x)^n \text{ for any integer } m \geq 1 \,.$$

From this,

$$S(x) = 1 + 4x + 6x(S(x) - 1) - 9x^2 S(x) + \frac{4x^3}{(1-x)^2} - \frac{4x^2}{1-x} \,,$$

which gives

$$(4.26) \qquad (1 - 6x + 9x^2) S(x) = \frac{(1-3x)(1-x-2x^2)}{(1-x)^2}$$

and

$$S(x) = \frac{1 - x - 2x^2}{(1-x)^2(1-3x)} \,.$$

(Here the denominator is not the reciprocal polynomial because the input function contributes a factor of $(1-x)^2$ and allows the cancellation

of $1-3x$.) The solution is completed as above, with the partial fraction decomposition

$$S(x) = \frac{1}{1-3x} + \frac{1}{(1-x)^2} - \frac{1}{1-x}$$

yielding the solution

$$s_n = 3^n + (n+1) - 1 = 3^n + n\,.$$

It is interesting to note that because 3 is not a double root of the denominator of the rational function in this example, the solution sequence is actually the solution to a *first-order* recurrence with $ch(x) = x - 3$ and so is not a full solution (refer to Exercise 3.5). In fact, from

$$s_{n+1} - 3s_n = (3^{n+1} + n + 1) - 3(3^n + n) = 1 - 2n$$

we see that $< s_n >$ is a solution to $s_{n+1} = 3s_n + 1 - 2n$.

We're now ready to determine the rational function which encodes the initial value problem

$$\text{(L)} \qquad s_{n+k} = c_1 s_{n+k-1} + c_2 s_{n+k-2} + \ldots + c_k s_n + \psi(n) \;\text{ with }\; S_0 = \alpha_0$$

for any input sequence $\psi(n)$ of complex numbers. Let $S(x) = \sum_{n\geq 0} s_n x^n$ and $F(x) = \sum_{n\geq 0} \psi(n) x^n$, the generating functions of $< s_n >$ and $< \psi(n) >$ respectively. Then

$$\begin{aligned}
S(x)\; ch^R(x) &= S(x)(1 - c_1 x - \ldots - c_k x^k) \\
&= \sum_{n\geq 0} s_n x^n - c_1 \sum_{n\geq 0} s_n x^{n+1} - \ldots - c_k \sum_{n\geq 0} s_n x^{n+k} \\
&= \sum_{n\geq 0} s_n x^n - c_1 \sum_{n\geq 1} s_{n-1} x^n - \ldots - c_k \sum_{n\geq k} s_{n-k} x^n \\
&= \sum_{n\geq k} \Big( s_n - c_1 s_{n-1} - \ldots - c_k s_{n-k} \Big) x^n + \sum_{n=0}^{k-1} s_n x^n \\
&\qquad - c_1 \sum_{n=1}^{k-1} s_{n-1} x^n - \ldots - c_{k-2} \sum_{n=k-2}^{k-1} s_{n-k+2} x^n - c_{k-1} s_0 x^{k-1}\,.
\end{aligned}$$

The first summand is

$$\sum_{n\geq 0} \Big( s_{n+k} - c_1 s_{k+n-1} - \ldots - c_k s_n \Big) x^{n+k} = x^k \sum_{n\geq 0} \psi(n) x^n = x^k F(x)\,,$$

which only depends on $F(x)$ and $k$. The remainder of the expression for $S(x) ch^R(x)$ contains the information about the initial conditions, and is

the polynomial

$$\begin{array}{lllll} s_0 & +s_1x & +s_2x^2+ & \dots & +s_{k-1}x^{k-1} \\ & -c_1s_0x & -c_1s_1x^2- & \dots & -c_1s_{k-2}x^{k-1} \\ & & -c_2s_0x^2- & \dots & -c_2s_{k-3}x^{k-1} \\ & & & \dots & \\ & & & & -c_{k-1}s_0x^{k-1} \end{array}.$$

Writing the coefficients as $d_0, d_1, \dots, d_{k-1}$, then $d_i = s_i - c_1s_{i-1} - \dots - c_is_0$ is the dot product

$$d_i = (0, \dots, 0, 1, -c_1, \dots, -c_i) \cdot (s_{k-1}, \dots, s_0),$$

which can be written as $(d_{k-1}, \dots, d_0)^T = M_0S_0$ where $M_0$ is the $k \times k$ matrix

$$M_0 = \begin{bmatrix} 1 & -c_1 & -c_2 & \dots & -c_{k-1} \\ 0 & 1 & -c_1 & \dots & -c_{k-2} \\ 0 & 0 & 1 & \dots & -c_{k-3} \\ \dots & & & & \\ 0 & 0 & 0 & \dots & 1 \end{bmatrix}. \tag{4.27}$$

This gives a rational representation of the generating function for (L):

**The Rational Representation**

*If $F(x)$ is the generating function of $< \psi(n) >$, then the generating function of the solution to (L) with initial values is*

$$S(x) = \frac{d(x) + x^kF(x)}{ch^R(x)}, \tag{4.28}$$

*where $d(x)$ is the polynomial with coefficients $(d_{k-1}, \dots, d_0)^T = M_0S_0$.*

From this we see that the representation is a rational function exactly when $F(x)$, the generating function for the input, can be written as a rational function.

Returning to the example in (4.24),

$$F(x) = 4\sum_{n\geq 0}(n-1)x^n = 4x\sum_{n\geq 0}nx^{n-1} - 4\sum_{n\geq 0}x^n = \frac{4x}{(1-x)^2} - \frac{4}{1-x} = \frac{4(2x-1)}{(1-x)^2},$$

and $M_0S_0 = (-2, 1)^T$ gives

$$ch^R(x)S(x) = 1 - 2x + \frac{4x^2(2x-1)}{(1-x)^2} = \frac{(1-3x)(1-x-2x^2)}{(1-x)^2},$$

as in (4.26).

## 4.3 A Review of Partial Fractions

If you've taken calculus you've probably seen the technique of partial fractions, which is used to decompose a rational function into a sum of rational functions which are simpler to integrate. For recurrences whose generating function $S(x)$ is a ratio of polynomials the partial fraction decomposition allows us to write $S(x)$ as a sum of formal power series which are in the form given in (4.25). Although your first exposure to partial fractions might have been in a calculus course, the fact that a partial fraction expansion *exists* is an algebraic result which we prove here.

Assume $p, q$ are polynomials with $\deg(p) < \deg(q)$ and that the leading coefficient of $q(x)$ equals 1. From the Fundamental Theorem of Algebra (refer to Appendix B) we know any polynomial with complex coefficients can be factored into linear polynomials, and

$$q(x) = (x - \rho_1)^{m_1} \dots (x - \rho_t)^{m_t}$$

for some $\rho_1, \dots, \rho_t \in \mathbb{C}$. The first step in our obtaining a useful partial fraction decomposition will be to show there exist polynomials $A_1(x), \dots, A_t(x)$ with each $\deg(A_i) < m_i$ such that

$$\frac{p(x)}{q(x)} = \frac{A_1(x)}{(x-\rho_1)^{m_1}} + \dots + \frac{A_t(x)}{(x-\rho_t)^{m_t}}\,. \tag{4.29}$$

We will not only show that such polynomials $A_i(x)$ exist and are unique, but also give a constructive method for finding them. Once the initial decomposition in (4.29) is found, each $A_i(x)$ can be expressed as a polynomial in $x-\rho_i$, say $A_i(x) = \alpha_{m-1}(x-\rho_i)^{m-1} + \dots + \alpha_1(x-\rho_i) + \alpha_0$ (for $m = m_i$), and then the summand $A_i(x)/(x-\rho_i)^m$ can be expanded as

$$\frac{A_i(x)}{(x-\rho_i)^m} = \frac{\alpha_{m-1}}{x-\rho_i} + \frac{\alpha_{m-2}}{(x-\rho_i)^2} + \dots + \frac{\alpha_0}{(x-\rho_i)^m}\,.$$

This is the **partial fraction decomposition** we will use, since each of these summands is in a form where (4.25) can be applied to obtain the power series for the generating function of $p(x)/q(x)$.

When $t = 1$, $q$ has only one root (which may or may not be simple) and $p(x)/q(x) = p(x)/(x-\rho_1)^{m_1}$ already has the form in (4.29). Our proof for the case when $q$ has more than one root uses the Euclidean Algorithm for polynomials. This algorithm relies on the fact that every non-zero coefficient in a complex polynomial has an inverse and so the **Division Algorithm** can be applied to any pair of non-zero polynomials $a, b$ to give unique polynomials $Q_1, R_1$ for which

$$a(x) = Q_1(x)b(x) + R_1(x) \;\text{ with }\; \deg(R_1) < \deg(b)\,.$$

This is the basis for the Euclidean Algorithm for polynomials, which is similar to the Euclidean Algorithm for integers that you may have already

seen and which we develop in Chapters 6 and 8. Once the Division Algorithm has been applied, the process can be continued with the division of $b(x)$ by $R_1(x)$ and in this way a sequence of polynomials $< R_i(x) >$ is obtained in which the degrees form a strictly decreasing sequence of natural numbers. This means the process ends after finitely many steps with $R_{n+1}(x) = 0$. That is, we have constructed two finite sequences of polynomials, $Q_1, \ldots, Q_{n+1}$ and $R_1, \ldots, R_n$ such that

$$a(x) = Q_1(x)b(x) + R_1(x)\,,\ b(x) = Q_2(x)R_1(x) + R_2(x), \ldots,$$
$$R_{n-2}(x) = Q_n(x)R_{n-1}(x) + R_n(x)\,,\ R_{n-1}(x) = Q_{n+1}(x)R_n(x) + 0\,.$$

This sequence of equations is the **Euclidean Algorithm for polynomials**. From the last equation, we note that $R_n$ divides $R_{n-1}$; using this information in the previous equation, $R_n$ also divides $R_{n-2}$; and continuing back $R_n$ divides both $a$ and $b$, and so $R_n$ is a common divisor of the original polynomials $a$ and $b$. (In Exercise 4.8 you show that $R_n$ has the largest degree among all polynomials which are common divisors. We don't use that property here.) Also, solving the penultimate equation for $R_n$ we can successively use the previous equations to obtain polynomials $B_1$ and $B_2$ such that

$$B_1(x)b(x) + B_2(x)a(x) = R_n(x)\,.$$

Let us now return to the problem at hand, justification of the existence of unique polynomials $A_1(x), \ldots, A_t(x)$ with $\deg(A_i) < m_i$ such that (4.29) holds. As we already mentioned there is nothing to prove when $t = 1$. For $t > 1$, the polynomials $a(x) = (x-\rho_1)^{m_1}$ and $b(x) = (x-\rho_2)^{m_2} \ldots (x-\rho_t)^{m_t}$ have no common factors and so the last remainder when the Euclidean Algorithm is applied to $a$ and $b$ is a constant polynomial, $R_n(x) = \alpha \in \mathbb{C}$. Also,

$$B_1(x)b(x) + B_2(x)a(x) = \alpha$$

can be multiplied by $\alpha^{-1}p(x)$ to obtain $C_1(x) = \alpha^{-1}p(x)B_1(x)$ and $C(x) = \alpha^{-1}p(x)B_2(x)$ with

$$C_1(x)b(x) + C(x)a(x) = p(x)\,, \tag{4.30}$$

giving

$$\frac{C_1(x)}{(x-\rho_1)^{m_1}} + \frac{C(x)}{b(x)} = \frac{p(x)}{q(x)}\,.$$

This process can then be inductively continued using $C(x)/b(x)$ to obtain polynomials $A_2(x), \ldots, A_t(x)$ such that (4.29) holds, provided $\deg(C) < \deg(b)$. For this, we note that by the Division Algorithm there exists a polynomial $t$ such that $A_1 = C_1 - ta$ has small degree, $\deg(A_1) < \deg(a)$. Substituting $A_1$ and $p_1 = C + tb$ in (4.30) we have

$$A_1(x)b(x) + p_1(x)a(x) = p(x)\,,$$

where

$$\deg(A_1 b) < \deg(a) + \deg(b) = \deg(q) \quad \text{and} \quad \deg(p) < \deg(q)$$

and

$$\deg(p_1) + \deg(a) = \deg(p_1 a) \leq \max\{\deg(p), \deg(A_1 b)\} < \deg(q)$$

combine to give $\deg(p_1) < \deg(b)$ as required.

In our application the denominator in the rational representation of the generating function $S(x) = p(x)/q(x)$ has a non-zero constant term, which means all of its roots $\rho_1, \ldots, \rho_t$ are non-zero and therefore each can be written as $\rho_i = 1/\lambda_i$ for some constant $\lambda_i$. The partial fraction representation becomes

$$\frac{p(x)}{q(x)} = \frac{A_1(x)}{(1-\lambda_1 x)^{m_1}} + \ldots + \frac{A_t(x)}{(1-\lambda_t x)^{m_t}} \tag{4.31}$$

where each $\lambda_i^{m_i}$ has been absorbed into the old $A_i$.

How do we actually find the polynomials $A_1(x), \ldots, A_t(x)$? Assume for the moment that the denominator $q$ has the property that it can be factored into the form

$$q(x) = (1-\lambda_1 x)\cdots(1-\lambda_k x),$$

where the non-zero constants $\lambda_1, \ldots, \lambda_k \in \mathbb{C}$ are different. We look at this case first only because its analysis is somewhat simpler. We want to determine *constants* $A_1, \ldots, A_k$ such that

$$\frac{p(x)}{q(x)} = \frac{A_1}{1-\lambda_1 x} + \ldots + \frac{A_k}{1-\lambda_k x}\,. \tag{4.32}$$

For this, set $\lambda = \lambda_1$ and multiply both sides of (4.32) by the polynomial $1-\lambda x$ to obtain

$$\frac{p(x)(1-\lambda x)}{q(x)} = A_1 + \frac{A_2(1-\lambda x)}{1-\lambda_2 x} + \ldots + \frac{A_k(1-\lambda x)}{1-\lambda_k x}$$

where $\lim_{x\to 1/\lambda} A_i(1-\lambda x)/(1-\lambda_i x)$ is zero for all $i = 2, \ldots, k$. This gives

$$\lim_{x\to 1/\lambda} \frac{p(x)(1-\lambda x)}{q(x)} = A_1\,.$$

Our method for computing the polynomials $A_1(x), \ldots, A_t(x)$ when $q(x)$ has repeated roots is only a slight modification of the one we've just given for simple roots. Setting $\lambda = \lambda_1$, $A(x) = A_1(x)$, $m = m_1$ and multiplying (4.31) by $(1-\lambda x)^m$, we have

$$\frac{p(x)(1-\lambda x)^m}{q(x)} = A(x) + (-\lambda)^m (x-a)^m T(x), \tag{4.33}$$

where

$$a = \frac{1}{\lambda} \text{ and } T(x) = \frac{A_2(x)}{(1-\lambda_2 x)^{m_2}} + \ldots + \frac{A_t(x)}{(1-\lambda_t x)^{m_t}}\,.$$

Since $\lambda \notin \{\lambda_2, \ldots, \lambda_t\}$, the function $T(x)$ is defined for all complex numbers $x$ within the circle

$$|x-a| < R \text{ with radius } R = \min\left\{ \left| \frac{1}{\lambda_i} - a \right| : i = 2, \ldots, t \right\} \neq 0$$

and the function $f(x) = \dfrac{p(x)(1-\lambda x)^m}{q(x)}$ therefore has a Taylor series (refer to Appendix B) which converges on $|x-a| < R$

$$f(x) = \sum_{i \geq 0} b_i (x-a)^i, \text{ for } b_i = \frac{D^i(f)(a)}{i!}\,.$$

(Here $D$ is the differentiation operator and $D^i(f)(a)$ is the evaluation of the $i^{\text{th}}$ derivative of $f$ at $x = a$.) Because $T(x)$ also has a Taylor series

$$T(x) = \sum_{i \geq 0} t_i (x-a)^i \text{ on } |x-a| < R\,,$$

from (4.33) we have

$$\sum_{i \geq 0} b_i (x-a)^i = f(x) = A(x) + \sum_{i \geq 0} (-\lambda)^m t_i (x-a)^{i+m}.$$

The uniqueness of the Taylor series implies that

$$A(x) = \sum_{i=0}^{m-1} b_i (x-a)^i = \sum_{i=0}^{m-1} \frac{D^i(f)(a)}{i!}(x-a)^i\,,$$

the Taylor polynomial of degree $m-1$. This result is summarized in the following.

**The Partial Fraction Decomposition**

*If* $q(x) = \prod_{i=1}^{t}(1-\lambda_i x)^{m_i}$ *(for distinct nonzero* $\lambda_1, \ldots, \lambda_t$*) then the partial fraction decomposition of any rational function* $p(x)/q(x)$ *with* $\deg(p) < q(x)$ *has the form*

$$\frac{p(x)}{q(x)} = \sum_{j=1}^{m_1} \frac{\alpha_{1j}}{(1-\lambda_1 x)^j} + \ldots + \sum_{j=1}^{m_t} \frac{\alpha_{tj}}{(q-\lambda_t x)^j}$$

*where for each* $i = 1, \ldots, t$, $\sum_{j=1}^{m_1} \alpha_{ij}(1-\lambda_i x)^{m_i - j}$ *is the Taylor polynomial about* $a_i = 1/\lambda_i$ *of degree* $m_i - 1$ *for* $f_i(x) = \dfrac{p(x)(1-\lambda_i x)^{m_i}}{q(x)}$.

Now that we have determined the form of the partial fraction decomposition, there are other ways to determine the constants $\alpha_{ij}$. For instance, using the notation above we again suppose $q$ has the factorization $q(x) = a(x)b(x)$ where

$$a(x) = (1-\lambda_1 x)^{m_1} \quad , \quad b(x) = (1-\lambda_2 x)^{m_2} \dots (1-\lambda_t x)^{m_t}$$

and $\lambda_1, \dots, \lambda_t$ are distinct and nonzero. Setting $q_1(x) = q(x)/(1-\lambda_1 x)$, from the form of the partial fraction decomposition we know there exists $\alpha_{1m} \in \mathbb{C}$ and a polynomial $p_1$ such that

$$\frac{p(x)}{q(x)} = \frac{\alpha_{1m}}{(1-\lambda_1 x)^m} + \frac{p_1(x)}{q_1(x)}\,.$$

Multiplying this by $(1-\lambda_1 x)^m$,

$$\alpha_{1m} = \frac{p(x)}{b(x)} - \frac{p_1(x)(1-\lambda_1 x)}{b(x)}$$

and since $b(1/\lambda_1) \neq 0$ then

$$\alpha_{1m} = \lim_{x \to 1/\lambda_1} \left( \frac{p(x)}{b(x)} - \frac{p_1(x)(1-\lambda_1 x)}{b(x)} \right) = \frac{p(1/\lambda_1)}{b(1/\lambda_1)}\,.$$

This determines $\alpha_{1m}$ and also gives $p(1/\lambda_1) - \alpha_{1m} b(1/\lambda_1) = 0$. Therefore, $1-\lambda_1 x$ is a factor of $P(x) = p(x) - \alpha_{1m} b(x)$, which means $p_1(x) = P(x)/(1-\lambda_1 x)$ is a polynomial for which $\deg(p_1) < \deg(b)$ and for $q_1 = b$

$$\frac{p(x)}{q(x)} = \frac{\alpha_{1m}}{(1-\lambda_1 x)^m} + \frac{p_1(x)}{q_1(x)}\,.$$

The process can be continued with $p_1(x)/q_1(x)$, and all the constants $\alpha ij$ can be found in this way.

We end this section with two examples. First we find the partial fraction decomposition of $1/(1-x)(1-2x)^2$ by calculating the required Taylor polynomials for $\rho = 1, 2$. For $\rho = 1$, we want the Taylor polynomial of degree 0 for $f_1(x) = 1/(1-2x)^2$ which is $f_1(1) = 1$, while for $\rho = 2$ the Taylor polynomial of degree 1 is required for $f_2(x) = 1/(1-x)$ which is

$$f_2(1/2) + f_2'(1/2)(x - 1/2) = 2 + 4(x - 1/2) = 2 - 2(1-2x)\,.$$

This gives the partial fraction decomposition

$$\frac{p(x)}{q(x)} = \frac{1}{1-x} + \frac{2}{(1-2x)^2} - \frac{2}{1-2x}\,.$$

Now let's use the second method for determining $\alpha_{11}, \alpha_{22}, \alpha_{21}$ such that

$$\frac{p(x)}{q(x)} = \frac{\alpha_{11}}{1-x} + \frac{\alpha 22}{(1-2x)^2} - \frac{\alpha_{21}}{1-2x}\,.$$

From $p(x) = 1$ and $b(x) = (1-2x)^2$, $\alpha_{11} = p(1)/b(1) = 1$. Also,

$$p_1(x) = \frac{1 - 1(1-2x)^2}{1-x} = 4x$$

means we want to find $\alpha_{22}, \alpha_{21}$ such that

$$\frac{p(x)}{q(x)} = \frac{\alpha 22}{(1-2x)^2} - \frac{\alpha_{21}}{1-2x}\,,$$

where now $p(x) = 4x$ and $q(x) = (1-2x)^2$. For this, the new $b(x) = 1$ and so $\alpha_{22} = p(1/2)/b(1/2) = 2$ and

$$p_1(x) = \frac{4x-2}{1-2x} = -2 = \alpha 21$$

as found above.

For another example, consider $p(x)/q(x)$ for $p(x) = -4 - 13x - 2x^2 - 7x^3 + 8x^4$ and $q(x) = 1 + x - 5x^2 - x^3 + 8x^4 - 4x^5$. To find the factorization of $q(x)$, we note that $q(1) = 0$ and divide $q(x)$ by $1 - x$ to get

$$q(x) = (1-x)(4x^4 - 4x^3 - 3x^2 + 2x + 1)\,.$$

The quotient is also divisible by $1 - x$, and in fact

$$q(x) = (1-x)^3(4x^2 + 4x + 1) = (1-x)^3(1+2x)^2\,.$$

From the above analysis we know that

$$\frac{p(x)}{q(x)} = \frac{A_1(x)}{(1-x)^3} + \frac{A_2(x)}{(1+2x)^2}\,.$$

where $A_1(x)$ is the Taylor polynomial of degree 2 about $x = 1$ for

$$f_1(x) = \frac{p(x)(1-x)^3}{q(x)} = \frac{p(x)}{(1+2x)^2}$$

and $A_2(x)$ is the Taylor polynomial of degree 1 for $f_2(x) = \dfrac{p(x)(1+2x)^2}{q(x)}$ about $x = -\frac{1}{2}$. Calculation gives

$$A_1(x) = -2 + 2(x-1) + (x-1)^2 = -2 - 2(1-x) + (x-1)^2$$

and

$$A_2(x) = 1 - 4(x + \frac{1}{2}) = 1 - 2(1+2x)\,.$$

This gives

$$\frac{p(x)}{q(x)} = -\frac{2}{(1-x)^3} - \frac{2}{(1-x)^2} + \frac{1}{1-x} + \frac{1}{(1+2x)^2} - \frac{2}{(1+2x)}\,. \tag{4.34}$$

## 4.4 The Generating Function Technique

The generating function technique we've just described allows us to solve the linear equation $L(s) = \psi$ when the generating function $F(x)$ for the input function is a rational function. For instance, when $\psi$ is a polynomial the method in the last sections allows us to write $F(x)$ as a sum of rational functions whose power series have the form given in (4.25). As an example, let us apply the generating function technique to solve

$$s_n = -s_{n-1} + 5s_{n-2} + s_{n-3} - 8s_{n-4} + 4s_{n-5}\,;\; S_0 = (9, -43, -13, -9, -4)^T\,,$$

which has characteristic polynomial $ch(x) = x^5 + x^4 - 5x^3 - x^2 + 8x - 4$, giving

$$\begin{bmatrix} 1 & 1 & -5 & -1 & 8 \\ 0 & 1 & 1 & -5 & -1 \\ 0 & 0 & 1 & 1 & -5 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix} \begin{pmatrix} 9 \\ -43 \\ -13 \\ -9 \\ -4 \end{pmatrix} = \begin{pmatrix} 8 \\ -7 \\ -2 \\ -13 \\ -4 \end{pmatrix}\,.$$

By (4.28) the generating function of $< s_n >$ is

$$S(x) = \frac{-4 - 13x - 2x^2 - 7x^3 + 8x^4}{1 + x - 5x^2 - x^3 + 8x^4 - 4x^5}$$

whose partial fraction decomposition is given in (4.34), and from (4.25)

$$s_n = \left(-2\binom{n+2}{2} - 2\binom{n+1}{1} + 1\right) + \left(\binom{n+1}{1} - 2\right)(-2)^n\,.$$

The rest of this section considers two nonlinear recurrences which arise in combinatorial problems and can be solved using generating functions.

### *4.4.1 The Catalan numbers*

For integer $n \geq 0$ consider a string of $n+1$ elements from a set with a binary operation. If the operation is not associative then the result of

$$a_0 a_1 \ldots a_n$$

is ambiguous until parentheses are inserted, and we can ask how many different arrangements of parentheses can be inserted in this string. This number is called the $n^{\text{th}}$ **Catalan number**. For instance, when $n = 2$ the possibilities are

$$a_0(a_1 a_2) \text{ and } (a_0 a_1)a_2\,,$$

giving $C_2 = 2$, and $C_3 = 5$ since the choices are

$$(a_0 a_1)(a_2 a_3)\,;\; a_0((a_1 a_2)a_3)\,;\; (a_0(a_1 a_2))a_3\,;\; a_0(a_1(a_2 a_3))\,;\; ((a_0 a_1)a_2)a_3\,.$$

Some authors, among them [16], claim this sequence was originally investigated by Euler [51]. In [21], Catalan considers this sequence and he points out that Lamé [80] already proved the sequence satisfies the nonlinear recurrence

$$\text{for all } n \geq 0\,, \;\; C_{n+1} = C_0C_n + C_1C_{n-1} + \cdots + C_nC_0\,,$$

with initial values $C_0 = 1\,,\, C_1 = 1$.

To see that $< C_n >$ does indeed satisfy this recurrence, consider any arrangement of parentheses in an $(n+2)$-string as defining a sequence of multiplications. The final multiplication is a product of a $(k+1)$-string and a $(n+1-k)$-string (for some $k$, $0 \leq k \leq n$) where the parentheses in the two factors can be arranged in $C_k$ and $C_{n-k}$ ways, respectively, and each yields a different final arrangement of parentheses. This means that for each $k \geq 0$ there is a total of $C_kC_{n-k}$ different arrangements of parentheses in which the first factor of the final multiplication is a $(k+1)$-string. Summing over all permitted $k$ $(k = 0, 1, \ldots, n)$ we obtain $C_{n+1} = C_0C_n + C_1C_{n-1} + \cdots + C_nC_0$, as required.

Let's use generating functions to show that $C_k = \dfrac{1}{k+1}\binom{2k}{k}$. Using the definition of multiplication of power series as in (3.12), the generating function $C(x)$ for the Catalan numbers satisfies

$$(C(x))^2 = \sum_{n\geq 0} (C_0C_n + C_1C_{n-1} + \cdots + C_nC_0)x^n = \sum_{n\geq 0} C_{n+1}x^n\,,$$

which gives

$$1 + xC(x)^2 = 1 + x\sum_{n\geq 0} C_{n+1}x^n = C_0 + \sum_{n\geq 1} C_nx^n = C(x)\,.$$

Then $z = C(x)$ is a power series solution to the quadratic equation $xz^2 - z + 1 = 0$ and Exercise 3.11 implies that $C(x)$ must be one of

$$C(x) = \frac{1 \pm \sqrt{1-4x}}{2x}\,.$$

By the generalized Binomial Theorem in (3.17) for $r = 1/2$, the power series of $\sqrt{1-4x}$ is $\sum_{n\geq 0}\binom{1/2}{n}(-4x)^n$ and we obtain

$$xC(x) = \frac{1 \pm \sum_{n\geq 0}\binom{1/2}{n}(-4x)^n}{2}\,.$$

Comparing the first coefficients of these power series, the negative sign is needed and

$$\begin{aligned} C(x) &= -\frac{1}{2}\sum_{n\geq 1}\binom{1/2}{n}(-4)^n x^{n-1} \\ &= -\frac{1}{2}\sum_{k\geq 0}(-1)^{k+1}\binom{1/2}{k+1}4^{k+1}x^k \\ &= \sum_{k\geq 0}(-1)^k\binom{1/2}{k+1}2^{2k+1}x^k, \end{aligned}$$

and

$$C_k = (-1)^k\binom{1/2}{k+1}2^{2k+1} = \frac{1}{k+1}\binom{2k}{k}$$

by Exercise 4.24.

### *4.4.2 Stirling Numbers of the Second Kind*

The second combinatorial problem that we consider is associated with partitioning a finite set, and our solution uses partial fractions. A **partition** of the $n$-set $S = \{1, 2, \ldots, n\}$ is a set of disjoint subsets (often called equivalence classes) whose union is $S$. For each pair of positive integers $k, n$ with $k \leq n$, the **Stirling number of the second kind** is denoted by $\left\{ {n \atop k} \right\}$ and is defined to be the number of partitions of a $n$-set in which there are $k$ classes. For instance, considering the partitions of a 4-set into 2 equivalence classes, there are 3 in which both equivalence classes have two elements (consider the possibilities for the class containing 1) and 4 which have a 1-set and a 3-set. This gives $\left\{ {4 \atop 2} \right\} = 7$.

First note that the number of such partitions in which $\{n\}$ is one of the equivalence classes is $\left\{ {n-1 \atop k-1} \right\}$. In the remaining partitions, $n$ is in an equivalence class with at least one other element and when $n$ is removed what remains is a partition of an $(n-1)$-set in which there are $k$ classes. There are $\left\{ {n-1 \atop k} \right\}$ such partitions, and $n$ might have originally been in any of the $k$ classes. Therefore, for all $1 \leq k \leq n$

$$\left\{ {n \atop k} \right\} = \left\{ {n-1 \atop k-1} \right\} + k\left\{ {n-1 \atop k} \right\}. \tag{4.35}$$

We extend this to all natural numbers $n, k$ by defining $\left\{ {n \atop k} \right\}$ to be zero when $k > n$ or $k = 0$ (except for $\left\{ {0 \atop 0} \right\} = 1$). For each fixed $k \geq 1$ define the generating function

$$S_k(x) = \sum_{n \geq 0} \left\{ {n \atop k} \right\} x^n = \sum_{n \geq k} \left\{ {n \atop k} \right\} x^n \,,$$

which from (4.35) becomes

$$S_k(x) = \sum_{n \geq k} \left\{ {n-1 \atop k-1} \right\} x^n + k \sum_{n \geq k} \left\{ {n-1 \atop k} \right\} x^n = x S_{k-1}(x) + kx S_k(x) \,.$$

This gives $S_k(x) = \dfrac{x}{1-kx} S_{k-1}(x)$, and inductively

$$S_k(x) = \frac{x^2 S_{k-2}(x)}{(1-kx)(1-(k-1)x)} = \ldots = \frac{x^k}{(1-kx)\ldots(1-x)}$$

since $S_0(x) = 1$. From the theory of partial fractions we know there exist $\alpha_j \in \mathbb{C}$ such that

$$\frac{1}{(1-x)(1-2x)\ldots(1-kx)} = \sum_{j=1}^{k} \frac{\alpha_j}{1-jx} \,,$$

which means that

$$\begin{aligned} S_k(x) &= x^k \sum_{j=1}^{k} \alpha_j \frac{1}{1-jx} = x^k \sum_{j=1}^{k} \alpha_j \sum_{n \geq 0} (jx)^n \\ &= \sum_{n \geq 0} \sum_{j=1}^{k} \alpha_j j^n x^{n+k} \\ &= \sum_{n \geq k} \Big( \sum_{j=1}^{k} \alpha_j j^{\,n-k} \Big) x^n \,, \end{aligned}$$

and combining this with $S_k(x) = \sum_{n \geq k} \left\{ {n \atop k} \right\} x^n$ gives

$$\left\{ {n \atop k} \right\} = \sum_{j=1}^{k} \alpha_j j^{\,n-k} \,.$$

For each $j = 1, \ldots, k$, $\alpha_j$ can be calculated (refer to Exercise 4.27) to be

$$\alpha_j = \frac{(-1)^{k-j} j^{\,k-1}}{(j-1)!(k-j)!} \quad (\text{for all } n, k > 0)\,,$$

and

$$\left\{ {n \atop k} \right\} = \sum_{j=1}^{k} \frac{(-1)^{k-j} j^{\,n-1}}{(j-1)!(k-j)!}\,.$$

## 4.5 Reversion of Generating Functions

In this section we explore another method for recovering the elements of a sequence from its generating function. (Be sure to review the material in Appendix B.) It's called the **Fourier Transform Method** and is a preview of the **Fast Fourier Transform** which we describe in detail in Chapter 9 (refer to Section 9.5). The basic idea is to replace the formal differentiation process by an appropriately selected weighted sum of some values of the generating function. Remember that in order to evaluate the generating function at $x = \alpha$, the power series must be more than simply a formal series: it must *converge* at $x = \alpha$. That is, its disk of convergence must contain a disk around each complex number at which the series is to be evaluated.

For any positive integer $n$, $\omega \in \mathbb{C}$ is called an **$\mathbf{n^{th}}$ root of unity** if $\omega^n = 1$. A **primitive $\mathbf{n^{th}}$ root of unity** is an $n^{\text{th}}$ root of unity which is not an $j^{\text{th}}$ root of unity for any $1 \le j < n$. For example, $\omega_2 = -1$ is the only a primitive second root of unity. For $n = 3$, $\omega_3 = \dfrac{-1 + i\sqrt{3}}{2}$ satisfies

$$\omega_3^2 = \frac{-1 - i\sqrt{3}}{2}\ ;\ \omega_3^3 = \omega_3^2 \cdot \omega_3 = \frac{-1 - i\sqrt{3}}{2} \cdot \frac{-1 + i\sqrt{3}}{2} = 1,$$

implying that $\omega_3$ is a primitive $3^{\text{rd}}$ root of unity. For general $n \ge 1$, since

$$\omega_n = e^{\frac{2\pi i}{n}} = \cos\left(\frac{2\pi}{n}\right) + i \sin\left(\frac{2\pi}{n}\right)$$

satisfies

$$\omega_n^k = e^{\frac{2k\pi i}{n}} = \cos\left(\frac{2k\pi}{n}\right) + i \sin\left(\frac{2k\pi}{n}\right),$$

then $\omega_n$ is always a primitive $n^{\text{th}}$ root of unity and is often called the **principal $\mathbf{n^{th}}$ root of unity**. When $\omega$ is any $n^{\text{th}}$ root of unity, then

$$0 = \omega^n - 1 = (\omega - 1)(\omega^{n-1} + \ldots + \omega + 1)$$

implies

$$\omega^{n-1} + \ldots + \omega + 1 = 0 \ \text{ provided } \omega \neq 1\,. \tag{4.36}$$

The next result shows how roots of unity can be used to recover the elements of *finite* sequences.

**Theorem 4.5.1.** *Let $< s_0, \ldots, s_{p-1} >$ be a finite sequence of length $p \geq 2$ with generating polynomial $S(x) = s_0 + s_1 x + \ldots + s_{p-1}x^{p-1}$. If $\omega$ is a primitive $p^{th}$ root of unity then $s_j$ can be computed by the formula*

$$s_j = \frac{1}{p}\sum_{n=0}^{p-1} \omega^{-jn} S(\omega^n)\,, \text{ for any } j = 1, \ldots, p-1.$$

*Proof.* For any fixed $j = 1, 2, \ldots, p-1$ we have

$$\omega^{-j}S(\omega) = s_0\omega^{-j} + s_1\omega^{1-j} + \ldots + s_{p-1}\omega^{p-1-j},$$

$$\omega^{-2j}S(\omega^2) = s_0\omega^{-2j} + s_1\omega^{2-2j} + \ldots + s_{p-1}\omega^{2(p-1)-2j},$$

$$\vdots$$

$$\omega^{-(p-1)j}S(\omega^{p-1}) = s_0\omega^{-(p-1)j} + s_1\omega^{p-1-(p-1)j} + \ldots + s_{p-1}\omega^{(p-1)^2-(p-1)^2 j}.$$

Adding these $p-1$ equations to $S(1) = s_0 + s_1 + \ldots + s_{p-1}$ we obtain

$$\sum_{n=0}^{p-1}\omega^{-jn}S(\omega^n) = s_0\sum_{n=0}^{p-1}\omega^{(0-j)n} + s_1\sum_{n=0}^{p-1}\omega^{(1-j)n} + \ldots + s_{p-1}\sum_{n=0}^{p-1}\omega^{(p-1-j)n},$$

where the coefficient of $s_i$ in this equation is $C_i = \sum_{n=0}^{p-1}\omega^{(i-j)n}$. When $i \neq j$ and $0 \leq i, j < p-1$, since $\omega$ is primitive then $\omega^{i-j} \neq 1$ and (4.36) implies $C_i = 0$. Therefore,

$$C_i = \begin{cases} 0 & \text{if } i \neq j \\ p & \text{if } i = j \end{cases},$$

and

$$\sum_{n=0}^{p-1}\omega^{-jn}S(\omega^n) = ps_j,$$

proving the theorem. □

This result can be easily extended to **periodic sequences**, infinite sequences $< s_n >$ formed by juxtaposing copies of a fixed finite sequence:

$$s_0, s_1, \ldots, s_{p-1}, s_0, s_1, \ldots, s_{p-1}, \ldots.$$

(The number $p$ is called the **period** of $< s_n >$ when it's the least integer with this property.) Since the first $p$ terms of such sequences $< s_n >$ are the

terms of the finite sequence $< s_0, s_1, \ldots, s_{p-1} >$, by Theorem 4.5.1 these terms can be calculated using

$$s_j = \frac{1}{p} \sum_{n=0}^{p-1} \omega^{-jn} S(\omega^n) \,.$$

The sequence $< s_n >$ is the infinite juxtaposition of these $p$ terms, and we've proved

**Corollary 4.5.2.** *Let $< s_n >$ be a periodic sequence with period $p$ and let $\omega$ be a primitive $p^{th}$ root of unity. If $S(x)$ is the generating function for $< s_0, s_1, \ldots, s_{p-1} >$ then for any $j = 0, 1, \ldots, p-1$*

$$s_{j+kp} = \frac{1}{p} \sum_{n=0}^{p-1} \omega^{-jn} S(\omega^n) \,, \text{ for all } k \,.$$

The technique used in these results can be extended to more general sequences, provided the generating function of the sequence converges in some neighborhood of $x = 0$. For this, we can no longer look at the series expansion as only a *formal* power series, but rather require it to satisfy the analytic property of convergence.

We first illustrate the technique by using it to recover $s_0$ and $s_1$. Suppose $S(x)$ converges on $|x| < R$. Then each of the series

$$S(x) = s_0 + s_1 x + \ldots + s_n x^n + \ldots$$

and

$$S(-x) = s_0 - s_1 x + \ldots + (-1)^n s_n x^n + \ldots$$

converges for every $|x| < R$, as does the sum

$$S(x) + S(-x) = 2(s_0 + s_2 x^2 + \ldots + s_{2n} x^{2n} + \ldots) = 2 f_2(x) \tag{4.37}$$

where $f_2(x)$ is the function $f_2(x) = \sum_{n \geq 0} s_{2n} x^{2n}$ discussed in Exercise 4.25(b). Therefore, (4.37) becomes

$$\lim_{x \to 0} \frac{S(x) + S(-x)}{2} = s_0 \,,$$

and we've recovered $s_0$! Also, considering the difference $S(x) - S(-x)$ and Exercise 4.25(c)-(d), we have

$$S(x) - S(-x) = 2x(s_1 + s_3 x^2 + \ldots + s_{2n+1} x^{2n} + \ldots) = 2x f_2(x) \,;$$

$$\lim_{x \to 0} \frac{S(x) - S(-x)}{2x} = \lim_{x \to 0} f_2(x) = s_1 \,, \tag{4.38}$$

the next element of the sequence.

To generalize this to a general reversion formula, notice that (4.38) can be rewritten as

$$\lim_{x\to 0}\frac{S(x)+\omega S(\omega x)}{2x} = s_1 \text{ when } \omega \text{ is the principal } 2^{\text{nd}} \text{ root of unity}\,.$$

Although this might seem like a complicated way to write (4.38), we'll see that this type of identity works in general.

As a warm-up for the general case, let $\omega$ be the principal $3^{\text{rd}}$ root of unity. Since

$$|x| = |\omega x| = |\omega^2 x| \quad \text{for all } x\,,$$

then for $x$ within $|x| < R$ the generating function $S(x)$ converges at all three of $x$ and $\omega x$ and $\omega^2 x$. Therefore, $\omega^3 = 1$ implies

$$S(\omega x) = s_0 + s_1\omega x + s_2\omega^2 x^2 + s_3 x^3 + s_4\omega x^4 + s_5\omega^2 x^5 + \dots\ ;$$

$$S(\omega^2 x) = s_0 + s_1\omega^2 x + s_2\omega x^2 + s_3 x^3 + s_4\omega^2 x^4 + s_5\omega x^5 + \dots\ .$$

Therefore,

$$\begin{aligned} S(x)+\omega S(\omega x)+\omega^2 S(\omega^2 x) &= s_0 + s_1 x + s_2 x^2 + s_3 x^3 + s_4 x^4 + s_5\omega x^5 + \dots \\ &\quad + s_0\omega + s_1\omega^2 x + s_2 x^2 + s_3\omega x^3 + s_4\omega^2 x^4 + s_5 x^5 + \dots \\ &\quad + s_0\omega^2 + s_1\omega x + s_2 x^2 + s_3\omega^2 x^3 + s_4\omega x^4 + s_5 x^5 + \dots \\ &= 0s_0 + 0xs_1 + 3s_2x^2 + 0s_3 + 0s_4x^4 + 3s_5x^5 + \dots\,, \end{aligned}$$

the last from (4.36). This can be reworded as

$$S(x)+\omega S(\omega x)+\omega^2 S(\omega^2 x) = 3x^2 f_3(x)\,, \tag{4.39}$$

where again from Exercise 4.25(d) we obtain

$$\lim_{x\to 0}\frac{S(x)+\omega S(\omega x)+\omega^2 S(\omega^2 x)}{3x^2} = s_2\,.$$

In Exercise 4.28 you show that the pattern in (4.39) generalizes to

$$S(x)+\omega S(\omega x)+\dots+\omega^{m-1}S(\omega^{m-1}x) = mx^{m-1}f_m(x)\,, \tag{4.40}$$

and from (4.40) and Exercise 4.25 we therefore obtain

$$\lim_{x\to 0}\frac{S(x)+\omega S(\omega x)+\dots+\omega^{m-1}S(\omega^{m-1}x)}{mx^{m-1}} = \lim_{x\to 0} f_m(x) = s_{m-1}\,,$$

which is

**The Reversion Formula**

*Suppose the generating function $S(x)$ for the sequence $< s_n >$ converges in some neighborhood of $x = 0$. Then for any integer $m \geq 2$,*

$$s_{m-1} = \lim_{x \to 0} \frac{1}{m x^{m-1}} \sum_{n=0}^{m-1} \omega_m^n S(\omega_m^n x), \tag{4.41}$$

*where $\omega$ is the principal $m^{th}$ root of unity.*

In the Reversion Formula $\omega$ is chosen to be the principal $m^{\text{th}}$ root of unity,

$$\omega = \cos\left(\frac{2\pi}{m}\right) + i \sin\left(\frac{2\pi}{m}\right),$$

because it allows us to use the very efficient Fast Fourier Transform (refer to Section 9.5) to evaluate the $S(\omega_m^n x)$.

As an example, we use the Reversion Formula to find the first two terms of the Fibonacci sequence. In Exercise 4.26 the radius of convergence for the Fibonacci generating function $F(x) = \dfrac{x}{1 - x - x^2}$ is shown to be $R = (\sqrt{5} - 1)/2$. We can therefore use $m = 1, 2$ in the Reversion Formula to obtain the first two terms of the sequence:

$$f_0 = \lim_{x \to 0} \frac{F(x) + F(-x)}{2} = \lim_{x \to 0} \frac{x^2}{(1 - x - x^2)(1 + x - x^2)} = 0,$$

and

$$f_1 = \lim_{x \to 0} \frac{F(x) - F(-x)}{2x} = \lim_{x \to 0} \frac{1 - x^2}{(1 - x - x^2)(1 + x - x^2)} = 1.$$

Of course for more complicated generating functions the limits in the Reversion Formula might be difficult to compute, and in such situations approximations to the limit give an estimate for the elements in the sequence.

## 4.6 Exercises

**Ex 4.1.** Using the alphabet $\{x, y\}$, find an exact formula for the number of $n$-strings with an even number of $x$'s.

In the next two exercises use the notation established in Section 4.1.

**Ex 4.2.** (a) Prove $s_{01}^{(n)} = s_{10}^{(n)}$ for all $n \geq 0$. Denote the common value by $t_n$; that is, $t_n = s_{01}^{(n)} = s_{10}^{(n)}$.

(b) Let $s_n = s_{00}^{(n)}$ and $t_n = s_{01}^{(n)} = s_{10}^{(n)}$. Use this and equations (4.4) and (4.10) through (4.13) to obtain

$$\begin{aligned} t_{n+1} &= 2t_n + 3t_{n-1} \; ; \; t_0 = 0, t_1 = 1 \, ; \\ s_{n+2} &= 3s_{n+1} + s_n - 3s_{n-1} \; ; \; s_0 = 1, s_1 = 1, s_2 = 3 \, ; \\ s_n &= s_{n-1} + 2t_{n-1} \; ; \; s_0 = 1, s_1 = 1 \, . \end{aligned}$$

**Ex 4.3.** Find exact formulas for $s_n$ and $t_n$.

**Ex 4.4.** Using the alphabet $\{x, y, z\}$, let $v_n$ be the number of $n$-strings in which the number of $y$'s is odd and the number of $z$'s is odd. Find an exact formula for $v_n$.

**Ex 4.5.** Use the technique of generating functions to find a formula for the $n^{\text{th}}$ term of the solution to the recurrence

$$s_0 = 0 \, , \, s_1 = 1 \quad , \quad s_n = 5s_{n-1} - 6s_{n-2}, \quad \text{for all } n \geq 2.$$

**Ex 4.6.** Let $f(x) = a_k x^k + \ldots + a_1 x + a_0$ be a nonconstant polynomial with complex coefficients and $a_0 \neq 0$. (For instance, the characteristic polynomial of (L) has these properties.) Let $f^R(x) = a_k + a_{k-1}x + \ldots + a_1 x^{k-1} + a_0 x^k$, the reciprocal polynomial of $f(x)$. Show that $\lambda$ is a non-zero root of $f(x) = 0$ if and only if $1/\lambda$ is a root of $f^R(x) = 0$.

**Ex 4.7.** Let $q(x) = (1 - \lambda x)(1 - \overline{\lambda} x)$ for nonreal $\lambda \in \mathbb{C}$.

(a) Show that for any $p(x) \in \mathbb{R}[x]$ with $\deg(p) < 2n$ there exists a unique $\beta(x) \in \mathbb{C}[x]$, $\deg(\beta) < n$, such that

$$\frac{p(x)}{(q(x))^n} = \frac{\beta(x)}{(1 - \lambda x)^n} + \frac{\overline{\beta(x)}}{(1 - \overline{\lambda} x)^n}, \tag{4.42}$$

where $\overline{\beta(x)}$ denotes the polynomial obtained from $\beta(x)$ by conjugating all coefficients.

(b) For any constant $\beta \in \mathbb{C}$

$$\frac{\beta}{1 - \lambda x} + \frac{\overline{\beta}}{1 - \overline{\lambda} x} = 2 \sum_{j \geq 0} Re(\beta \lambda^j) x^j, \tag{4.43}$$

where $Re(\beta\lambda^j)$ denotes the real part of the complex number $\beta\lambda^j$.

**Ex 4.8.** . Let $a(x), b(x) \in \mathbb{C}[x]$ be two non-zero polynomials. Show any common divisor of $a(x)$ and $b(x)$ divides the last non-zero remainder $R_n(x)$ in the Euclidean Algorithm applied to $a(x)$ and $b(x)$. If $\alpha$ is the leading coefficient of $R_n(x)$ then $\alpha^{-1} R_n(x)$ is the **greatest common divisor of a(x) and b(x)**.

**Ex 4.9.** As established in Chapter 3, any operator of the form $L = I - c_1\sigma - \ldots - c_k\sigma^k$ is an invertible operator on $\mathcal{S}^+$ and is also invertible as a formal power series. At the end of the chapter we claimed that the coefficients of $L^{-1} = ch^R(\sigma)^{-1} = \sum_{n\geq 0} a_n\sigma^n$ satisfy

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \ldots + c_k a_{n-k} \text{ for all } n \geq k\,.$$

Use results from generating functions to show this.

**Ex 4.10.** If $L(\sigma)$ is the shift operator associated with the initial value problem

$$s_0 = 0\,,\ s_1 = 1\,;\ s_n = 4s_{n-1} - 4s_{n-2}\,,$$

show that its inverse is the operator

$$L^{-1} = \sum_{n\geq 0} (n+1)2^n\sigma^n\,.$$

**Ex 4.11.** Verify that partial fraction expansion of the generating function for

$$s_0 = 0\,,\ s_1 = 1\,,\ s_n = 4s_{n-1} - 4s_n + 3^n(n-1)$$

is

$$\frac{19}{2}\frac{1}{(1-2x)^2} + \frac{71}{2}\frac{1}{1-2x} + \frac{9}{(1-3x)^2} - \frac{54}{1-3x}\,.$$

**Ex 4.12.** (a) Find the generating function of $s_n = n^2$.

(b) Show that the generating function of the sequence $s_n = 1^2 + 2^2 + \ldots + n^2$ is

$$\frac{2}{(1-x)^4} - \frac{3x}{(1-x)^3} - \frac{2}{(1-x)^2}\,.$$

**Ex 4.13.** (a) If $S(x)$ is the generating function for the sequence $< s_n >$ show that the the generating function for the partial sums $\sum_{j=0}^{n} s_j$ is $S(x)/(1-x)$.

(b) Use part(a) to prove the Fibonacci identity

$$F_0 + F_1 + \ldots + F_n = F_{n+2} - 1\,.$$

**Ex 4.14 (L'Hôpital's Rule).** Let $p(x), q(x) \in \mathbb{C}[x]$ be polynomials.

(a) If $\alpha \in \mathbb{C}$ such that $q(\alpha) = 0$, show that $q(x) = (x-\alpha)q_1(x)$ for some $q_1(x) \in \mathbb{C}[x]$ and $q'(\alpha) = q_1(\alpha)$.

(b) Suppose $p(\alpha) = 0$ and that $\alpha$ is a *simple* zero of $q(x)$. Show that

$$\lim_{x\to\alpha} \frac{p(x)}{q(x)} = \lim_{x\to\alpha} \frac{p'(x)}{q'(x)}.$$

**Ex 4.15.** Let $s_0 = 5\,,\ s_1 = 13,\ s_n = -4s_{n-1} + 5s_{n-2}$ for all $n \geq 2$. Use the generating function method to find an exact formula for the $n^{\text{th}}$ term of $< s_n >$.

**Ex 4.16.** Let $s_0 = -1$ , $s_1 = 2$ , $s_2 = 14$, $s_n = s_{n-1} - 4s_{n-2} + 4s_{n-3}$ for all $n \geq 3$. Use the generating function method to find a formula for the $n^{\text{th}}$ term of $< s_n >$.

**Ex 4.17.** Expand $\dfrac{4x^2 - 2x - 1}{(1+x)(1-x-3x^2)}$ into a power series.

**Ex 4.18.** Find the sequence with generating function $\dfrac{1}{1-2x} + \dfrac{1}{(1-3x)^3}$.

**Ex 4.19.** Use the generating function technique to show that

$$s_n = \begin{cases} 2^n & \text{if } n \text{ is even} \\ 2^n + 2 & \text{if } n \equiv 1 \pmod 4 \\ 2^n - 2 & \text{if } n \equiv 3 \pmod 4 \end{cases}$$

is the solution to

$$(s_0, s_1, s_2) = (1, 4, 4), \quad s_{n+3} = 2s_{n+2} - s_{n+1} + 2s_n.$$

**Ex 4.20.** (a) Use the technique of generating functions to solve

$$s_0 = 1\,,\ s1 = 0\,, \quad s_n = 4s_{n-1} - 4s_n + 3^n(n-1)\,.$$

(Note that the equation is quite in the form of (L).)

(b) Find all solutions to

$$s_n = 4s_{n-1} - 4s_n + 3^n(n-1)\,.$$

**Ex 4.21.** We consider a nonhomogeneous recurrence with characteristic polynomial $ch(x)$ and forcing function $\psi(n) = \lambda^n p(n)$ for some polynomial $p(x)$ of degree $m \geq 1$.

(a) Show there exists a polynomial $Q(x)$ with $\deg(Q) \leq m$ such that the generating function of every initial value problem for this recurrence has the form

$$S(x) = \frac{d(x)(1-\lambda x)^{m+1} + x^k Q(x)}{ch^R(x)(1-\lambda x)^{m+1}} \tag{4.44}$$

for some polynomial $d(x)$ with $\deg(d) < \deg(ch)$.

(b) Show that for any polynomial $d(x)$ with $\deg(d) < \deg(ch)$ with $\deg(d) < \deg(ch)$ there exists an initial value problem whose generating function has the form given in (4.44).

**Ex 4.22.** Let $s_n = \alpha\lambda_0^n + a_1(n)\lambda_1^n + \ldots + a_{k-1}(n)\lambda_{k-1}^n$ be a solution to the $k^{\text{th}}$ order recurrence

$$s_n = c_1 s_{n-1} + c_2 s_{n-2} + \ldots + c_k s_{n-k}$$

and suppose $\lambda_0$ is a simple eigenvalue of the recurrence. Then $s_n = \alpha\lambda_0^n + a_1(n)\lambda_1^n + \ldots + a_{k-1}(n)\lambda_{k-1}^n$ where $\alpha$ is a constant and each $\alpha_i(n)$ is a polynomial in $n$. Show

$$\alpha = \lim_{x\to\lambda_0} \frac{(d_0x^{k-1} + d_1x^{k-2} + \ldots + d_{k-1})(x-\lambda_0)}{ch(x)}$$

for $d_i = s_i - c_1s_{i-1} - \ldots - c_{i-1}s_0$.

**Ex 4.23 (Generalized Fibonacci sequence).** For $k \geq 2$ define the sequence $< f_n >$ for all $n \geq 0$ by

$$f_0 = f_1 = \ldots = f_{k-2} = 0\, f_{k-1} = 1 \quad \text{and} \quad f_n = f_{n-1} + f_{n-2} + \cdots + f_{n-k}\,.$$

(a) Show that for all $x \neq 1$

$$ch(x) = \frac{x^k(x-2)+1}{x-1}\,.$$

(b) Use the last exercise and part (a) to show that if $\lambda_0$ is the largest eigenvalue of the recurrence (later we'll prove it is the *only* positive eigenvalue) then $f_n = \alpha\lambda_0^n + d_n$ where $d_n$ has no $\lambda_0^n$ component and $\alpha$ equals

$$\alpha = \frac{\lambda_0 - 1}{\lambda_0[(k+1)\lambda_0 - 2k]}\,.$$

**Ex 4.24.** Show that for all positive integers $k$,

$$\binom{1/2}{k+1} 2^{2k+1} = \frac{(-1)^k}{k+1}\binom{2k}{k}\,.$$

For the next two exercises, refer to the information on the disk of convergence given in Appendix B.

**Ex 4.25.** Let $\gamma(x) = \sum_{n\geq 0} a_n x^n$ be a fixed power series with disk of convergence $|x| \leq R$.

(a) For any infinite strictly increasing sequence $< n_i >$ of positive integers, consider the power series $\gamma_1(x) = \sum_{i\geq 0} a_{n_i} x^{n_i - n_0}$. If $\alpha \in \mathbb{C}$ such that $|\alpha| < R$ , show that the power series $\gamma_1(x)$ also converges at $x = \alpha$.

(b) For any $|x| < R$, show that

$$\lim_{x\to 0} \frac{\gamma(x) + \gamma(-x)}{2} = a_0\,.$$

(c) For any positive integer $m$ use part (a) to define the complex-valued function $f_m$ on the open disk $|x| < R$ by

$$f_m(x) = \sum_{i\geq 0} a_{(i+1)m-1} x^{im}\,.$$

(d) Show that $\lim_{x\to 0} \dfrac{f_m(x)}{mx^m} = a_{m-1}$.

**Ex 4.26.** Show that $(\sqrt{5}-1)/2$ is the radius of convergence of the Fibonacci generating function.

**Ex 4.27.** For any $k \geq 1$, show the partial fraction decomposition of

$$\frac{1}{(1-x)(1-2x)\dots(1-kx)} = \frac{\alpha_1}{1-x} + \dots + \frac{\alpha_k}{1-kx}$$

has

$$\alpha_j = \frac{(-1)^{k-j} j^{k-1}}{(j-1)!(k-j)!}\,.$$

**Ex 4.28.** Verify the pattern in (4.40) and thereby construct an argument to convince yourself that the Reversion Formula holds for general $m \geq 2$.

**Ex 4.29.** Let $S(x) = \dfrac{2}{1-4x^2}$.

(a) Use the Fourier Transform Method to calculate the first three elements of the sequence with generating function $S(x)$.

(b) Use the partial fraction decomposition of $S(x)$ to verify you computed the first three terms correctly.

**Ex 4.30.** For any sequence $< a_n >$, define the **exponential generating function** by

$$\mathcal{E}(a_n) = \sum_{n\geq 0} \frac{a_n}{n!} x^n\,.$$

In particular, $\mathcal{E}((-1)^n) = e^{-x}$ and $\mathcal{E}(n!) = \dfrac{1}{1-x}$. Show that

(a) For any positive integer $m$, $\mathcal{E}(a_{n+m}) = \mathcal{D}^m(\mathcal{E}(a_n))$.

(b) For any polynomial $p(x)$, $\mathcal{E}(p(n)a_n) = p(x\mathcal{D})(\mathcal{E}(a_n))$.

(c) $\mathcal{E}(a_n)\mathcal{E}(b_n) = \mathcal{E}(\sum_{k=0}^{n} \binom{n}{k} a_k b_{n-k})$.

**Ex 4.31.** For the problem

$$a_{n+1} = (n+1)a_n + (-1)^n \text{ with } a_0 = 1$$

use the recurrence to show directly that $\mathcal{E}(a_n) = \mathcal{E}((-1)^n)\mathcal{E}(n!)$, and from this use a result in the last problem to prove that $a_n = \displaystyle\sum_{i=0}^{n} \frac{n!}{i!}(-1)^n$.

**Ex 4.32.** A **derangement** of the finite set $\{1, 2, \dots, n\}$ is a permutation of the set in which every element is moved from its original position. Let $d_n$ equal the number of derangements of the set $\{1, 2, \dots, n\}$. For instance, $d_1 = 0$, $d_2 = 1$, $d_3 = 2$. Let $d_0 = 1$.

(a) Show $d_n$ equals the number of derangements of the set $\{1, 2, \ldots, n+1\}$ in which $n+1$ is in the first position and 1 is not in the last place. Use this idea to show that $< d_n >$ satisfies the second-order recurrence

$$d_{n+1} = n(d_n + d_{n-1}) .$$

(b) Setting $b_{n+1} = d_{n+1} - (n+1)d_n$, show that $b_{n+1} + b_n = 0$ holds for all $n \geq 0$. That is, $< d_n >$ satisfies the first-order recurrence in the last problem.

(c) Writing the probability that a permutation is a derangement as a finite sum, show that this probability converges to $1/e$.

# 5
# Nonnegative Difference Equations

In this chapter we consider **nonnegative difference equations**. Our prototype, the Fibonacci sequence

$$f_n = f_{n-1} + f_{n-2} \quad ; \quad f_0 = 0\,,\, f_1 = 1$$

is a nonnegative system because all coefficients and all initial values are nonnegative. We've shown earlier that elements of the Fibonacci sequence have the form

$$f_n = \frac{1}{\sqrt{5}}(\lambda_0^n - \lambda_1^n)$$

where $\lambda_0 = (1+\sqrt{5})/2$ and $\lambda_1 = (1-\sqrt{5})/2$ are the roots of the characteristic polynomial, $ch(x) = x^2 - x - 1$. Although $\lambda_0$ and $\lambda_1$ are irrational, $f_n$ is an integer and in fact

$$f_n = Round(\lambda_0^n/\sqrt{5}) \text{ for all } n \geq 0,$$

where $Round(X)$ is the function which returns the integer nearest to $X$. (Refer to Chapter 1.) This implies the asymptotic size $f_n = \Theta(\lambda_0^n)$ where $\lambda_0$ is the positive eigenvalue of the recurrence. We'd like to generalize this example and discover which properties hold for solutions of the generalized problem. For example, for integer $k \geq 3$ we can consider the **generalized Fibonacci sequence** which satisfies the $k^{\text{th}}$-order recurrence

$$f_n^{(k)} = f_{n-1}^{(k)} + f_{n-2}^{(k)} + \cdots + f_{n-k}^{(k)}$$

with initial values

$$f_0^{(k)} = 0, f_1^{(k)} = 1, f_2^{(k)} = 2, f_3^{(k)} = 2^2, \cdots = f_{k-1}^{(k)} = 2^{k-2}\,.$$

We might guess that $f_n^{(k)} = \Theta(\lambda_0^n)$ where $\lambda_0 = \lambda_0(k)$ is some nonnegative number which depends on $k$. A further generalization is

$$s_n = c_1 s_{n-1} + c_2 s_{n-2} + \cdots + c_k s_{n-k}$$

where the $c_i$'s are nonnegative and all initial conditions $s_0, \ldots, s_{k-1}$ are nonnegative. (We also assume that at least one of these initial conditions is positive, or else the solution is the zero sequence.) We will investigate whether any additional conditions are needed to ensure the existence of a nonnegative number $\lambda_0$ such that $s_n = \Theta(\lambda_0^n)$. For the Fibonacci numbers it is true that $f_n = \mathit{Round}(\lambda_0^n/\sqrt{5})$ and we want to know what other integer sequences have the property that there exists an $\alpha > 0$ so that $s_n = \mathit{Round}(\alpha\lambda_0^n)$ holds.

We also consider the asymptotic size of solutions to *nonhomogeneous* equations

$$\text{(NN)} \qquad s_n = c_1 s_{n-1} + c_2 s_{n-2} + \cdots + c_k s_{n-k} + g(n)$$

where the $c_i$'s are nonnegative, the initial conditions are nonnegative, and $g(n)$ are functions with nonnegative values. Here we ask when $s_n = \Theta(g(n))$ or $s_n = \Theta(\lambda_0^n)$ for some $\lambda_0$.

As in previous chapters, much information about the recurrence is encoded in its characteristic polynomial which is analyzed in the next section.

## 5.1 Nonnegative Polynomials

The purpose of this section is to study polynomials of the form

$$(5.1)\quad p(x) = x^k - c_1 x^{k-1} - \cdots - c_{k-1}x - c_k \ , \quad \text{with all } c_i \geq 0 \text{ and } c_k > 0\,.$$

These polynomials are called **nonnegative**, and are the characteristic polynomials of nonnegative linear recurrences, that is, recurrences which have the form (NN). The study of these polynomials takes us through some classical areas of mathematics which are often omitted from an undergraduate education but which are currently experiencing at least a small revival because of applications to computing and robotics as well as to other areas.

### *5.1.1 The Dominant Root*

From the Fundamental Theorem of Algebra (refer to Appendix B) we know that any polynomial $p(x)$ has $\deg(p)$ complex roots when the roots are counted according to multiplicity. It's difficult to say how many of these roots are real. One of the oldest results in this direction is Descartes' Rule of Signs, which relates the number of roots of a polynomial with *real* coefficients to the number of sign changes among its coefficients. For us, a

**sign change** occurs in the polynomial when some $c_i$ is positive and the next non-zero coefficient is negative, or vice versa. For instance, $x^2-2x+1$ has two sign changes, and $x^{30}+x^{20}-5x^{10}-7$ has just one sign change. Descartes' Rule bounds the number of positive real roots of a polynomial with *real* coefficients by its number of sign changes. Here's the statement of the rule and a proof.

**Theorem 5.1.1 (Descartes' Rule of Signs).** *The number of positive real roots (counted according to multiplicity) of a polynomial with real coefficients is no more than the number of sign changes among its coefficients.*

*Proof.* We proceed by induction on both the number of sign changes $n$ in the polynomial *and* its degree $k$. When there are no sign changes $p(x) = c_0x^k+\ldots+c_k$ has the same sign at every positive real number and therefore has no positive root. Because of this we may assume $n > 0$. The induction hypothesis is that any polynomial with fewer than $n$ sign changes or with exactly $n$ sign changes but whose degree is less than $k$ has at most $n$ positive (real) roots.

If $c_k = 0$, then $p(x) = xP(x)$ for some polynomial $P(x)$ which also has $n$ sign changes and whose degree is $k-1$. The polynomial $P(x)$ is covered by our induction hypothesis and its number of positive roots is therefore at most $n$. Since $p(x) = xP(x)$, every non-zero root of $p(x)$ is a root of $P(x)$. This gives the conclusion for such $p(x)$ and allows us to assume that $c_k \neq 0$.

Without loss of generality we may assume $p(x)$ has at least one positive root, and set $\lambda_0$ to be the least of its positive roots. Consider the derivative of $p(x)$,

$$p'(x) = kc_0x^{k-1} + (k-1)c_1x^{k-2} + \cdots + 2c_{k-2}x + c_{k-1}\,.$$

The number of sign changes in $p'(x)$ either equals the number for $p(x)$ or is one fewer, depending on whether $c_{k-1}$ and $c_k$ have the same or opposite sign. By the continuity of $p(x)$ and $p'(x)$, between every two different consecutive real roots of $p(x)$ there must be a local extremum of $p(x)$ and so at least one root of $p'(x)$. Together with Exercise 5.1 this implies that $p(x)$ can have at most one more positive root than $p'(x)$ on the interval $[\lambda_0, \infty)$. The proof is completed by showing that $p'(x)$ has at most $n-1$ roots in this interval since then the number of positive roots of $p(x)$ is at most $n$ since $\lambda_0$ was chosen as its least positive root.

If $c_k$ and $c_{k-1}$ have opposite signs, then $p'(x)$ has $n-1$ sign changes. Our induction hypothesis therefore limits the number of positive real roots of $p'(x)$ to $n-1$, as claimed. If $c_k$ and $c_{k-1}$ have the same sign, both $p(x)$ and $p'(x)$ have $n$ sign changes and by induction $p'(x)$ has at most $n$ positive roots. We will show that $p'(x)$ has at least one root in the interval $(0, \lambda_0)$. This follows from a bit of calculus. First considering the case when $c_k > 0$, the values of both $p(x)$ and $p'(x)$ on any sufficiently small interval

containing zero are dominated by their constant terms and therefore are both positive. So $p(x)$ is increasing and positive at $x = 0$ but must decrease to $p(\lambda_0) = 0$, and $p(x)$ has a relative maximum (which is a root of $p'(x)$) on the interval $(0, \lambda_0)$. When $c_k < 0$, this argument can be applied to $q(x) = -p(x)$ to obtain a relative minimum for $p(x)$. In either case, $p(x)$ has a relative extremum on $(0, \lambda_0)$ and continuity yields a root of $p'(x)$ in the interval $(0, \lambda_0)$ as claimed. □

In 1637 Descartes stated this rule of signs without proof in his famous book *La Géométrie* [130]. During the next two centuries several people proved and refined the rule. Among these was Carl Friedrich Gauss [61] who in 1828 provided the additional information that the difference between the number of positive roots and the number of sign changes is always an even number. **Sturm's Theorem** (*Mémoire sur la résolution des équations numériques*, published in 1829) is another classic theorem in this vein. In its simplest version this theorem yields an algorithm for counting the number of real roots of $P(x)$ in any interval. More complicated versions allow the determination of the number of roots of a polynomial $P(x)$ subject to sign constraints on another polynomial $Q(x)$. These theorems fell into relative obscurity until revived by Tarski in 1940 [138] to prove an abstract result in mathematical logic.

Nonnegative polynomials, $p(x) = x^k - c_1 x^{k-1} - \ldots c_{k-1}x - c_k$, have exactly one sign change and so $p(x)$ can have *at most* one positive real root. On the other hand, since $p(0) = -c_k < 0$ and $\lim_{t \to +\infty} p(x) = +\infty$ hold, the Intermediate Value Theorem implies that $p(x)$ has *at least* one positive real root. Therefore,

**Corollary 5.1.2.** *A nonnegative polynomial has exactly one positive real root and it is a simple root.*

We next show that among all roots of a nonnegative polynomial its sole positive root has the largest complex modulus.

**Theorem 5.1.3.** *If $\lambda_0$ is the positive real root of a nonnegative polynomial $p(x)$ then $\lambda_0$ is a* **dominant root**, *that is, any other root $\lambda \in \mathbb{C}$ satisfies $|\lambda| \leq \lambda_0$.*

*Proof.* If $\lambda$ is any root of $p(x)$, then

$$\lambda^k = c_1\lambda^{k-1} + c_2\lambda^{k-2} + \cdots + c_k\,.$$

The triangle inequality then yields

$$\begin{aligned} |\lambda|^k = |\lambda^k| &\leq |c_1\lambda^{k-1}| + |c_2\lambda^{k-2}| + \cdots + |c_k| \\ &= c_1|\lambda|^{k-1} + c_2|\lambda|^{k-2} + \cdots + c_k\,. \end{aligned}$$

We conclude that

$$p(|\lambda|) = |\lambda|^k - c_1|\lambda|^{k-1} - \cdots - c_k \leq 0\,,$$

and from Exercise 5.2 this implies $|\lambda| \leq \lambda_0$. □

Do nonnegative polynomials ever have other roots of modulus $\lambda_0$? Not only is it possible to show that the answer is *yes*, but we can actually find all roots with modulus $|\lambda_0|$ without doing much work at all. This is described in the next theorem. We make use of the $m^{\text{th}}$ **roots of unity** (refer to Appendix B), that is, complex numbers $\zeta$ such that $\zeta^m = 1$. The following result gives the relationship among the roots of maximum modulus.

**Theorem 5.1.4.** *Let $\lambda_0$ be the positive real root of the nonnegative polynomial $p(x) = x^k - c_1 x^{k-1} - \cdots - c_k$. Let $g = \gcd\{i : c_i > 0\}$ be the greatest common divisor of the set of indices of non-zero coefficients, which is called the* **index of imprimitivity**. *Then $p(x)$ has exactly $g$ roots of modulus $\lambda_0$, the $g$ complex numbers of the form $\lambda_0\zeta$ where $\zeta$ is a primitive $g^{th}$ root of unity.*

We will consider a few examples before proving this result. For a polynomial of the form $p_1(x) = x^k - 1$ we have $g = k$ and the roots of the polynomial *are* the $g^{\text{th}}$ roots of unity. For example, $x^4 - 1$ has the four roots $\pm 1, \pm i$ which can be written as $i^0, i^1, i^2, i^3$ since $i$ is a primitive $4^{\text{th}}$ root of unity. Next, let's look at the polynomial $p_2(x) = x^4 - x^2 - 1 = (x^2)^2 - (x^2) - 1$. To find its positive real root, we apply the quadratic formula:

$$x^2 = \frac{1 \pm \sqrt{5}}{2},$$

which leads to its positive root

$$\lambda_0 = \sqrt{\frac{1 + \sqrt{5}}{2}}.$$

Since the only non-zero coefficients are $c_2$ and $c_4$, then $g = 2$, $\pm 1$ are the $g^{th}$ roots of unity, and the theorem correctly says that there are exactly two roots of modulus $\lambda_0$, namely $\pm\lambda_0$. For a more interesting example, consider the nonnegative polynomial $p_3(x) = x^9 - 3x^6 - 7$. Even though we don't know the value of $\lambda_0$, from the theorem we know that it has exactly three roots of modulus $\lambda_0$ since $\gcd\{6, 9\} = 3$.

*Proof of Theorem 5.1.4.* If $\zeta \in \mathbb{C}$ is both a square root *and* a cube root of unity, $\zeta$ is in the set $\{-1, 1\}$ and also in the set $\{1, e^{2\pi/3}, e^{4\pi/3}\}$ which implies that $\zeta = 1$. This observation is generalized in Exercise 5.6 where you show that if $g$ is the greatest common divisor of a finite set $G$ of integers, then $\zeta$ is a $g^{\text{th}}$ root of unity if and only if $\zeta$ is an $i^{\text{th}}$ root of unity for every $i \in G$. We apply this here with $G = \{i : c_i > 0\}$, the set of indices of the negative coefficients of $p(x)$. Since $\lambda_0$ is the dominant root of $p(x)$ it suffices to prove $\omega$ is a root of $p(x)$ with $|\omega| = \lambda_0$ if and only if $\zeta = \omega/\lambda_0$ is an $i^{\text{th}}$ root of unity for all $i \in G$

If $\zeta$ is a complex number with $\zeta^i = 1$ for all $i \in G$, then

$$\begin{aligned}\zeta^{-k} \cdot p(\lambda_0 \zeta) &= \lambda_0^k - \sum_{i \in G} c_i \lambda_0^{k-i} \zeta^{-i} \\ &= \lambda_0^k - \sum_{i \in G} c_i \lambda_0^{k-i} = p(\lambda_0) = 0\,,\end{aligned}$$

and $\omega = \lambda_0 \zeta$ is indeed a root of $p(x)$. On the other hand, suppose $\omega = \zeta \lambda_0$ is a root of $p(x)$ with $|\omega| = \lambda_0$. Then

$$0 = p(\zeta \lambda_0) = \zeta^k \lambda_0^k - c_1 \zeta^{k-1} \lambda_0^{k-1} - \cdots - c_k\,;$$

$$\lambda_0^k = \sum_{i \in G} c_i \lambda_0^{k-i} \zeta^{-i}\,.$$

Now $p(\lambda_0) = 0$ implies

$$\lambda_0^k = c_1 \lambda_0^{k-1} + \cdots + c_k = \sum_{i \in G} c_i \lambda_0^{k-i}\,,$$

and subtracting these two equations gives

$$0 = \sum_{i \in G} c_i \lambda_0^{k-i} (1 - \zeta^{-i})\,. \tag{5.2}$$

Notice that since each $\zeta^{-i}$ lies on the unit circle, the real part of each $1 - \zeta^{-i}$ is $1 - cos(-i\theta)$ and must be nonnegative. Taking the real part of both sides of (5.2), the left-hand side is 0 and the right-hand side contains only nonnegative terms since all $\lambda_0^{k-i}$ are positive (remember that $\lambda_0 > 0$). Therefore, the real part of $1 - \zeta^{-i}$ must equal zero for all $i \in G$. Again using the fact that each $\zeta^{-i}$ lies on the unit circle, we note that the real part of $1 - \zeta^{-i}$ is zero if and only if $\zeta^{-i} = 1$, and we've proved $\zeta$ must be an $i^{\text{th}}$ root of unity for all $i \in G$. □

Returning to the examples we considered before the proof, the fact that the index of imprimitivity for the polynomials $p_2(x)$ and $p_3(x)$ doesn't equal 1 allows us to write $p_2(x) = (x^2)^2 - (x^2) - 1$ and $p_3(x) = (x^3)^3 - 3(x^3)^2 - 7$, displaying a sort of periodicity. Because of this, we call a nonnegative polynomial **periodic** when its index of imprimitivity does not equal one, and otherwise we call it **primitive (or aperiodic)**. Combining Theorem 5.1.3 with Theorem 5.1.4,

**Corollary 5.1.5.** *Any primitive nonnegative polynomial has a unique root of largest modulus, which is often referred to as the* **strictly dominant root**.

## 5.2 Homogeneous Nonnegative Recurrences

In this section we consider the asymptotic size of solutions to *homogeneous* nonnegative recurrences, which can be written in the form:

(HNN) $s_n = c_1 s_{n-1} + c_2 s_{n-2} + \cdots + c_k s_{n-k}$ where all $c_i \geq 0$ and $c_k \neq 0$.

Since the characteristic polynomial of this equation is a nonnegative polynomial, the results of the previous section can be applied, and we let $\lambda_0$ be the dominant eigenvalue of the recurrence. Remember also that we're assuming the initial conditions are nonnegative.

An example of a simple *periodic* nonnegative difference equation is

$$s_n = 4s_{n-2}$$

which has periodic characteristic polynomial $ch(x) = x^2 - 4$ and dominant root $\lambda_0 = 2$. With initial conditions $s_0 = 1 = s_1$ the solution is

$$s_n = 4^{\lfloor n/2 \rfloor},$$

and $s_n = \Theta(2^n) = \Theta(\lambda_0^n)$. When the initial conditions are $s_0 = 1$, $s_1 = 0$ the solution is

$$s_n = \begin{cases} 4^{n/2} & \text{when } n \text{ is even} \\ 0 & \text{when } n \text{ is odd} \end{cases}$$

and $s_n = O(2^n)$ but $s_n \neq \Theta(\lambda_0^n)$ since every odd position in the sequence is 0.

The next theorem and its corollary elucidate the difference between periodic and primitive equations.

**Theorem 5.2.1.** *Let $\lambda_0$ be the dominant eigenvalue of the homogeneous nonnegative recurrence (HNN) with nonnegative initial values $s_0, \ldots, s_{k-1}$. If $< s_n >$ is the solution then $s_n = O(\lambda_0^n)$. If in addition there are $k$ consecutive elements in $< s_n >$ which are positive, then $s_n = \Theta(\lambda_0^n)$.*

*Proof.* For any natural number $N$ we define the finite set of real numbers

$$S_N = \{s_i/\lambda_0^i \,:\, N \leq i < N + k\},$$

and set $\alpha_N = \min(S_N)$, $\beta_N = \max(S_N)$. Then

$$\alpha_N \lambda_0^i \leq s_i \leq \beta_N \lambda_0^i \text{ for all } N \leq i < N + k. \tag{5.3}$$

We next prove

$$s_n \leq \beta_0 \lambda_0^n \text{ for all } n \geq 0 \tag{5.4}$$

by induction on $n$. By construction, (5.4) holds for all $0 \leq n < k$. If (5.4) holds for all $0 \leq n < K$ then

$$s_K = \sum_{i=1}^{k} c_i s_{K-i} \leq \beta_0 \sum_{i=1}^{k} c_i \lambda_0^{K-i} = \beta_0 \lambda_0^K,$$

since $\sum_{i=1}^{k} c_i \lambda_0^{K-i} = \lambda_0^K$. Therefore, $s_K \leq \beta_0 \lambda_0^K$, and we've proved (5.4), and so $s_n = O(\lambda_0^n)$. We can obtain $s_n \geq \alpha_0 \lambda_0^n$ by a similar argument, but with only nonnegativity of the initial conditions $\alpha_0$ might be zero. By hypothesis we know there are $k$ consecutive positive values, say $s_N, \ldots, s_{N+k-1}$ and then $\alpha_N > 0$ holds. Using the above argument beginning with $n = N$ instead of $n = 0$, gives

$$s_{N+k} = \sum_{i=0}^{k-1} c_i s_{N+i} \geq \alpha_N \sum_{i=0}^{k-1} c_i \lambda_0^{N+i} = \alpha_N \lambda_0^{N+k}$$

and by induction $s_n \geq \alpha_N \lambda_0^n$ for all $n \geq N$, and so $s_n = \Theta(\lambda_0^n)$. □

For example, the Fibonacci recurrence is a homogeneous nonnegative recurrence in which all terms after the first are positive. Therefore this result says that $f_n = \Theta(\lambda_0^n)$ where $\lambda_0 = \frac{1+\sqrt{5}}{2}$ is the unique positive eigenvalue. Another simple example is

$$s_n = s_{n-1} + 2s_{n-2} \text{ with } s_0 = 1\, s_1 = 2.$$

Since $ch(x) = x^2 - x - 2 = (x+1)(x-2)$ and the first two conditions are positive, the solution is $\Theta(2^n)$. In Exercise 5.3 you consider this recurrence under various initial conditions, including when some initial conditions are negative.

**Corollary 5.2.2.** *Let $< s_n >$ be the solution to a homogeneous nonnegative recurrence with nonnegative initial conditions where not all initial conditions are zero. If the characteristic polynomial is primitive then $< s_n >$ is eventually positive and $s_n = \Theta(\lambda_0^n)$.*

*Proof.* From the preceding result it suffices to prove the sequence is eventually positive. Let $s_n = c_1 s_{n-1} + \ldots + c_k s_{n-k}$ and we may assume that $s_0 > 0$. Setting $G = \{i : c_i > 0\}$, we know that $s_0 > 0$ implies $s_i > 0$ for all $i \in G$, and so also $s_j > 0$ for all $j$ which are nonnegative sums of elements in $G$. By primitivity, $\gcd(G) = 1$ and this in turn means that every sufficiently large integer can be expressed as a nonnegative sum of elements of $G$. (Refer to Exercise 5.5.) □

We should note that a periodic equation whose index of imprimitivity is $g > 1$ (such as $s_n = 4s_{n-2}$, the example at the beginning of this section with $g = 2$) can be expressed as a system of $g$ primitive equations because

$$s_n = c_g s_{n-g} + c_{2g} s_{n-2g} + \ldots + c_{rg} s_{n-rg}$$

can be written as the primitive system

$$\begin{aligned} t_n^{(0)} &= c_g t_{n-1}^{(0)} + c_{2g} t_{n-2}^{(0)} + \ldots + c_{rg} t_{n-r}^{(0)} \\ t_n^{(1)} &= c_g t_{n-1}^{(1)} + c_{2g} t_{n-2}^{(1)} + \ldots + c_{rg} t_{n-r}^{(1)} \\ &\vdots \\ t_n^{(g-1)} &= c_g t_{n-1}^{(g-1)} + c_{2g} t_{n-2}^{(g-1)} + \ldots + c_{rg} t_{n-r}^{(g-1)} \end{aligned}$$

where the initial conditions for each $< t_n^{(j)} >$ are the initial conditions of $< s_n >$ whose subscripts equal $j \pmod g$. If all initial conditions for a particular $j$ are zero, the sequence $< t_n^{(j)} >$ is the zero sequence. Otherwise, at least one initial condition for $< t_n^{(j)} >$ is positive and by the corollary $t_n^{(j)} = \Theta(\lambda_0^{ng})$. Translating this back to the original sequence $< s_n >$, we see that for a fixed $j$ the subsequence whose subscripts satisfy the arithmetic progression $n \equiv j \pmod g$ can either be the zero sequence or it will grow like $\lambda_0^n$. In some sense we can therefore regard $< s_n >$ as a periodic sequence with period $g$, or under some special circumstances the period of $< s_n >$ may be a divisor of $g$.

### *5.2.1 Roundability or When are integer solutions rounded powers of an eigenvalue?*

We've shown that the Fibonacci sequence $< f_n >$ satisfies

$$f_n = \mathit{Round}(\lambda_0^n/\sqrt{5}) \text{ for all } n \geq 0\,, \tag{5.5}$$

where $\mathit{Round}(X)$ returns the integer nearest to $X$. Is this a special property of the Fibonacci sequence or are there more general sequences with this feature? The proof of (5.5) used some properties of the Fibonacci sequence. For instance, the absolute value of the nondominant eigenvalue is relatively small when compared to $\lambda_0$ and the initial deviations are small. When we speak of **deviation** here we mean the difference between the actual element of the sequence and its approximation. (Occasionally in the following, we may use the term deviation when we mean the absolute value of the deviation.) For example,

$$d_n = f_n - \lambda_0^n/\sqrt{5}$$

are the Fibonacci deviations. More generally, for an integer recurrence sequence $< s_n >$ we want an approximation of the form $\alpha\lambda_0^n$ where $\lambda_0$ is the dominant eigenvalue and $\alpha$ is a constant which depends on the initial values. We'd like the absolute values of the deviations $d_n = s_n - \alpha\lambda_0^n$ to start small and stay small, although they need not be decreasing.

Let's first consider the special case when the eigenvalues, $\lambda_0, \ldots, \lambda_{k-1}$, are all simple. From previous work we know that $s_n$ can be written as

$$s_n = \sum_{i=0}^{k-1} \alpha_i \lambda_i^n$$

for some constants $\alpha_0, \ldots, \alpha_{k-1} \in \mathbb{C}$. For the approximation of $s_n$ by $\alpha\lambda_0^n$ the deviations satisfy

$$d_n = s_n - \alpha_0\lambda_0^n = \sum_{i=1}^{k-1} \alpha_i \lambda_i^n,$$

and the familiar triangle inequality gives

$$|d_n| \leq \sum_{i=1}^{k-1} |\alpha_i||\lambda_i|^n. \tag{5.6}$$

Inspecting this, we see for instance that if each nondominant eigenvalue were to satisfy $|\lambda_i| \leq 1$ then $s_n = \mathit{Round}(\alpha_0\lambda_0^n)$ would hold whenever $\sum_{i=1}^{k-1} |\alpha_i| < 1/2$. This result isn't as good as it might seem at first because in order to apply it we would have to calculate all $\alpha_1, \ldots, \alpha_{k-1}$!

In an effort to obtain a more effective result, let's perform a more thorough analysis of (5.6) from a graphical point of view. The upper bound in (5.6) is a decreasing function of $n$ and the deviations can be plotted as dots which can dance around (perhaps erratically) beneath the envelope formed by the bounding function. If the deviations behave very irregularly, any upper bound on them may be a dramatic overestimation. For instance, one possible type of irregularity is a **spiking behavior**, that is, the deviations might be close to zero for a while, say for $d_0, d_1, \ldots, d_5$, but then $d_6$ is relatively large in absolute value. Such spiking could occur if $\lambda_1 = (1-\epsilon)\omega$ where $\epsilon$ is a small positive number and $\omega$ is a $6^{th}$ root of unity. Longer period spiking is also possible. For instance, for $\omega_i$ an $i^{\text{th}}$ root of unity if $\lambda_1 = (1-\epsilon_1)\omega_3$ and $\lambda_2 = (1-\epsilon_2)\omega_5$, then spiking behavior with period 15 can occur if the period 3 spike adds to the period 5 spike to give a large spike of period 15. It's possible to construct sequences with even longer periods because a number of short periods could multiply together to give a long period. To summarize, the dots can be very irregular, and possibly very few dots come near the curve. In other words, one pays for the smoothness of the bound since the estimating curve is not very close to the deviations.

For general recurrences the enveloping curve given by the upper bound in (5.6) may well be the best *easy* estimate on the devations. We'll see below that putting an additional restriction on the characteristic polynomial gives a stronger result that is often quick and easy to check. For this, first

notice that since $\lambda_0$ is a *simple* eigenvalue then any solution to a general homogeneous nonnegative recurrence has the form

$$s_n = \alpha\lambda_0^n + a_1(n)\lambda_1^n + \ldots + a_{k-1}(n)\lambda_{k-1}^n \text{ for all } n \geq 0\,,$$

for some constant $\alpha$ and polynomials $a_1(x), \ldots, a_{k-1}(x)$.

**Theorem 5.2.3.** *Let $s_n = \alpha\lambda_0^n + a_1(n)\lambda_1^n + \ldots + a_{k-1}(n)\lambda_{k-1}^n$ be an integer solution to (HNN) with dominant eigenvalue $\lambda_0$. Set $d_n = s_n - a\lambda_0^n$ for all $n \geq 0$ and let $M = \max\{|d_0|, \ldots, |d_{k-1}|\}$, the maximum of the initial deviations. We further assume $f(x) = (x-1)ch(x)/(x-\lambda_0)$ is a nonnegative polynomial.*

*(a) If $M < \frac{1}{2}$ then $s_n = Round(\alpha\lambda_0^n)$ for all $n \geq 0$.*

*(b) If $f(x)$ is primitive there exists $n_0$ with $\max\{|d_{n_0}|, \ldots, |d_{n_0+k-1}|\} < 1/2$, which gives $s_n = Round(\alpha\lambda_0^n)$ for all sufficiently large $n$.*

*Proof.* There exist nonnegative $b_1, \ldots, b_k$ such that $f(x) = x^k - b_1x^{k-1} - \cdots - b_k$ where $1 - b_1 - b_2 - \cdots - b_k = 0$ since $\lambda = 1$ is a root, the dominant root of $f(x)$. We rewrite this as $\sum b_i = \sum |b_i| = 1$ and apply Exercise 5.9 to obtain that any solution $< y_n >$ of the homogeneous recurrence with characteristic polynomial $f(x)$ must satisfy

$$|y_n| \leq \max\{|y_0|, |y_1|, \cdots, |y_{k-1}|\} \quad \text{for all } n\,. \tag{5.7}$$

The sequence $< d_n >$ has the form

$$d_n = s_n - \alpha\lambda_0^n = a_1(n)\lambda_1^n + \ldots + a_{k-1}(n)\lambda_{k-1}^n\,,$$

and therefore is a solution to both (HNN) and the difference equation with characteristic polynomial $f(x)$. Therefore (5.7) holds for $y_n = d_n$, that is, all deviations satisfy $|d_n| \leq M < 1/2$, and $s_n = Round(\alpha\lambda_0^n)$.

In part (b), the fact that $f$ is nonnegative and primitive means that its only positive root $\lambda = 1$ is strictly dominant. Ordering the roots by

$$\lambda_0 = 1 > |\lambda_1| \geq |\lambda_2| \geq \cdots \geq |\lambda_{k-1}|,$$

we have

$$|d_n| \leq \sum_{i=1}^{k-1} |a_i(n)||\lambda_i|^n \leq |\lambda_1|^n \sum_{i=1}^{k-1} |a_i(n)|. \tag{5.8}$$

Since each $a_i(n)$ is a polynomial whose degree is less than the multiplicity of $\lambda_i$, then $\deg(a_i) < k$ and each $a_i(n) = O(n^k)$. Therefore, we can find universal positive constants $N, C$ such that each $|a_i(n)| < Cn^k$ for all $n \geq N$ and (5.8) gives

$$|d_n| \leq |\lambda_1|^n Ckn^k \text{ for all } n \geq N \quad .$$

Since $|\lambda_1|^n$ is exponentially decreasing to 0 while $n^k$ is growing at a polynomial rate, the inequality $|d_n| < 1/2$ eventually holds, and part (a) yields $s_n = Round(\alpha\lambda_0^n)$ for sufficiently large $n$. □

The only problem with this result is that it seems relatively difficult to show that the polynomial $f(x)$ is nonnegative. In particular, do we have to explicitly find $f(x)$? This would seem to involve the calculation of $\lambda_0$. Fortunately, there is a relatively easy condition that assures $f$ is nonnegative.

For this, define $\psi(n)$ to be the sequence whose first $k$ terms are $c_1, \ldots, c_k$ and the remaining terms are zero and let $< H_n >$ be the sequence defined by the first order recurrence

$$H_0 = 1 \quad ; \quad H_n = \lambda_0 H_{n-1} - \psi(n) \text{ for all } n \geq 1 \,.$$

Since $\lambda_0$ is a root of $ch(x) = x^k - c_1 x^{k-1} - \ldots - c_k$, then $H_k = \lambda_0 H_{k-1} - c_k = 0$ and from Theorem 3.1.2 we have

$$H_n = \begin{cases} \lambda_0^n - \sum_{i=1}^{n} c_i \lambda_0^{n-i} & \text{for } n < k \\ 0 & \text{for } n \geq k \end{cases} . \tag{5.9}$$

Also, it can be checked that

$$(x - \lambda_0)(x^{k-1} + H_1 x^{k-2} + \cdots + H_{k-1}) = ch(x)$$

which gives

$$\begin{aligned} f(x) &= (x-1)\frac{ch(x)}{x - \lambda_0} = (x-1)(x^{k-1} + H_1 x^{k-2} + \cdots + H_{k-1}) \\ &= x^k - (1 - H_1)x^{k-1} - (H_1 - H_2)x^{k-2} - \cdots - (H_{k-2} - H_{k-1})x - H_{k-1}. \end{aligned}$$

The nonnegativity of $f(x)$ is therefore equivalent to

$$1 \geq H_1 \geq H_2 \geq \cdots \geq H_{k-1} > 0 \,, \tag{5.10}$$

where the condition $H_{k-1} > 0$ is free because

$$0 = H_k = \lambda_0 H_{k-1} - c_k \quad \text{and} \quad H_{k-1} = c_k / \lambda_0 > 0.$$

It might seem that testing $H_1 = \lambda_0 - c_1 \leq 1$ would require knowledge of $\lambda_0$. But remember (recall Exercise 5.2) that for positive $x$, $ch(x) \geq 0$ if and only if $x \geq \lambda_0$, which gives the check

$$c_1 + 1 \geq \lambda_0 \iff ch(c_1 + 1) \geq 0.$$

Testing this condition is practical and can be done in at most $k$ multiplications. (Refer to Horner's Method in Section 5.5.) When we require the strict inequality $ch(c_1 + 1) > 0$ then the coefficient of $x^{k-1}$ in $f(x)$ must be nonnegative and the primitivity of $f(x)$ is ensured. As far as the other conditions in (5.10), from (5.9) we know that for all $n < k - 1$

$$H_{n+1} = \lambda_0^{n+1} - c_1 \lambda_0^n - \sum_{i=2}^{n+1} c_i \lambda_0^{n+1-i} = \lambda_0^n (\lambda_0 - c_1) - \sum_{i=1}^{n} c_{i+1} \lambda_0^{n-i} \,,$$

which gives

$$H_n - H_{n+1} = \lambda_0^n(c_1 + 1 - \lambda_0) + \sum_{i=1}^{n} (c_{i+1} - c_i)\lambda_0^{n-i}\,.$$

Therefore, $H_n < H_{n+1}$ is implied by the conditions

$$c_1 + 1 > \lambda_0 \quad \text{and} \quad c_{n+1} \geq c_{n+1} \geq \cdots \geq c_2 \geq c_1\,,$$

which are easy to check restrictions on the coefficients of the original characteristic polynomial $ch(x)$. These observations are collected in the following theorem.

**Theorem 5.2.4 (The Rounding Theorem).** *Let $\lambda_0$ be the dominant eigenvalue of the homogeneous nonnegative recurrence (HNN) whose coefficients satisfy*

$$c_{k-1} \geq \cdots \geq c_1 \quad \textit{and} \quad ch(c_1 + 1) \geq 0. \tag{5.11}$$

*Suppose $< s_n >$ is a integer solution of the form $s_n = \alpha\lambda_0^n + d_n$ for some constant $\alpha$ where $d_n$ has no $\lambda_0^n$ component.*

*(a) If $\max\{|d_0|, \ldots, |d_{k-1}|\} < 1/2$ then $s_n = Round(\alpha\lambda_0^n)$ for all $n \geq 0$.*

*(b) If the strict inequality $ch(c_1 + 1) > 0$ holds then $s_n = Round(\alpha\lambda_0^n)$ for all $n \geq n_0$ where $n_0$ is such that $\max\{|d_{n_0}|, \ldots, |d_{n_0+k-1}|\} < 1/2$.*

### *5.2.2 Using the Rounding Theorem*

For any integer $k \geq 3$, the **generalized Fibonacci sequence** is

$$f_0 = 0, f_1 = 1, f_2 = 2, \cdots, f_{k-1} = 2^{k-2}\,,\ f_n = f_{n-1} + f_{n-2} + \cdots + f_{n-k}\,,$$

where the coefficients are all 1 and the first condition of the Rounding Theorem is satisfied. Also, $c_1 + 1 = 2$ with

$$ch(2) = 2^k - 2^{k-1} - \cdots - 2 - 1 = 2^k - (2^k - 1) = 1 > 0.$$

Hence the Rounding Theorem assures us that there exist some $\alpha$ and $n_0 \geq 0$ so that $f_n = Round(\alpha\lambda_0^n)$ for $n \geq n_0$. Notice that we haven't yet used the initial conditions, and we know that the values of $\alpha$ and $n_0$ depend on the initial conditions. From $f_0 = 0, f_1 = 1, f_2 = 1, f_3 = 2, \cdots, f_{k-1} = 2^{k-3}$ we can work backwards in the sequence and obtain an equivalent set of initial conditions, $f_{-(k-2)} = f_{-(k-3)} = \cdots = f_{-1} = f_0 = 0$ and $f_1 = 1$. This translation doesn't change the value of either $\alpha$ or $n_0$ and allows an easier computation of $\alpha$ and the deviations. From Exercise 4.23,

$$\alpha = \frac{\lambda_0 - 1}{\lambda_0[(k+1)\lambda_0 - 2k]}$$

with corresponding deviations

$$\begin{aligned} d_{-(k-2)} &= 0 - \alpha\lambda_0^{-(k-2)} \\ d_{-(k-3)} &= 0 - \alpha\lambda_0^{-(k-3)} \\ &\vdots \\ d_0 &= 0 - \alpha \\ d_1 &= 1 - \alpha\lambda_0. \end{aligned}$$

Notice that $\max\{|d_{-(k-2)}|, |d_{-(k-3)}|, \cdots, |d_0|, |d_1|\} = \max\{|d_0|, |d_1|\}$ because $ch(1) = -(k-1)$ implies $\lambda_0 > 1$. Also, $d_1 > 0$ because otherwise $d_n$ would always be negative and so would have a $\lambda_0^n$ component. If we can show that both $\alpha < 1/2$ and $1 - \alpha\lambda_0 < 1/2$, then we can take $n_0 = -(k-2)$, and the generalized Fibonacci numbers can be calculated by $f_n = Round(\alpha\lambda_0^n)$ for all $n \geq -(k-2)$.

To show that $1 - \alpha\lambda_0 < 1/2$, it suffices to have $\dfrac{2(\lambda_0 - 1)}{(k+1)\lambda_0 - 2k} > 1$. From Exercise 5.4, the denominator is positive and the requirement can be written as $0 > (k-1)(\lambda_0 - 2)$ which is true because $2 > \lambda_0$.

To show $1/2 > \alpha$, we need $\frac{1}{2} > \dfrac{\lambda_0 - 1}{\lambda_0[(k+1)\lambda_0 - 2k]}$ which can be rewritten as $2 > (k+1)\lambda_0(2 - \lambda_0)$ (since the denominator is positive). Using $2 - \lambda_0 = \lambda_0^{-k}$, this is equivalent to the inequality $2\lambda_0^{k-1} > k + 1$. For $k = 2$, this reduces to $\lambda_0 > 3/2$ which is easy to verify. For $k > 2$, we use $\lambda_0^{k-1} = \lambda_0^{k-2} + \cdots + 1 + \frac{1}{\lambda_0}$ to get $2\lambda_0^{k-1} = 2(\lambda_0^{k-2} + \cdots + \frac{1}{\lambda_0}) > 2(k-1)$ using the fact that $\lambda_0 > 1$. Finally, $2(k-1) \geq k + 1$ if $k \geq 3$, and $1/2 > \alpha$ is established.

As another example, let's consider

$$s_n = 2s_{n-1} + 2s_{n-2} + 3s_{n-3}. \tag{5.12}$$

The characteristic polynomial is $ch(x) = x^3 - 2x^2 - 2x - 3$, a nonnegative polynomial with dominant root $\lambda_0 = 3$. Here $k = 3$, $c_{k-1} = c_2 = 2 \geq c_1$, and $c_1 + 1 = 2 + 1 = 3 = \lambda_0$. Because $c_1 + 1$ equals the dominant root, eventual roundability is not ensured but depends on the deviations which in turn are specified by the initial conditions. In Exercise 5.10 you show that $\alpha = \frac{1}{13}(s_0 + s_1 + s_2)$.

For initial conditions $s_0 = 1, s_1 = 3, s_2 = 9$, then $\alpha = 1$ and the deviations are

$$\begin{aligned} d_0 &= s_0 - \alpha = \quad 1 - 1 = 0 \\ d_1 &= s_1 - \alpha\lambda_0 = 3 - 3 = 0 \\ d_2 &= s_2 - \alpha\lambda_0^2 = 9 - 9 = 0. \end{aligned}$$

Therefore, the Rounding Theorem implies

$$s_n = Round(\alpha\lambda_0^n) = Round(3^n) = 3^n.$$

Of course this result could have been found directly without using the Rounding Theorem.

Using recurrence (5.12) with the initial conditions $s_0 = 0, s_1 = 0, s_2 = 1$, the coefficient is $\alpha = 1/13$ with deviations

$$\begin{aligned} d_0 &= 0 - 1/13 = -1/13 \\ d_1 &= 0 - 3/13 = -3/13 \\ d_2 &= 1 - 9/13 = 4/13. \end{aligned}$$

Since the absolute values of the initial deviations are bounded by $1/2$, then $s_n = Round(\frac{1}{13}\lambda_0^n)$ for all $n \geq 0$. For this choice roundability is not obvious without the theorem.

Let us consider one more example of initial conditions for (5.12), namely, $s_0 = 0, s_1 = 3, s_2 = 9$. Here, $\alpha = 12/13$ and the deviations are:

$$\begin{aligned} d_0 &= 0 - \frac{12}{13} = -\frac{12}{13} \\ d_1 &= 3 - \frac{12}{13} * 3 = \frac{3}{13} \\ d_2 &= 9 - \frac{12}{13} * 3^2 = \frac{9}{13}. \end{aligned}$$

In this case, the initial deviations are not all less than $1/2$ in absolute value and neither immediate rounding nor eventual rounding is promised by the Rounding Theorem since $c_1 + 1 = \lambda_0$. It is easy to calculate that

$$\begin{aligned} d_3 &= 24 - \frac{12}{13} * 3^3 = -\frac{12}{13} \\ d_4 &= 75 - \frac{12}{13} * 3^4 = \frac{3}{13} \\ d_5 &= 225 - \frac{12}{13} * 3^5 = \frac{9}{13}. \end{aligned}$$

In Exercise 5.11 you verify this pattern continues, that is, the sequence of deviations is periodic with period 3. Because the deviations do not decrease below $1/2$, even eventual rounding doesn't occur.

## 5.3 Nonhomogeneous Nonnegative Equations

In this section we look at *nonhomogeneous* nonnegative equations which have one of three types of non-zero inputs.

**Theorem 5.3.1.** *Consider a nonnegative difference equation*

$$s_n = c_1 s_{n-1} + c_2 s_{n-2} + \cdots + c_k s_{n-k} + g(n) \tag{5.13}$$

*where $g$ is a function such that the value of $g(n)$ is nonnegative for each natural number $n$. Let $\lambda_0$ be the dominant eigenvalue of the recurrence. If $s_n > 0$ for all sufficiently large $n$, then*

- $s_n = \Theta(\lambda_0^n)$ *when* $g(n) = O(\lambda_1^n)$ *for some positive* $\lambda_1 < \lambda_0$*;*
- $s_n = \Theta(g(n))$ *when* $g(n) = \Theta(\lambda_2^n)$ *for some* $\lambda_2 > \lambda_0$*;*
- $s_n = \Theta(n^{d+1}\lambda_0^n)$ *when* $g(n) = f(n)\lambda_0^n$ *for a polynomial* $f$ *with* $\deg(f) = d$.

These three types of nonnegative forcing functions respectively correspond to the situations when $g(n)$ is much less than $\lambda_0^n$, much greater than $\lambda_0^n$, and equal to $\lambda_0^n$ times a polynomial in $n$. For the last two types of forcing functions the hypothesis requiring $s_n$ to be eventually positive is superfluous. In the first type, the required eventual positivity follows once there are $k$ consecutive positive terms, and that can result from positive initial conditions or positive $g(n)$ or some combination of these forms of positiveness.

Notice that the solutions asymptotically behave about as one would expect, except that the behavior in the third case may be a little unexpected. In that case, the response to forcing by a polynomial times $\lambda_0^n$ results in a polynomial of one higher degree times $\lambda_0^n$. The most common occurrence of this is with $g(n) = \lambda_0^n$, for which $s_n = \Theta(n\lambda_0^n)$ holds. The remainder of this section is occupied with proving the theorem.

*Proof of the Theorem.* Let $N \geq 0$ be such that $s_n \neq 0$ for all $n \geq N$. Similarly to what was done for the homogeneous case (refer to page 97), for any positive $\lambda$ we define

$$S_\lambda = \{s_i/\lambda^i \,:\, N \leq i < N + k\},$$

and set $\alpha_\lambda = \min(S_\lambda)$, $\beta_\lambda = \max(S_\lambda)$. Notice that $\alpha_\lambda$ and $\beta_\lambda$ are functions of $\lambda$. In particular,

$$\alpha_\lambda \lambda^i \leq s_i \leq \beta_\lambda \lambda^i \text{ for all } N \leq i < N + k. \tag{5.14}$$

**Case A**. $g(n) = O(\lambda_1^n)$ *for some positive* $\lambda_1 < \lambda_0$. Setting $\alpha_0 = \alpha_{\lambda_0}$ and $\beta_0 = \beta_{\lambda_0}$,

$$\alpha_0 \lambda_0^n \leq s_n \leq \beta_0 \lambda_0^n \text{ for all } N \leq n < N + k.$$

As in the proof of the homogeneous case, if $s_n \geq \alpha_0 \lambda_0^n$ for all $N \leq n < K$ then

$$s_K = \sum_{i=1}^{k} c_i s_{K-i} + g(n) \geq \alpha_0 \sum_{i=1}^{k} c_i \lambda_0^{K-i} = \alpha_0 \lambda_0^K,$$

since $g(K) \geq 0$ and $ch(\lambda_0) = 0$. Therefore, by induction we have that $s_n \geq \alpha\lambda_0^n$ for all $n \geq N$.

Since $g(n) = O(\lambda_1^n)$, there exist positive $C, N_1$ such that $0 \leq g(n) < C\lambda_1^n$ for all $n \geq N_1$, and we can increase $C$ if necessary (when $N < N_1$) to obtain

$$0 \leq g(n) < C\lambda_1^n \text{ for all } n \geq N.$$

From the fact that $\lambda_1 < \lambda_0$, Exercise 5.2 implies that $-C\lambda_1^k/ch(\lambda_1)$ is a positive constant, which we call $\gamma_1$. Setting $\beta_1 = \beta_0 + \gamma_1$, we'll prove

$$s_n \leq \beta_1\lambda_0^n - \gamma_1\lambda_1^n \text{ for all } n \geq N$$

by induction on $n$ and thereby obtain $s_n = \Theta(\lambda_0^n)$. For this, we note

$$\beta_1\lambda_0^n - \gamma_1\lambda_1^n = \beta_0\lambda_0^n + \gamma_1(\lambda_0^n - \lambda_1^n) > \beta_0\lambda_0^n \geq s_n \text{ for all } N \leq n < N + k.$$

Suppose we've shown $s_n \leq \beta_1\lambda_0^n - \gamma_1\lambda_1^n$ for all $N \leq n < K$. Then

$$\begin{aligned}
s_K &= \sum_{i=1}^{k} c_i s_{K-i} + g(K) \\
&\leq \sum_{i=1}^{k} c_i(\beta_1\lambda_0^{K-i} - \gamma_1\lambda_1^{K-i}) + C\lambda_1^K \\
&= \beta_1\lambda_0^{K-k}\sum_{i=1}^{k} c_i\lambda_0^{k-i} - \gamma_1\lambda_1^{K-k}\sum_{i=1}^{k} c_i\lambda_1^{k-i} + C\lambda_1^K \\
&= \beta_1\lambda_0^K + \gamma_1\lambda_1^{K-k}(ch(\lambda_1) - \lambda_1^k) + C\lambda_1^K \\
&= \beta_1\lambda_0^K - \gamma_1\lambda_1^K + \lambda_1^{K-k}(\gamma_1\, ch(\lambda_1) + C\,\lambda_1^k) \\
&= \beta_1\lambda_0^K - \gamma_1\lambda_1^K\,,
\end{aligned}$$

by construction of $\gamma_1$, and we've shown that $s_n = \Theta(\lambda_0^n)$ does hold.

**Case B**. $g(n) = \Theta(\lambda_2^n)$ *for* $\lambda_2 > \lambda_0$. Let $\alpha, \beta > 0$ be such that

$$\alpha\lambda_2^n < g(n) < \beta\lambda_2^n\,, \text{ for all } n \geq N_1.$$

For $n \geq N_1$,

$$s_n = \sum_{i=1}^{k} c_i s_{n-i} + g(n) \geq g(n) \geq \alpha\lambda_2^n,$$

giving a lower bound on $s_n$. For an upper bound on $s_n$, from (5.14) with $\lambda = \lambda_2$ we obtain $\beta_2$ such that

$$s_n \leq \beta_2\lambda_2^n \text{ for all } N \leq n < N + k. \tag{5.15}$$

Increase $\beta_2$ if necessary to obtain $\beta_2 \geq \lambda_2^k \beta / ch(\lambda_2)$, which is positive because $\lambda_2 > \lambda_0$ (recall Exercise 5.2). Remembering (5.15), we assume $s_n \leq \beta_2 \lambda_2^n$ for all $N \leq n < M$ and prove this also holds for $n = M$, since

$$\begin{aligned} s_M &= \sum_{i=1}^{k} c_i s_{M-i} + g(M) \\ &\leq \sum_{i=1}^{k} c_i \beta_2 \lambda_2^{M-i} + \beta \lambda_2^M \\ &= \beta_2 \lambda_2^{M-k} \sum_{i=1}^{k} c_i \lambda_2^{k-i} + \beta \lambda_2^M \\ &= \beta_2 \lambda_2^{M-k} (\lambda_2^k - ch(\lambda_2)) + \beta \lambda_2^M \\ &= \beta_2 \lambda_2^M + \lambda_2^{M-k} (\beta \lambda_2^k - \beta_2 ch(\lambda_2)) \leq \beta_2 \lambda_2^M, \end{aligned}$$

from the choice of $\beta_2$. Therefore, for all $n \geq N$ we have

$$\alpha \lambda_2^n \leq s_n \leq \beta \lambda_2^n,$$

and we've proved $s_n = \Theta(\lambda_2^n)$.

**Case C**. $g(n) = \lambda_0^n f(n)$. For this case, we divide the recurrence in (5.13) by $\lambda_0^n$ to get

$$\frac{s_n}{\lambda_0^n} = \sum_{i=1}^{k} \frac{c_i}{\lambda_0^i} \frac{s_{n-i}}{\lambda_0^{n-i}} + f(n).$$

Defining $t_n = s_n / \lambda_0^n$ and $b_i = c_i / \lambda_0^i$, this becomes the new recurrence

$$t_n = \sum_{i=1}^{k} b_i t_{n-i} + f(n)$$

where we note that

$$\sum_{i=1}^{k} b_i = \sum_{i=1}^{k} \frac{c_i}{\lambda_0^i} = \frac{1}{\lambda_0^k} \sum_{i=1}^{k} c_i \lambda_0^{k-i} = 1,$$

since $\lambda_0^k = \sum_{i=1}^{k} c_i \lambda_0^{k-i}$. Because $s_n = t_n \lambda_0^n$, it suffices to prove $t_n = \Theta(n^{d+1})$.

Since $f(n) = \Theta(n^d)$, there exist constants $\gamma_1, \gamma_2$ and positive integer $N$ such that $\gamma_1 n^d \leq f(n) \leq \gamma_2 n^d$ for all $n \geq N$. Define $S = \{t_n / n^{d+1} : N \leq n < N+k\}$ and let $\alpha > 0$ be the minimal element of $S \cup \{\gamma_1/(k+1)^{d+1}\}$ and $\beta$ the maximal element of $S \cup \{\gamma_2\}$, and we'll prove $\alpha n^{d+1} \leq t_n \leq \beta n^{d+1}$ by induction on $n$. From the construction of $\alpha$ and $\beta$, this holds for all

$N \le n < N + k$. Assuming $\alpha n^{d+1} \le t_n \le \beta n^{d+1}$ for all $n < K$, we prove this inequality also holds for $n = K$. Then $t_K$ is bounded as follows

$$\sum_{i=1}^{k} b_i \alpha (K-i)^{d+1} + \gamma_1 K^d \le t_K = \sum_{i=1}^{k} b_i t_{K-i} + f(K)$$
$$\le \sum_{i=1}^{k} b_i \beta (K-i)^{d+1} + \gamma_2 K^d ,$$

where $K - k \le K - i \le K - 1$ and $\sum_{i=1}^{k} b_i = 1$ give

$$\alpha (K-k)^{d+1} + \gamma_1 K^d \le t_K \le \beta (K-1)^{d+1} + \gamma_2 K^d .$$

Therefore, the choice of $\gamma_2 \le \beta$ implies

$$t_K \le \beta K^d (K - 1 + 1) = \beta K^{d+1}$$

and $\alpha \le \gamma_1/(k+1)^{d+1}$ gives

$$t_K \ge \alpha((K-k)^{d+1} + (k+1)^{d+1} K^d)) \ge \alpha K^{d+1}$$

by Exercise 5.14. □

The following special case arises often in practice.

**Corollary 5.3.2.** *For any positive constant $b$, any nonnegative difference equation of the form*

$$s_n = c_1 s_{n-1} + c_2 s_{n-2} + \cdots + c_k s_{n-k} + b\lambda_0^n$$

*has solution $s_n = \Theta(n\lambda_0^n)$ where $\lambda_0$ is the dominant eigenvalue.*

There are many variations of the results in Theorem 5.3.1. Because these variations are proved in essentially the same way as the results in the theorem, we simply state them here and leave their proofs as exercises.

- If $g(n) = O(\lambda_1^n)$ for $\lambda_1 < \lambda_0$ then $s_n = O(\lambda_0^n)$.
  - (a) If the difference equation is primitive and $g(n) = O(\lambda_1^n)$ for $\lambda_1 < \lambda_0$ then $s_n = \Theta(\lambda_0^n)$.
  - (b) If $s_n > 0$ for $k$ consecutive values of $n$ and $g(n) = O(\lambda_1^n)$ for $\lambda_1 < \lambda_0$ then $s_n = \Theta(\lambda_0^n)$.
- Let $G(n) = \max\{g(n), \lambda_2 g(n-1), \ldots, \lambda_2^{n-1} g(1)\}$. If $g(n) = \Omega(\lambda_2^n)$ for $\lambda_2 > \lambda_0$ then
  - (a) $s_n = \Omega(\lambda_2^n)$.
  - (b) $s_n = \Omega(g(n))$.
  - (c) $s_n = O(G(n))$ .
  - (d) If $G(n) = O(g(n))$ then $s_n = \Theta(g(n))$.

(e) If $g(n) = \Omega(G(n))$ then $s_n = \Theta(G(n))$.

- If $g(n) = h(n)\lambda_0^n$ for some nonnegative $h(n)$ then $s_n = O(ng(n))$ and $s_n = \Omega(g(n))$.
- If $g(n) = h(n)\lambda_0^n$ for some nonnegative $h(n)$ which is nondecreasing then $s_n = \Theta(\sum_{j=1}^{n} \lambda_0^{n-j} g(j))$.

## 5.4 Estimating Roots

### *5.4.1 Estimation of the dominant root*

In order to find the long-term behavior of the solutions to a nonnegative difference equation we would like to estimate its dominant root. (Recall Theorem 5.3.1.) For nonnegative difference equations $s_n = \Theta(\lambda_0^n)$ often holds, and when the nonnegative equation is *primitive*, the dominant eigenvalue $\lambda_0$ is simple and $\lim_{n\to\infty}(s_n/\lambda_0^n) = \alpha$ holds for some constant $\alpha$. For the nonnegative polynomial $p(x) = x^k - c_1x^{k-1} - \cdots - c_{k-1}x - c_k$, Exercise 5.2 can be used to roughly locate its dominant root $\lambda_0$.

**Lemma 5.4.1.** *The root $\lambda_0$ depends on $c_1 + c_2 + \cdots + c_k$ in the following way:*

(1) *if $c_1 + \cdots + c_k = 1$, then $\lambda_0 = 1$;*

(2) *if $c_1 + \cdots + c_k > 1$, then $1 < \lambda_0 < c_i + \cdots + c_k$;*

(3) *if $c_1 + \cdots + c_k < 1$, then $c_1 + \cdots + c_k < \lambda_0 < 1$.*

*Proof.* We observe that $p(1) = 1 - S$, where $S = c_1 + \ldots + c_k$. Since $\lambda_0$ is the only positive root of $p(x)$, then $\lambda_0 = 1$ if and only if $S = 1$, which proves (1) and allows us to assume $\lambda_0 \neq 1$. From Exercise 5.2,

$$\lambda_0 > 1 \iff 0 > p(1) = 1 - S \iff S > 1$$

which is the required relative ordering of $\lambda_0$ and 1. Further,

$$\begin{aligned} p(S) &= S^k - c_1S^{k-1} - \ldots - c_{k-1}S - c_k \\ &= (S - c_1)S^{k-1} - c_2S^{k-2} - \ldots - c_{k-1}S - c_k \\ &= (c_2 + \ldots + c_k)S^{k-1} - c_2S^{k-2} - \ldots - c_{k-1}S - c_k \\ &= c_2S^{k-2}(S-1) + c_3S^{k-3}(S^2-1) + \ldots + c_{k-1}S(S^{k-2}-1) + c_k(S^{k-1}-1)\,, \end{aligned}$$

and the facts that each $c_i$ is nonnegative and $c_k > 0$ implies

$$S > 1 \iff p(S) > 0 \iff S > \lambda_0\,,$$

as required. □

This lemma gives useful bounds on the dominant root, positioning it between 1 and $c_1 + \ldots + c_k$, a useful first approximation for Newton's method (refer to Section 5.5.) Newton's method can also be used to approximate the other roots of a nonnegative polynomial but these roots have some extra complications. For instance, applying Newton's Method can be difficult because the second root may be nonreal and also because we do not know a good initial estimate for the second root. While Newton's Method can be used for complex numbers, there are other methods which are more effective [1].

### *5.4.2 Estimation of the second root*

While estimates of the dominant root of the characteristic polynomial are used to find the asymptotic behavior of a nonnegative recurrence, bounds on the second eigenvalue tell how quickly and in what sense the solution approaches the long-term behavior. When the nonnegative equation is primitive, $\lambda_1$ is the maximum of the nondominant eigenvalues and there exists a polynomial $\beta(x)$ such that

$$|s_n - \alpha\lambda_0^n| < \beta(n)|\lambda_1|^n .$$

From this we see that when $|\lambda_1| < 1$ there exist positive constants $\beta, M$ with $M < 1$ such that

$$|s_n - \alpha\lambda_0^n| < \beta M^n , \tag{5.16}$$

and $< s_n >$ converges to $\alpha\lambda_0^n$ in the sense of **absolute error**. On the other hand, when $|\lambda_1| \geq 1$ there exist $\beta$ and $M$, $1 < M < \lambda_0$, such that

$$\left|\frac{s_n}{\lambda_0^n} - \alpha\right| < \beta\Big(\frac{M}{\lambda_0}\Big)^n , \tag{5.17}$$

and $< s_n >$ is said to converges to $\alpha\lambda_0^n$ in the sense of **relative error**.

Can we find such a bound $M$ which is less than 1? In Appendix D we describe an algorithm for counting the number of roots of a polynomial within the (complex) unit circle $|z| < 1$. If the algorithm finds the $k^{th}$ degree characteristic polynomial has at least $k-1$ roots insie the unit circle, then $\lambda_1 < 1$ and the above analysis gives absolute error convergence of $< s_n >$ to $\alpha\lambda_0^n$. On the other hand if it's found that there are fewer than $k-1$ roots within the circle, only relative error convergence is obtained.

The method in Appendix D does not actually compute an $M$ with $\lambda_0 > M > |\lambda_1|$. One way to try to find such a bound is to take the nonnegative polynomial $p(x)$ and its positive root $\lambda_0$ and find the least $M$ for which $(x - M)p(x)/(x - \lambda_0)$ is a nonnegative polynomial. Unfortunately, this method may only yield $M = \lambda_0$. Next we give some upper bounds on the nondominant roots of a nonnegative polynomial.

**Theorem 5.4.2.** *Consider a polynomial* $p(x) = x^k - c_1x^{k-1} - \ldots - c_k$ *where all* $c_i$ *are strictly positive (and so* $p(x)$ *is primitive.) Then*

$$M = \max\left\{\frac{c_2}{c_1}, \frac{c_3}{c_2}, \ldots, \frac{c_k}{c_{k-1}}\right\}$$

*is an upper bound on* $|\lambda_1|$.

*Proof.* Since the conclusion follows if $M > \lambda_0$, we may assume $M \leq \lambda_0$. Multiplying $p(x) = x^k - c_1x^{k-1} - \cdots - c_{k-1}x - c_k$ by $x - M$ gives

$$x^{k+1} - (c_1 + M)x^k + (Mc_1 - c_2)x^{k-1} + \cdots + (Mc_{k-1} - c_k)x + c_kM\,,$$

and we call this polynomial $q(x)$. Since any root of $p(x)$ is also a root of $q(x)$, then $q(\lambda_1) = 0$ which means

$$\lambda_1^{k+1} + (Mc_1 - c_2)\lambda_1^{k-1} + \cdots + c_kM = (c_1 + M)\lambda_1^k\,.$$

Taking absolute values and recalling the definition of $M$ gives

$$|\lambda_1|^{k+1} + (Mc_1 - c_2)|\lambda_1|^{k-1} + \cdots + c_kM \geq (c_1 + M)|\lambda_1|^k\,,$$

and $q(|\lambda_1|) = (|\lambda_1| - M)p(|\lambda_1|) \geq 0$. Because $p(x)$ is a nonnegative polynomial, its values change from negative to positive at $x = \lambda_0$ and primitivity implies that $\lambda_0$ is strictly dominant. This means that $p(|\lambda_1|) < 0$, and $q(|\lambda_1|) \geq 0$ gives $|\lambda_1| \leq M$. □

For example, if $c_1 > c_2 > c_3 > \cdots > c_k > 0$, then $M < 1$, $|\lambda_i| < 1$ and (5.16) says any nonnegative solution $< s_n >$ has limiting behavior in the *absolute* error sense, that is, $\lim_{n\to\infty} |s_n - \alpha\lambda_0^n| = 0$.

If any of $c_1, \cdots, c_{k-1}$ is zero then Theorem 5.4.2 cannot be used to get an upper bound on $\lambda_1$. Another result is

**Theorem 5.4.3.** *Let* $p(x) = x^k - c_1x^{k-1} - \cdots - c_k$ *be a nonnegative polynomial with* $\lambda_0$ *its positive real root. If* $H_i = \lambda_0^i - c_1\lambda_0^{i-1} - \cdots - c_i$ *for all* $i = 1, \ldots, k$ *(as seen earlier in (5.9)) then the maximum of the absolute values of the roots of* $p(x)$ *excluding* $\lambda_0$ *satisfies*

$$|\lambda_1| \leq \max\left\{H_1, \frac{H_2}{H_1}, \frac{H_3}{H_2}, \ldots, \frac{H_{k-1}}{H_{k-2}}\right\}. \tag{5.18}$$

*Proof.* In Exercise 5.20 you show that each $H_i$ is positive, and we may assume that the maximum in (5.18) is strictly less than $\lambda_0$ since otherwise the conclusion holds. The roots of the polynomial $p(x)/(x - \lambda_0)$ are $\lambda_1, \ldots, \lambda_{k-1}$ and

$$\frac{p(x)}{x - \lambda_0} = x^{k-1} + H_1x^{k-2} + \cdots + H_{k-1}\,.$$

Also, for any positive $w$ the polynomial $g(x) = (x-w)\dfrac{p(x)}{x-\lambda_0}$ has only one positive real root and

$$g(x) = x^k - (w - H_1)x^{k-1} - (H_1 w - H_2)x^{k-2} \\ - \ldots - (H_{k-2}w - H_{k-1})x - w\,H_{k-1}\,.$$

Restricting to

$$w > \max\left\{H_1, \frac{H_2}{H_1}, \ldots, \frac{H_{k-1}}{H_{k-2}}\right\},$$

the polynomial $g(x)$ is primitive and nonnegative with the strictly dominant root $w$. Since $w$ can be taken arbitrarily close to the maximal element of (5.18), $|\lambda_1|$ must be less than or equal to this maximum. □

*Example* 5.4.1. Consider the **generalized Fibonacci polynomial** $x^k - x^{k-1} - \cdots - 1$ whose dominant root satisfies $\lambda_0 < 2$. (Refer to Exercise 5.4.) To apply the last theorem we need to bound the various $H$'s. It is true that $1 > H_1 > H_2 > \cdots > H_{k-1}$ because $1 > H_1 = \lambda_0 - 1$ (which is true since $2 > \lambda_0$), and $H_i > H_{i+1}$ is equivalent to $\lambda_0^i > \lambda_0^{i+1} - \lambda_0^i$ (which again is true because $2 > \lambda_0$.) Finally, $H_1 = \lambda_0 - 1 > \lambda_0 - \frac{1}{H_i} = \frac{H_{i+1}}{H_i}$ because $1 > H_i$. So for the generalized Fibonacci polynomial, the maximum in (5.18) occurs at $H_1$ and the result gives $|\lambda_1| \le \lambda_0 - 1$. Since $\lambda_0 < 2$, this gives a slightly better bound than $|\lambda_1| < 1$.

Note that in general, $\dfrac{H_{i+1}}{H_i} = \lambda_0 - \dfrac{c_{i+1}}{H_i}$ holds, and this shows

**Corollary 5.4.4.**

$$|\lambda_1| \le \lambda_0 - \min\left\{c_1, \frac{c_2}{H_1}, \ldots, \frac{c_{k-1}}{H_{k-2}}\right\}.$$

This corollary can also be written in another form. For this, set

$$g_i = \sum_{j=i}^{k} c_j \lambda_0^{k-j} = c_i\lambda_0^{k-i} + \cdots + c_{k-1}\lambda_0 + c_k$$

where $0 = p(\lambda_0) = \lambda_0^{k-i}H_i - g_{i+1}$ and so

$$H_i = \frac{g_{i+1}}{\lambda_0^{k-i}} \text{ and } \frac{H_{i+1}}{H_i} = \frac{g_{i+2}}{\lambda_0^{k-i-1}} \cdot \frac{\lambda_0^{k-1}}{g_{i+1}} = \frac{g_{i+2}}{g_{i+1}}\lambda_0\,.$$

This gives

**Corollary 5.4.5.**

$$|\lambda_1| \le \lambda_0 \max\left\{\frac{g_2}{\lambda_0^k}, \frac{g_3}{g_2}, \frac{g_4}{g_3}, \ldots, \frac{g_k}{g_{k-1}}\right\}.$$

## 5.5 Calculation of the Roots

As we said earlier, we would like to have an efficient algorithm for computing the dominant root to desired accuracy because the dominant root is used to find the long-term behavior of solutions. Beginning with an interval $[L, U]$ containing $\lambda_0$, another interval approximation of $\lambda_0$ can be obtained by computing $p(M)$ for $M = (U + L)/2$. (Refer to Figure 5.1.) If $p(M) = 0$ then $M = \lambda_0$ and we've located the root. If $p(M) > 0$, we replace $U$ by $M$ and otherwise replace $L$ by $M$. This method, usually called the **Bisection Method**, is guaranteed (up to round-off error) to produce a sequence of decreasing intervals which always contain $\lambda_0$. Since the interval is halved at each step, repeating the bisection method for $n$ steps will produce an interval of size $(U - L)/2^n$ which is known to contain $\lambda_0$. Bisection requires that $U$ and $L$ isolate one root of the polynomial. For a nonnegative polynomial, by Lemma 5.4.1 the single positive real root is between 1 and $c_1 + \ldots + c_r$, so starting values of $U$ and $L$ are immediately available.




FIGURE 5.1. A nonnegative polynomial with the U and L to be used in calculating the root $\lambda$.

**Bisection Method**

The following code finds an interval of width $E$ which contains the root of the polynomial $p(x)$ when started with an interval $[L, U]$ where $p(L)$ and $p(U)$ have different sign.

```
A := p(U)
B := p(L)
WHILE U − L > E DO
        MID := (U + L)/2
        C := p(MID)
        IF A ∗ C > 0, THEN A := C
                           U := MID
                      ELSE B := C
                           L := MID
```

The decision rule used in the Bisection Method requires polynomial evaluation. If direct substitution is used, every evaluation of an arbitrary polynomial of $p(x) = c_0x^d + c_1x^{d-1} + \ldots + c_d$ of degree $d$ at some $x = a$ would require $\Theta(d^2)$ multiplications. The following code, which is usually called **Horner's Method**, uses only $d$ multiplications.

**Horner's Method**

A polynomial of the form $p(x) = c_0x^d + c_1x^{d-1} + \cdots + c_d$ can be evaluated at $x = a$ by the following code which uses $d$ multiplications and $d$ additions.

```
H := c_0
FOR i = 1 TO d DO
        H := H ∗ a + c_i
```

Note that when the polynomial $p(x)$ is monic (namely, $c_0 = 1$) the terms generated by Horner's Method in the evaluation of $p(x)$ at $\lambda_0$ are exactly the $H_i$ defined earlier in (5.9), and

$$p(x) = (x - \lambda_0)(x^{d-1} + H_1x^{d-2} + H_2x^{d-3} + \ldots + H_{d-1}),$$

which is why Horner's method is sometimes called **synthetic division**. In 1964 Pan [107] showed that Horner's Method uses the fewest number of operations to evaluate a polynomial at an arbitrary point when no precomputing using the coefficients is allowed. This does not preclude evaluating

a polynomial at $m$ points using fewer than $md$ operations. (Also refer to Exercise 38 in Section 4.6.4 in Knuth [77].)

Because a polynomial of degree $d$ can be evaluated using Horner's Method in $d$ multiplications and $d$ additions, the Bisection Method will compute the root to accuracy $(U-L)/2^n$ in $\Theta(nd)$ steps. Since multiplications often take much more time than additions, the computing time is often given as only the number of multiplications. Therefore, if $E$ is the designated error bound for the root and $E_0$ the initial error bound, the root can be found within error $E$ using $d\log(E_0/E)$ multiplications.

While the Bisection Method works reasonably well, there is another method, **Newton's Method** (sometimes called the **Newton-Raphson method**), which usually locates a root more quickly. Again consider the graph of a nonnegative polynomial, but this time include the tangent to the curve $y = p(x)$ at $x = U$ and extrapolate the tangent until it intersects the $x$-axis at some $x$-value which we name $U_1$. (Refer to Figure 5.2.) This can always be done for nonnegative polynomials because in Exercise 5.2 (c) it is shown that for nonnegative polynomials $p(x)$ the tangent line always has positive slope when $x \geq \lambda_0$. This also guarantees that $U_1 < U$. From the graph we see that $U_1$ is greater than the positive root of $p(x)$, which means that it lies between $\lambda_0$ and $U$. This construction can be repeated using $U_1$ in place of $U$, and continued. By construction,

$$U_{n+1} = U_n - \frac{p(U_n)}{p'(U_n)} \tag{5.19}$$

because the value of the derivative $p'(U_n)$ equals the slope of the tangent line, $\dfrac{p(U_n)-0}{U_n - U_{n+1}}$. The formula in (5.19) is a nonlinear recurrence relation and defines the iterative procedure used in Newton's Method. Although there is little chance that we will be able to write down a closed form solution to this recurrence, we would like the recurrence to produce a sequence which converges to $\lambda_0$ as well as an estimation of how well the root is approximated.

The following is the standard Newton theorem for a general polynomial (for a proof refer to [18, 136]).

**Theorem 5.5.1 (Newton's Method).** *For any polynomial and any root $R$ there is a small interval around $R$ so that if Newton's method is started at any point within this interval, the approximations found by Newton's method will stay in the interval and converge to $R$. Further, if the error $E_i$ is the distance from $R$ at the $i^{th}$ step of Newton's method, then $E_{i+1} = O(E_i^2)$. This is called* **quadratic convergence**.

There are several practical problems with this theorem. One is that it only states the existence of the interval and it does not tell you how to find it. Also, it only promises rapid *asymptotic* convergence. The graph in Figure 5.2 suggests that the situation might be better for locating the




FIGURE 5.2. A nonnegative polynomial with two tangent lines indicated. The tangents cross the axis at the successive Newton approximations to the root of the polynomial.

positive root of a nonnegative polynomial, and in particular we hope that Newton's Method might yield a sequence which is rapidly decreasing to $\lambda_0$.

Recall that we plan to initialize Newton's Method with either $U_0 = 1$ or $U_0 = c_1 + \ldots + c_k$, according to whether $p(1)$ is positive or negative. We showed above that the constructed sequence $< U_n >$ will be decreasing provided each $U_n$ is greater than $\lambda_0$, that is, provided each error $E_n = U_n - \lambda_0$ is positive. The sequence of errors has the form

$$E_{n+1} = U_{n+1} - \lambda_0 = E_n - \frac{p(U_n)}{p'(U_n)} = \frac{E_n p'(U_n) - p(U_n)}{p'(U_n)}\,, \tag{5.20}$$

and expanding each of $p(x)$ and $p'(x)$ into its Taylor series about $\lambda_0$ (refer to Appendix B) gives

$$p(x) = \sum_{i=1}^{d} \frac{p^{(i)}(\lambda_0)}{i!}(x-\lambda_0)^i\,; \quad p'(x) = \sum_{i=0}^{d-1} \frac{p^{(i+1)}(\lambda_0)}{i!}(x-\lambda_0)^i\,,$$

which means

$$\begin{aligned} E_n p'(U_n) - p(U_n) &= E_n \sum_{i=0}^{d-1} \frac{p^{(i+1)}(\lambda_0)}{i!}(E_n)^i - \sum_{i=1}^{d} \frac{p^{(i)}(\lambda_0)}{i!}(E_n)^i \\ &= \sum_{i=1}^{d} \Big(\frac{1}{(i-1)!} - \frac{1}{i!}\Big) p^{(i)}(\lambda_0) E_n^i = \sum_{i=2}^{d} \frac{i-1}{i!} p^{(i)}(\lambda_0) E_n^i\,. \end{aligned}$$

Therefore,

$$E_{n+1} = \sum_{i=2}^{d} \frac{i-1}{i!} \frac{p^{(i)}(\lambda_0)}{p'(U_n)} E_n^i.$$

where $U_n = \lambda_0 + E_n$ and for sufficiently small $E_n$, $p'(U_n) \approx p'(\lambda_0)$ and $E_{n+1} \approx \dfrac{p^{(2)}(\lambda_0)}{2p'(\lambda_0)} E_n^2$. Since both $p^{(i)}(\lambda_0)$ and $p'(U_n)$ are positive, $E_n$ and $E_{n+1}$ must have the same sign. Therefore Newton's method does yield a sequence which is monotonically decreasing to $\lambda_0$ provided the initial value is greater than $\lambda_0$.

If Newton's method is applied to functions which are not nonnegative polynomials, then various types of odd behavior are possible (refer to [123], [141]). But even for nonnegative polynomials, Newton's method is only guaranteed to converge to the positive real root when the initial value is larger than $\lambda_0$. For example, consider the nonnegative polynomial $f(x) = x^3 - 5x$, which has the three distinct real roots, $-\sqrt{5}$, 0, and $\lambda_0 = \sqrt{5}$. Newton's formula for this polynomial can be written as

$$N(x) = x - \frac{x(x^2-5)}{3x^2-5)} = \frac{2x^3}{3x^2-5}\,,$$

where $\sqrt{5} < N(x) < x$ holds for $x > \sqrt{5}$, and when the iteration is started at some $x > \sqrt{5}$, the iteration will monotonically decrease to $\sqrt{5}$. But if the iteration is started at $x = 1$ then

$$N(1) = -1 \quad \text{and} \quad N(-1) = 1\,,$$

which means the Newton iteration oscillates with period 2. To see what is happening here, such initial values of period 2 satisfy $x = N^{(2)}(x)$, the two-fold iteration of $N$. Using the Newton iteration, such values must be zeros of the polynomial

$$x(x^2-1)(x^2-5)(4x^4-15x^2+25)\,,$$

where $x = 0$ and $x = \pm\sqrt{5}$ are fixed points (have period 1). As we saw above, the points $x = \pm 1$ are points of period 2, and this polynomial has four nonreal roots which are also points of period 2.

It is straightforward but extremely tedious to derive a $27^{\text{th}}$-degree polynomial from $x = N^{(3)}(x)$, the third iterate of $N$. If we did, after eliminating the three points of period 1 there would be 24 points of period 3, some of which might not be real. Similarly, for any $n$ one can derive a polynomial from $x = N^{(n)}(x)$, a polynomial whose roots have period dividing $n$. Once the points which have a period which is a proper divisor of $n$ are eliminated, all remaining points will oscillate with period $n$ under the Newton iteration .

The point is that even something as seemingly simple as Newton's method applied to a nonnegative polynomial can have complicated behavior. One

should exercise care in picking a good starting point to assure rapid convergence. In practice, these complications are unlikely to show up because round-off error will usually be sufficient to keep the iteration from settling down into periodic behavior. A more practical difficulty is starting at or near a zero of the derivative. This can throw the iteration into some unexpected region and may lead to numerical underflow or overflow. For nonnegative polynomials, starting at $x = U_0$ ensures the derivatives will have non-zero values.

In summary, Newton's method will be well-behaved when used to find the positive real root of a nonnegative polynomial, but you must remember to use a starting value which is larger than the root.

### *5.5.1 The rate of convergence in Newton's method*

To obtain additional insight into the behavior of Newton's method for nonnegative polynomials $p(x)$ we look at the sequence of errors. For this it's helpful to write $E(x) = x - \lambda_0$ and $\hat{E} = E(N(x))$ for $x > \lambda_0$. Suppressing all occurrences of the variable $x$, the error equation (5.20) becomes

$$\hat{E} = E^2 \alpha \,,$$

where $\alpha = (Ep' - p)/p'E^2$ is a function of $x$. Since $E = x - \lambda_0$ is a factor of $p(x)$, there exists a polynomial $q$ such that $p = Eq$. Using the product rule to calculate the derivative $p' = Eq' + q$, then (5.20) becomes

$$\alpha = \frac{Ep' - p}{p'E^2} = \frac{q'}{p'} \tag{5.21}$$

and allows us to prove the following result.

**Theorem 5.5.2.** *For a nonnegative polynomial $p(x)$ with positive root $\lambda_0$, the error in Newton's method started at any $x > \lambda_0$ obeys*

$$\hat{E} < E^2 \frac{p''(\lambda_0)}{2\,p'(\lambda_0)} \,.$$

*Proof.* Since $E(\lambda_0) = 0$ gives $p''(\lambda_0) = 2q'(\lambda_0)$, the result follows from (5.21) if we can show that $\alpha(x)$ is decreasing for $x \geq \lambda_0$. For this we prove that $\alpha'(x)$ is nonpositive for all $x > \lambda_0$,

$$p'q'' - p''q' = T_1 q'' - p'' T_2 \leq 0 \,, \tag{5.22}$$

for

$$T_1 = p' - \frac{x}{k-1} p'' \,; \quad T_2 = q' - \frac{x}{k-1} q'' .$$

For any polynomial $f(x) = \sum_{j=0}^{d} a_j x^{d-j}$ it is true that

$$\begin{aligned} f'(x) - \frac{x}{d-1} f''(x) & \\ &= \sum_{j=0}^{d-1} (d-j) a_j x^{d-1-j} - \frac{x}{d-1} \sum_{j=0}^{d-1} (d-j)(d-1-j) a_j x^{d-2-j} \\ &= \sum_{j=1}^{d-1} (d-j) a_j x^{d-1-j} \left(1 - \frac{d-1-j}{d-1}\right) \\ &= \sum_{j=1}^{d-1} (d-j) a_j x^{d-1-j} \frac{j}{d-1} \,. \end{aligned}$$

This identity implies

$$T_1 = \sum_{j=1}^{d-1} \frac{d-j}{d-1} (-c_j) x^{d-1-j} \,,$$

which is negative for all positive $x$. Also, using the sequence obtained from Horner's method, $H_0 = 1, H_1, \ldots, H_{s-1}$ are negative with $q(x) = \sum_{i=0}^{s-1} H_i x^{s-1-i}$ and the above identity gives $T_2 \geq 0$. Recalling that $p'(x)$ is a nonnegative polynomial whose dominant root is less than $\lambda_0$, Exercise 5.2(c) implies $p''(x) > 0$ for all $x > \lambda_0$. Therefore, (5.22) is satisfied for all $x > \lambda_0$, and the result is proved. □

This result is helpful because it says that the error will satisfy $E_i < \alpha(\lambda_0)^{-(2^i+1)}$ after $i$ steps if initially $E_0 < 1/\alpha(\lambda_0)^2$. That is, in $i$ steps the error will satisfy $E_i < \alpha(\lambda_0)^{-(2^i+1)}$. The difficulty is we don't know how long it will take for Newton's Method to reduce the error to less than $1/\alpha(\lambda_0)^2$.

Until now we've considered only *absolute* error, that is, the difference (or deviation) of our estimate from the actual value. Often comparing the size of the actual error with the size of $\lambda_0$ is more appropriate, and that's what is meant by **relative error**, that is, $\delta(x) = E(x)/\lambda_0$. Rephrasing the above result in terms of relative error:

$$\hat{\delta}(x) = \frac{\hat{E}(x)}{\lambda_0} = \frac{(\delta\lambda_0)^2 \alpha(x)}{\lambda_0} < \delta^2 \frac{\lambda_0 p''(\lambda_0)}{2\, p'(\lambda_0)}.$$

The homogeneity of the convergence factor associated with the relative error criterion allows the following

**Theorem 5.5.3.** *The relative error in Newton's method for approximating the dominant root of a $k^{th}$ degree nonnegative polynomial obeys*

$$\hat{\delta} < (k-1)\delta^2 \,.$$

*In fact, the convergence factor* $\lambda_0 p'(\lambda_0)/2p''(\lambda_0)$ *is guaranteed to lie in the interval* $[(k-1)/2, k-1)$.

*Proof.* First we note that for any polynomial $f(x) = \sum_{i=0}^{k-1} a_i x^{k-1-i}$ of degree $k-1$ we have

$$(k-1)f(x) - xf'(x) = \sum_{i=1}^{k-1} a_i i x^{k-1-i}\,. \tag{5.23}$$

Using this identity for $f(x) = p'(x) = kx^{k-1} - c_1(k-1)x^{k-2} - \ldots - c_{k-1}$, we see that no coefficient in the polynomial $(k-1)p'(x) - xp''(x)$ is positive, and $x = \lambda_0$ gives the lower bound of $(k-1)/2$ for the convergence factor.

From $p'(\lambda_0) = q(\lambda_0)$ and $p''(\lambda_0) = 2q'(\lambda_0)$ we get another representation of the convergence factor which is not computationally helpful unless the quotient $q(x)$ can be determined. However, this representation can be used to successfully obtain the upper bound we want to prove here, namely that

$$\lambda_0 q'(\lambda_0) < (k-1)q(\lambda_0)\,.$$

Recalling that

$$q(\lambda_0) = \lambda_0^{k-1} + H_1\lambda_0^{k-2} + \cdots + H_{k-1}$$

where each Horner element $H_i$ is positive, (5.23) with $f(x) = q(x)$ gives the required upper bound. □

This result gives doubling convergence for *quadratic* polynomials. It can be extended to polynomials of any degree.

**Corollary 5.5.4.** *Let* $\delta_0$ *be the initial relative error when Newton's method is used to approximate the dominant root of a* $k^{th}$ *degree nonnegative polynomial. If* $\delta_0$ *satisfies* $\delta_0 < 1/k$ *then the number of correct digits double at each iteration when the computation is done in base* $b = k/(k-1)$.

*Proof.* From the theorem, $\delta_{i+1} < (k-1)\delta_i$ and

$$\delta_1 < (k-1)/k^2, \quad \delta_2 < \frac{(k-1)^3}{k^4}, \ldots, \delta_r < \frac{(k-1)^{2^r-1}}{k^{2^r}}\,.$$

□

It remains to locate an initial value for which the relative error is sufficiently small. For this we consider the notion of "scaling". In this context, scaling means replacing $x$ by a new variable $y$ of the form $x = \beta y$. The standard example is approximating a square root. (Refer to Section 9.4.4 for an analysis of the running time when Newton's method is used to calculate a square root.) The square root of $A > 0$ can be computed by using Newton's

method to find the dominant root $\lambda_0$ of the polynomial $p(x) = x^2 - A$ . The Newton iteration formula is

$$N(x) = x - \frac{x^2 - A}{2x},$$

or the computationally better

$$N(x) = \frac{x}{2} + \frac{A}{2x}\,.$$

Next we scale the expression $x^2 - A$ by replacing $x$ by $x = 2^l y$ where $4^{l-1} \leq A < 4^l$ (such $l$ might be negative). Then $x^2 - A = 0$ iff $y^2 - \frac{A}{4^l} = 0$. Once we scale, the root $\mu_0 = \sqrt{A}/2^l$ of $h(y) = y^2 - A/4^l$ satisfies $1/2 < \mu_0 < 1$ and when $y = 1$ is chosen as the initial value, $E_0 < 1/2$. In this example, the convergence factor for the relative error is

$$\frac{\mu_0 h''(\mu_0)}{h'(\mu_0)} = 1,$$

from which we obtain doubling convergence. Hence, the number of correct bits when Newton's Method is used to find $\mu_0$ at least doubles at each iteration, that is, $E < 1/2^r$ implies $\hat{E} < 1/2^{2r}$. After the root $\mu_0$ has been determined with desired accuracy, the root $\lambda_0$ is easily obtained by multiplying by $2^l$, a binary shift. Scaling helped to identify a good initial value for the Newton iteration, and information about the scaled polynomial easily translates back to information about the original polynomial.

In general, if $p(x) = x^k - \sum_{i=1}^{k} c_i x^{k-i}$ is a nonnegative polynomial with dominant root $\lambda_0$, scaling it by $\beta$ yields the new polynomial

$$h(y) = \beta^{-k}((\beta y)^k - \sum_{i=1}^{k} c_i(\beta y)^{k-i}) = y^k - \sum_{i=1}^{k} \beta^{-i} c_i y^{k-i},$$

with dominant root $\mu_0 = \lambda_0/\beta$. It can be checked that the relative convergence factor remains unchanged, that is,

$$\lambda_0 p''(\lambda_0)/p'(\lambda_0) = \mu_0 h''(\mu_0)/h'(\mu_0)\,.$$

How is this used? First, if it happens that $\sum c_i > 1$, then we repeatedly scale by a factor of $\beta = k/(k-1)$ until the coefficient sum either equals 1 (in which case the dominant root equals 1) or is less than 1. The last scaling places the scaled root $\mu_0$ in the interval $((k-1)/k,\, 1)$. If it happens that $\mu_0$ lies in the subinterval $((k-1)/k, k/(k+1))$ of length $1/k(k+1)$, then $\delta_0 < 1/(k^2-1) < 1/k$ holds for the initial value $U_0 = k/(k+1)$. Otherwise, when $\mu_0 \in (k/(k+1), 1)$, the usual $U_0 = 1$ gives $\delta_0 < 1/(k+1)$. In either case, we've identified an initial value for which the relative error satisfies $\delta_0 < 1/k$ and doubling convergence is ensured. Once the correct

degree of accuracy is attained for $\mu_0$, we scale back to $\lambda_0$ by a shift in base $\beta = k/(k-1)$. Since our scaled values are less than 1, the relative error is an upper bound on the absolute error and the number of correct digits really does double at each iteration.

## 5.6 Exercises

**Ex 5.1.** Let $p(x)$ be a polynomial with real coefficients, and let $\lambda$ be a root of $p(x)$ whose multiplicity is $m \geq 2$. Show that $\lambda$ is a root of $p'(x)$ with multiplicity $m-1$.

**Ex 5.2.** Show that

(a) If $p(x)$ is a nonnegative polynomial with dominant real root $\lambda_0$ then the values of $p(x)$ change from negative to positive at $x = \lambda_0$.

(b) If $p(x)$ is a nonnegative polynomial with positive root $\lambda_0$, then $p'(x)$ is positive for all $x \geq \lambda_0$. Use this to conclude that $p(x) > 0$ for $x > \lambda_0$.

**Ex 5.3.** Consider the recurrence

$$s_n = s_{n-1} + 2s_{n-2}.$$

(a) Show the solution is $\Theta(2^n)$ for $s_0 = 0$, $s_1 = 15$.

(b) When $s_0 = -4$ and $s_1 = 7$, show that $s_n > 0$ for all $n \geq 3$ and that $s_n = \Theta(2^n)$ still holds.

(c) When $s_0 = -1$, $s_1 = 1$, show that $s_n \neq \Theta(2^n)$.

**Ex 5.4.** In this problem for $k \geq 3$ we consider the $k^{\text{th}}$-order Fibonacci recurrence

$$f_n = f_{n-1} + f_{n-2} + \ldots + f_{n-k}$$

with primitive nonnegative characteristic polynomial $ch(x)$ and dominant root $\lambda_0$.

(a) Use Newton's Method initialized at $x_0 = 2$ to show that $\lambda_0 \leq 2 - 1/(2^k - 1)$.

(b) Show that $f(x)$ has a minimum at $x = 2k/(k+1) = 2 - 2/(k+1)$ on the interval $x \geq 1$. Noting that $f(x) = (x-1)ch(x)$, show that $\lambda_0 \geq 2 - 2/(k+1)$.

**Ex 5.5.** (a) Let $a$ and $b$ be fixed positive integers. Show that an integer $n$ is expressible as a nonnegative combination of $a$ and $b$ (that is, there exist *nonnegative* integers $i, j$ such that $n = ia + jb$) iff $n$ is in

one of the following sequences

$$\begin{aligned}
&0, b, 2b, \dots \\
&a, b+a, 2b+a, \dots \\
&2a, b+2a, 2b+2a, \dots \\
&\dots \\
&(b-1)a, b+(b-1)a, 2b+(b-1)a, \dots
\end{aligned}$$

(b) When $\gcd(a,b) =$ show that each of the above sequences is in a different congruence class modulo $b$. Use that to show that every integer $n \geq (a-1)(b-1)$ is in one of these sequences and so can be written as a nonnegative combination of $a$ and $b$.

(c) If $G$ is any finite set of positive integers with $\gcd(G) = 1$ show that every sufficiently large integer can be written as a nonnegative sum of elements in $G$.

**Ex 5.6.** Let $G$ be a finite set of positive integers and let $g = \gcd(G)$ be the greatest common divisor of the elements of $G$. Then $\zeta$ is an $i^{\text{th}}$ root of unity for every $i \in G$ if and only if $\zeta$ is a $g^{\text{th}}$ root of unity.

**Ex 5.7.** Show that

(a) $p(x) = (x-3)(x+1)^3$ is a primitive nonnegative polynomial.

(b) $p(x) = x^4 - x^3 - 5x^2 - x - 6$ is a primitive nonnegative polynomial with two roots on the unit circle.

**Ex 5.8.** Consider

$$p(x) = x^5 - \frac{1}{2}x^3 - \frac{9}{2}x^2 - \frac{55}{16}x - \frac{25}{8}\,.$$

Show that 2 is the unique positive root of $p(x)$ and that there exists a quadratic polynomial $q(x)$ such that $p(x) = (x-2)[q(x)]^2$ and every root $\lambda$ of $q(x)$ satisfies $|\lambda| = \sqrt{5}/2$. With this you see that even a primitive nonnegative polynomial can have multiple subdominant eigenvalues.

**Ex 5.9.** Show that any solution to the recurrence

$$y_n = b_1 y_{n-1} + \cdots + b_k y_{n-k}$$

with $\sum_{i=1}^{k} |b_i| \leq 1$ satisfies $|y_n| \leq \max\{|y_0|, |y_1|, \cdots, |y_{k-1}|\}$.

**Ex 5.10.** In this problem we consider the sequence $< s_n >$ satisfying the nonnegative homogeneous recurrence

$$s_n = 2s_{n-1} + 2s_{n-2} + 3s_{n-3}.$$

Use the method of Exercise 4.22 to show that the coefficient of the dominant eigenvalue in the closed form of the solution is $\alpha = (s_0 + s_1 + s_2)/13$.

**Ex 5.11.** Show that the sequence of deviations is periodic for the solution to $s_n = 2s_{n-1} + 2s_{n-2} + 3s_{n-3}$ with $s_0 = 0, s_1 = 3, s_2 = 9$.

**Ex 5.12.** Show that the nonnegative solutions to

$$x_n = 5x_{n-1} + 4x_{n-2}$$

converge to $\alpha\lambda_0^n$ in the absolute value sense. If the initial conditions are nonnegative integers, will $x_n = Round(\alpha\lambda_0^n)$?

**Ex 5.13.** (a) Let $p(x) = x^k + c_1x^{k-1} + \ldots + c_{k-1}x + c_k$ be a polynomial with integer coefficients. Show that any real root of $p(x)$ is either an integer or is an irrational number.

(b) Give an efficient algorithm to determine if the dominant root of a nonnegative polynomial with integer coefficients is an integer.

**Ex 5.14.** Write $n^{d+1}$ as $[(n-k)+k]^{d+1}$ and use the binomial expansions of this and of $(k+1)^{d+1}$ to show

$$(n-k)^{d+1} \geq n^{d+1} - (k+1)^{d+1}n^d\,.$$

**Ex 5.15.** Find the asympotic size of the solution to the nonnegative equation

$$s_n = 2s_{n-1} + 2s_{n-2} + 3s_{n-3} + g(n)$$

for each of the following choices of $g(n)$:

$$g(n) = 2^n\,; \quad g(n) = 4^n\,; \quad g(n) = n3^n\,.$$

**Ex 5.16.** Find the asymptotic size of the solution to

$$f_n = f_{n-1} + f_{n-2} + g(n) \quad f_0 = 0, f_1 = 1\,,$$

where $g(n)$ satisfies the recurrence

$$g(n) = g(n-1) + g(n-2) \quad g(0) = 0, g(1) = 1\,.$$

**Ex 5.17.** Let $< s_n >$ be the solution of the nonnegative recurrence

$$s_n = c_1s_{n-1} + c_2s_{n-2} + \cdots + c_ks_{n-k} + g(n)\,.$$

Show that

(a) If $g(n) = O(\lambda_1^n)$ for $\lambda_1 < \lambda_0$ show that $s_n = O(\lambda_0^n)$.

(b) If the difference equation is primitive and $g(n) = O(\lambda_1^n)$ for $\lambda_1 < \lambda_0$ then $s_n = \Theta(\lambda_0^n)$.

(c) If $s_n > 0$ for $k$ consecutive values of $n$ and $g(n) = O(\lambda_1^n)$ for $\lambda_1 < \lambda_0$ then $s_n = \Theta(\lambda_0^n)$.

**Ex 5.18.** Let $< s_n >$ be the solution of the nonnegative recurrence

$$s_n = c_1 s_{n-1} + c_2 s_{n-2} + \cdots + c_k s_{n-k} + g(n) .$$

Let $G(n) = \max\{g(n), \lambda_2 g(n-1), \ldots, \lambda_2^{n-1} g(1)\}$. If $g(n) = \Omega(\lambda_2^n)$ for $\lambda_2 > \lambda_0$ show that

(a) $s_n = \Omega(\lambda_2^n)$.
(b) $s_n = \Omega(g(n))$.
(c) $s_n = O(G(n))$ .
(d) If $G(n) = O(g(n))$ then $s_n = \Theta(g(n))$.
(e) If $g(n) = \Omega(G(n))$ then $s_n = \Theta(G(n))$.

**Ex 5.19.** Let $< s_n >$ be the solution of the nonnegative recurrence

$$s_n = c_1 s_{n-1} + c_2 s_{n-2} + \cdots + c_k s_{n-k} + g(n) .$$

(a) If $g(n) = h(n)\lambda_0^n$ for some nonnegative $h(n)$ then $s_n = O(ng(n))$ and $s_n = \Omega(g(n))$.
(b) If $g(n) = h(n)\lambda_0^n$ for some nonnegative $h(n)$ which is nondecreasing then $s_n = \Theta(\sum_{j=1}^{n} \lambda_0^{n-j} g(j))$.

**Ex 5.20.** Let $p(x) = c_0 x^d + c_1 x^{d-1} + \ldots + c_d$ be a nonnegative polynomial with dominant root $\lambda_0$. Show that each $H_i$ generated by Horner's Method in the evaluation of $p(x)$ at $\lambda_0$ is positive.

**Ex 5.21.** Show that the polynomial $f(x) = x^3 + 3x^2 - 13x + 17$ has one real root and that this real root is negative. Try various positive initial conditions and see if Newton's method converges to the root. Conclude from this that the assumption of nonnegativity is needed in order to be assured that any initialization of Newton's method above the root will converge to the root.

**Ex 5.22.** Show that there exists an initial value for which Newton's Method applied to $f(x) = x^3 + 3x^2 - 13x + 17$ oscillates with period 3.

# 6
# Leslie's Population Matrix Model

## 6.1 Leslie's Model

In the Fibonacci Story (see Chapter 1) the rabbits are immortal, but with a few exceptions like the Energizer Bunny we know that real rabbits have a small finite lifetime. We can make this model more reasonable by reinterpreting the meaning of the age classes while still maintaining the mathematical form of the model. Recalling the original model, if $A_t$ is the number of adults at time $t$ and $Y_t$ is the number of young at time $t$ then

$$\begin{pmatrix} A_{t+1} \\ Y_{t+1} \end{pmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} \begin{pmatrix} A_t \\ Y_t \end{pmatrix}.$$

The assumed immortality of the adults is represented by the 1 in the upper left of the matrix. The young disappear by becoming adults, but they are replaced by each adult producing a youth. What happens if we simply interchange adults and young leaving the matrix alone? We get

$$\begin{pmatrix} Y_{t+1} \\ A_{t+1} \end{pmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} \begin{pmatrix} Y_t \\ A_t \end{pmatrix},$$

but our interpretation is now quite different. New young are produced by both young and adults. The adults disappear, but they are replaced by young who grow into adulthood. So by reinterpreting this model we can replace the unrealistic assumption of immortality with the biologically realistic assumptions of aging and death. As a leading biologist said at a

recent meeting, "the facts are always changing, but a good model goes on forever."

The reinterpreted model still has some unrealistic features. Should every youth survive to adulthood? Should the number of offspring produced by an adult and a youth be the same? The model can be generalized to avoid these unrealistic features by replacing the 1's in the matrix with other constants. Since the lower left 1 represents a youth surviving to become an adult, we could replace this 1 by $s$, which could be called the **survival rate**, and reasonably assume that $1 \geq s > 0$. The 1's in the top row represent the number of offspring produced by a youth and an adult respectively. We can replace these 1's with the **fertility rates** $f_1$ and $f_2$ and simply assume that $f_1 \geq 0$ and $f_2 \geq 0$. The generalized model is

$$\begin{pmatrix} Y_{t+1} \\ A_{t+1} \end{pmatrix} = \begin{bmatrix} f_1 & f_2 \\ s & 0 \end{bmatrix} \begin{pmatrix} Y_t \\ A_t \end{pmatrix}.$$

Several special cases suggest themselves. What happens if $f_1 = f_2 = 0$? Then, if we start with the population $\begin{pmatrix} Y_0 \\ A_0 \end{pmatrix}$, we next get $\begin{pmatrix} 0 \\ sY_t \end{pmatrix}$ and then $\begin{pmatrix} 0 \\ 0 \end{pmatrix}$. So as expected if the fertility rates are 0, the population dies out, and in this model the population dies out in two time steps. What happens if $f_2 = 0$, but $f_1 \neq 0$? Then starting with $\begin{pmatrix} Y_t \\ A_t \end{pmatrix}$, we get $\begin{pmatrix} f_1 Y_t \\ sY_t \end{pmatrix}$ and then $\begin{pmatrix} Y_t f_1^2 \\ Y_t s f_1 \end{pmatrix}$. These last two vectors can be rewritten as $Y_t \begin{pmatrix} f_1 \\ s \end{pmatrix}$ and $Y_t f_1 \begin{pmatrix} f_1 \\ s \end{pmatrix}$. Notice that these are both multiples of the vector $\begin{pmatrix} f_1 \\ s \end{pmatrix}$, and that neither depends on $A_t$. These observations agree with the expectations that the individuals past reproductive age will eventually have no effect on the composition of the population, and the eventual shape of the population will depend on the structural parameters $f$'s and $s$, and not on the initial population. What happens if $f_1 = 0$ and $f_2 \neq 0$? Then, starting with $\begin{pmatrix} Y_t \\ A_t \end{pmatrix}$, we get $\begin{pmatrix} f_2 A_t \\ sY_t \end{pmatrix}$ and then $\begin{pmatrix} s f_2 Y_t \\ s f_2 A_t \end{pmatrix}$. Unlike the other cases, the shape of the population oscillates between multiples of $\begin{pmatrix} Y_t \\ A_t \end{pmatrix}$ and $\begin{pmatrix} f_2 A_t \\ sY_t \end{pmatrix}$. In the second and third cases the population may increase or decrease depending on whether the multiplier, $f_1$ or $sf_2$, is or is not bigger than 1. These cases show that this simple model does have some features that we would like in a population model. The population size may increase or decrease. The population may die out. The shape of the population may approach a stable form or the shape of the population may oscillate. By doing some simple calculations on the parameters, we would like to be able to determine which of these various situations will occur.

We can generalize the Fibonacci model with two age classes to a model with $k$ age classes. In population biology the model with $k$ age classes is usually called **Leslie's model**. In 1945, Leslie [84] published one of the most influential papers in population biology. In it he introduced vectors and matrices to a generation of biologists. The model Leslie described is quite similar to the renewal model [53] and the method of life-tables which were devised in the 1700s by such mathematicians as the Bernoullis, Euler, and Halley (refer to Boyer's *History of Mathematics*[13]). These methods form the basis of an entire industry—insurance. Further, these methods were not unknown to earlier biologists. For example, Lotka's 1925 book, *Elements of Mathematical Biology*[88], devotes an entire chapter to them. On the other hand, Leslie presented a concise formulation of the model, and the post-war generation of biologists were a receptive audience.

The model can be concisely stated as

$$X_{t+1} = LX_t$$

where $X_t$ and $X_{t+1}$ are population vectors and $L$ is a Leslie matrix. The model assumes discrete time and the time unit must be chosen appropriately for the organism being modeled. For bacteria the time unit might be 20 minutes. For many insects an appropriate time unit might be one week. For many vertebrates a one year time unit might be used. For human populations a five year time unit is often used. The population vectors have a number of components. Each component is the number of individuals of a particular age. For example, if

$$X_t = (x_1, x_2, x_3, x_4)^T ,$$

then at time $t$, $x_1$ is the number of individuals in the first age class, $x_2$ is the number of individuals in the second age class, and so forth. Said another way, $x_i$ is the number of individuals of age $i-1$ because newborns are usually assigned age 0 and not 1. If one is not really happy with the discrete time assumption and assumes that discrete time is only an approximation to an underlying continuous time, then one could say that $x_i$ represents the individuals with ages $a$ where $i-1 \leq a < i$. One could instead include in $x_i$ the ages where $i-1 < a \leq i$ and also include newborns in $x_1$. Or one could say that ages where $i-1 \leq a \leq i$ are represented by $x_i$ and claim there is no ambiguity because there are no individuals of exactly age $i$ for any $i$. The main point is that the width of an age class is one time unit. If the data are arranged for five year age classes, this model will not calculate the population one year in the future.

Generalizing from the Fibonacci example, the **Leslie matrix** will contain both survival rates and fertility rates. Specifically:

$$L = \begin{bmatrix} f_1 & f_2 & \dots & \dots & f_k \\ s_1 & 0 & \dots & \dots & 0 \\ & s_2 & & & \vdots \\ & & \ddots & & \vdots \\ & & & s_{k-1} & 0 \end{bmatrix},$$

that is, the first row of $L$ consists of fertility rates where $f_i$ is the number of offspring (newborn) produced by an individual of age class $i$ in one time unit, and the subdiagonal of $L$ contains the survival rates where $s_i$ is the probability that an individual in age class $i$ will survive to age class $i+1$. All other entries in $L$ are 0.

The usual assumptions on these parameters are that for each $i$, $0 < s_i \leq 1$ and $f_i \geq 0$. The first assumption makes sense if one interprets $s_i$ as a probability and assumes that there is some possibility for an individual to survive a particular age class. Further if any $s_i$ were 0, then in $k-i$ steps the population would become a population in which the last $k-i$ age classes were empty, and all future developments would occur within and depend only on the first $i$ age classes. For similar reasons, one usually assumes that $f_k \geq 0$. That is, if one or several of the oldest age classes have 0 fertility, then the composition of these age classes will have no effect on the rest of the population, and in a small number of steps the composition of these age classes will be determined by the compositions of the younger age classes with no effect from the original composition of these oldest age classes.

An extra assumption made in Leslie's original model and often used in demographic applications is that at least 2 adjacent fertility rates are positive. This assumption is often enforced by averaging fertilities. That is, the number of offspring from females in each age class is measured but a fraction of these are attributed to females in the next age class because the females are assumed to be aging as the measurements are taken. A mathematically more appropriate assumption, which includes the Leslie assumption as a special case, is that there is a power of the Leslie matrix which is strictly positive. Luckily this can be checked easily using the greatest common divisor of the indices of positive fertility rates.

**Theorem 6.1.1.** *There exists $m$ so that $L^m \gg 0$ iff $\gcd\{i \mid f_i > 0\} = 1$, where $A \gg 0$ means that every entry in the matrix $A$ is strictly positive.*

This theorem says that an $m$ making $L^m$ strictly positive exists. The following theorem gives bounds on the smallest power $m$ which makes $L^m$ strictly positive.

**Theorem 6.1.2.** *Let $L$ be an $n \times n$ Leslie matrix with* $\gcd\{i \mid f_i > 0\} = 1$, *and let $m_0$ be the least nonegative integer so that $L^{m_0} \gg 0$, and let $l$ be the least positive integer so that $f_l > 0$, then*

$$m_0 \leq l(n-2) + n \leq (n-1)^2 + 1\,.$$

We will prove both of these theorems in Chapter 7 in the more general context of nonnegative matrices (which might not be in Leslie form.) The bounds given in Theorem 6.1.2 are tight. For any $n \geq 2$, consider a Leslie matrix in which $f_1 = f_2 = \cdots = f_{n-2} = 0$, $f_{n-1} > 0$ and $f_n > 0$. Here, $\gcd(n, n-1) = 1$ and it is relatively easy to show that $m_0 = (n-1)^2 + 1$. (See Exercise 6.8.) These two theorems can be generalized to apply to all nonnegative matrices. A nonnegative matrix $A$ is called **primitive** if there exists positive $m$ so that $A^m \gg 0$. A graph $G$ can be associated with $A$ in which vertices $v_i$ and $v_j$ have an edge $v_i \leftarrow v_j$ iff $a_{ij} > 0$. (The other direction, $v_i \rightarrow v_j$, could be used.) A graph is **strongly connected** iff for each pair of vertices $v_i$ and $v_j$ there is a directed path from $v_i$ to $v_j$ and a directed path from $v_j$ to $v_i$. When the directed edges are represented by arrows, a directed path always follows the arrows from tail to head. The generalized theorems are:

**Theorem 6.1.3.** *A nonnegative $n \times n$ matrix $A$ is primitive iff the associated graph $G$ is strongly connected and the greatest common divisor of the cycle lengths in $G$ is 1.*

**Theorem 6.1.4.** *If $A$ is a primitive $n \times n$ matrix, then $m_0$, the least nonnegative integer so that $A^{m_0} \gg 0$, obeys*

$$m_0 \leq l(n-2) + n \leq (n-1)^2 + 1$$

*where $l$ is the length of the shortest cycle in the associated graph $G$.*

Again we will defer the proofs of these theorems until Chapter 7 where we will also describe in more detail the graph associated with a nonnegative matrix.

### *6.1.1 How To Tell If A Leslie Matrix Is Primitive*

How does one determine if a nonnegative matrix has a power which is strictly positive? For a Leslie matrix, strong connectedness holds (since all $s_i$ are positive), so only the gcd condition needs to be checked. Luckily for Leslie matrices, every cycle in the graph contains the vertex 1 and the necessary and sufficient condition for primitivity in Theorem 6.1.3 becomes $\gcd\{i | f_i > 0\} = 1$ for Leslie matrices.

Notice that if $L^m$ is positive then $L^{m_1}$ will be positive for all $m_1 > m$. So one way to test if a Leslie matrix is primitive is to raise the matrix to the $(n-1)^2 + 1$ power and see if this power of the matrix is strictly positive.

This technique will require a number of matrix multiplications and hence may not be very efficient. A more efficient algorithm can be based on the gcd condition. Compute $\gcd\{i|f_i > 0\}$ by using the **Euclidean greatest common divisor algorithm**. This algorithm is given in the next two boxes.

**Recursive form of Euclid's algorithm for Greatest Common Divisor**

$$\gcd(a,b) = \begin{cases} b & \text{if } a = 0 \\ \gcd(b \bmod a,\ a) & \text{if } a \neq 0\,. \end{cases}$$

It can be shown (see Exercise 6.2) that Euclid's algorithm computes the gcd of two numbers which are less than $n$ by using $O(\log n)$ arithmetic operations. Since Euclid's algorithm will be used at most $O(n)$ times, the whole algorithm takes $O(n \log n)$.

**Algorithm to determine if a Leslie Matrix is primitive**

```
Let {i_1, i_2, ..., i_r} = {i|f_i > 0}
    G := i_1
    FOR J := 2 TO r DO
        G := gcd(G, i_J)
    END FOR
    IF G = 1 THEN PRIMITIVE
    ELSE NOT PRIMITIVE
```

## 6.2 Leslie's Convergence Theorem

The content of Leslie's Convergence Theorem is that asymptotically a multidimensional model becomes a one-dimensional model. Specifically, the theorem states that under reasonable circumstances the population distribution will converge toward a single distribution regardless of the initial distribution. In general this convergence will be convergence in the sense of relative error. With an additional assumption, the convergence can become convergence in the sense of absolute error.

The asymptotic growth rate of the population will be determined by a single number, $\lambda_0$, which depends on the fertility rates and survival rates, but is independent of the initial population distribution. There is no closed form formula for $\lambda_0$, but it can be calculated quickly to high numerical accuracy using Newton's method (see Section 5.4).

Will the population be asymptotically increasing or decreasing? Obviously this depends on $\lambda_0$, but a highly accurate estimate of $\lambda_0$ is not needed to answer this question. All one needs to know is whether $\lambda_0 > 1$ or $\lambda_0 < 1$, and this can be determined using only $O(n)$ arithmetic operations.

What will the asymptotic distribution, usually called the **stable age distribution**, look like? This distribution can, and will, be given as an explicit formula in $\lambda_0$, and the fertility rates and the survival rates. Further, assuming that $\lambda_0 > 1$, this stable age distribution will have "inverted pyramid" form. That is, the largest age class will be the newborns and the size of each age class will decrease as age increases.

**Theorem 6.2.1 (Leslie's Convergence Theorem).** *Let $L$ be an $n \times n$ Leslie matrix in which* $\gcd\{i \mid f_i > 0\} = 1$*, let $X$ be a nonnegative vector, then there is a unique positive eigenvalue $\lambda_0$ of $L$ so that*

$$\lim_{t\to\infty} \frac{L^t X}{\lambda_0^t} = \gamma \begin{pmatrix} \frac{c_1}{\lambda_0} \\ \vdots \\ \frac{c_i}{\lambda_0^i} \\ \vdots \\ \frac{c_n}{\lambda_0^n} \end{pmatrix}$$

*where*

$$c_i = \begin{cases} 1 & \text{if } i = 1 \\ s_1 s_2 \dots s_{i-1} & \text{if } i = 2, \dots, n \end{cases}$$

*and*

$$\gamma = \sum_{j=1}^{n} \frac{\lambda_0^{j-1} g_j(\lambda_0) x_j}{ch'_L(\lambda_0) c_j}$$

*where*

$$ch_L(\lambda) = \lambda^n - \sum_{i=1}^{n} c_i f_i \lambda^{n-i}$$

*and*

$$g_j(\lambda) = \sum_{i=j}^{n} c_i f_i \lambda^{n-i}.$$

This form of the theorem is slightly stronger than the Leslie's original theorem because he assumed that there was some $i$ so that both $f_i > 0$ and $f_{i+1} > 0$. Clearly, his assumption implies the gcd condition, but there are Leslie matrices which satisfy the gcd condition and do not have two adjacent positive fertility rates. We will not prove the theorem at this point. It will follow as a corollary of some more general theorems which we will prove in this chapter. The above form for the limiting vector was chosen to display the inverted pyramid form.

**Corollary 6.2.2.** *If $\lambda_0 > 1$ and each $s_i \leq 1$ or $\lambda_0 \geq 1$ and each $s_i < 1$, then the limiting vector has inverted pyramid form; that is, the entries of the vector decrease as one goes from top to bottom.*

The other important point is that $\gamma > 0$ if $X > 0$. This follows because $ch'_L(\lambda_0) > 0$ and for each $j$ both $c_j$ and $g_j(\lambda_0)$ are positive. (For this, refer to Exercise 5.2.) Notice that if $X = 0$, then $\gamma = 0$ and $L^t X$ is always 0. The convergence is in the relative error sense since the theorems on nonnegative polynomials (Section 5.1) only guarantee that $\lambda_0$ is greater than the absolute value of the other eigenvalues. For convergence in the absolute error sense, one would want that $\lambda_0 \geq 1$, and for all other eigenvalues $\lambda_i$, $|\lambda_i| < 1$. For example, the Fibonacci matrix satisfies these hypotheses and so not only

$$\lim_{t\to\infty} \frac{F^t X}{\lambda_0^t} = \gamma \begin{pmatrix} 1/\lambda_0 \\ 1/\lambda_0^2 \end{pmatrix} = \hat{\gamma} \begin{pmatrix} \lambda_0 \\ 1 \end{pmatrix}$$

but also

$$\lim_{t\to\infty} \left| F^t X - \hat{\gamma} \lambda_0^t \begin{pmatrix} \lambda_0 \\ 1 \end{pmatrix} \right| = 0$$

where $F$ is the Fibonacci matrix, $\hat{\gamma}$ depends on $X$, and the absolute value of a vector equals the maximum absolute value of its coordinates.

## 6.3 Nonprimitive Leslie Matrices

Not all Leslie matrices converge. Some can oscillate. The idea of using oscillating matrices for populations even predates Leslie's paper. For example, Bernadelli [9] described a hypothetical population of beetles and modeled this population with an oscillating matrix model. The periodic cicadas (the famous 17 year locusts) can also be described by the oscillating matrices which we will cover in this section.

### *6.3.1 A Simple Example*

Consider the two-dimensional Leslie matrix

$$\begin{bmatrix} 0 & 8 \\ 1/2 & 0 \end{bmatrix}.$$

There is only one positive fertility rate, $f_2$, so $g = 2$ and we expect an oscillation of period 2. The characteristic polynomial is $\lambda^2 - 4$, and the eigenvalues are $+2$ and $-2$. Letting $M$ be this matrix we have

$$M^{2m} = 4^m I \quad \text{and} \quad M^{2m+1} = 4^m M\,.$$

So in some sense, the powers of the matrix are periodic. To be more precise, we can normalize by dividing $M$ by its positive eigenvalue, $\lambda_0 = 2$, giving

$$\frac{M}{2} = \begin{bmatrix} 0 & 4 \\ 1/4 & 0 \end{bmatrix}$$

and clearly

$$\left(\frac{M}{2}\right)^{2m} = I \quad \text{and} \quad \left(\frac{M}{2}\right)^{2m+1} = \frac{M}{2}\,.$$

Thus, $\frac{M}{2}$ is periodic with period 2. In general, we say that a matrix $A$ is **periodic with period** $p$ iff $A^{p+i} = A^i$ for all $i \geq 0$, and $p$ is the smallest positive integer which makes the equation true. So while in our example it is true that $\left(\frac{M}{2}\right)^{512+i} = \left(\frac{M}{2}\right)^i$ for all $i$, we do *not* say that $\left(\frac{M}{2}\right)$ has period 512, because the periodicity equations are also satisfied by $p = 2$. When we say that the period $p = 2$, we are also implying that $A^{1+i} \neq A^i$; that is, the matrix does not have period 1.

### *6.3.2 Special Case – Only one positive fertility rate*

**Theorem 6.3.1.** *Let $M$ be an $n$-dimensional Leslie matrix with $f_n$ its only nonzero fertility rate, let $\lambda_0$ be the positive eigenvalue of $M$, then $M/\lambda_0$ is periodic with period $n$. Also $\lambda_0 = (s_1 \ldots s_{n-1} f_n)^{1/n}$.*

*Proof.* By direct calculation, the characteristic polynomial of $M$ is the nonnegative polynomial $\lambda^n - s_1 \ldots s_{n-1} f_n$. So $\lambda_0 = (s_1 \ldots s_{n-1} f_n)^{1/n}$ is the (unique) positive eigenvalue. By the Cayley-Hamilton Theorem (see Appendix C), $M^n = \lambda_0^n I$ and $n$ satisfies the periodicity equations. To see that the minimal polynomial $min_M(\lambda)$ equals $ch_M(\lambda)$, that is, that $n$ is the smallest such positive integer, consider the orbit of $e_n$ under $M$: $e_n, Me_n, M^2e_n, \ldots, M^{n-1}e_n$ where $e_1, e_2, \ldots, e_n$ are the standard coordinate vectors. For $j > 0$,

$$M^j e_n = s_1 \ldots s_{j-1} f_n e_j\,,$$

implying the period divides $n$. Now if $M/\lambda_0$ were periodic with period $p$, then for every vector $X$, $M^p X$ would be a scalar multiple of $X$. But no two $e_j$'s are scalar multiples of one another. Therefore, for any $p < n$, $M^p e_n$ is not a scalar multiple of $e_n$, so $M$ cannot have period less than $n$. □

### *6.3.3 Asympotically Periodic Leslie Matrices*

Fortunately or unfortunately, most matrices, even most Leslie matrices, are not truly periodic. Even nonprimitive Leslie matrices are usually not periodic. (Recall that a matrix is primitive if and only if a power of the matrix has only positive entries.) All powers of a nonprimitive Leslie matrix

will contain some 0's. These 0's will move through the powers of the matrix in a periodic fashion. The nonzero entries will also appear periodically, but the values of the nonzero entries will be changing, preventing true periodicity. If a Leslie matrix is $n$-dimensional with at least 2 positive fertility rates, then $g$ is less than $n$, and the matrix will have $g$ eigenvalues of largest magnitude and $n-g$ eigenvalues of smaller magnitude. These smaller magnitude eigenvalues prevent the matrix from being truly periodic, but we can expect the contribution of these smaller eigenvalues to disappear (at least in a ratio sense) as we take higher powers of the matrix.

For example, consider the Leslie matrix

$$A = \begin{bmatrix} 0 & 1 & 0 & 2 \\ 1/2 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1/2 & 0 \end{bmatrix}.$$

The characteristic polynomial is $\lambda^4 - \frac{1}{2}\lambda^2 - \frac{1}{2}$ and the eigenvalues are 1, -1, $\frac{i}{\sqrt{2}}$, $\frac{-i}{\sqrt{2}}$, and $g = 2$. For this example, we expect that taking high powers of $A$ will lead to a matrix which is close to a periodic matrix of period 2. We can use the characteristic polynomial to display this behavior. Since $A^4 = \frac{1}{2}(A^2 + I)$,

$$\begin{aligned} A^{K+2} - A^K &= A^{K-2}(A^4 - A^2) = A^{K-2}(\frac{1}{2}A^2 + \frac{1}{2}I - A^2) \\ &= -\frac{1}{2}A^{K-2}(A^2 - I) = -\frac{1}{2}(A^K - A^{K-2})\,. \end{aligned}$$

So iterating this formula gives

$$A^{K+2} - A^K = \begin{cases} (-\frac{1}{2})^{K/2}\ (A^2 - I) & \text{if } K \text{ is even} \\ (-\frac{1}{2})^{\lfloor K/2 \rfloor}\ A(A^2 - I) & \text{if } K \text{ is odd}\,. \end{cases}$$

In either case, $A^2 - I$ or $A(A^2 - I)$ are fixed matrices independent of $K$, while $(\frac{-1}{2})^{K/2}$ is decreasing exponentially toward 0, so

$$\lim_{K \to \infty} (A^{K+2} - A^K) = 0$$

where this convergence is componentwise; that is, each entry in the matrix $A^{K+2} - A^K$ goes to 0. (Other notions of convergence are possible for matrices.)

Using this example as our paradigm, we say that $A$ is an **asymptotically periodic matrix** with period $g$ iff

$$\lim_{K \to \infty} (A^{K+g} - A^K) = 0$$

and $g$ is the smallest positive integer which satisfies this equation.

To show that our example $A$ has asymptotic period 2, we must show that $A$ cannot have a smaller period. If $A$ had asymptotic period 1, then for every vector $X$

$$\lim_{K\to\infty}(A^{K+1}-A^K)X=0\,.$$

(This time, the 0 on the right-hand side is a vector not a matrix.) Consider the vector

$$X=(4,-2,2,-1)^T\,.$$

For our example matrix $A$, calculation shows that $AX$ is equal to $-X$. Hence $A^KX=(-1)^KX$, and

$$(A^{K+1}-A^K)X=(-1)^{K+1}X-(-1)^KX=2(-1)^{K+1}X\,,$$

which does not go to zero. Hence, $A$ has asymptotic period 2.

## 6.4 Companion Matrices

To set the stage for our proof of the Leslie Convergence Theorem, we will take a detour through companion matrices. Our intention is to prove convergence results about *matrices* and from these derive convergence results for *vectors*. In fact we will be able to generalize Leslie's theorem by considering both primitive and nonprimitive matrices.

Let $A$ be a matrix in **companion** form; that is,

$$A=\begin{bmatrix}c_1 & c_2 & \dots & c_n\\ 1 & 0 & \dots & 0\\ & \ddots & & \\ 0 & & 1 & 0\end{bmatrix}$$

where $c_n\neq 0$. Clearly companion matrices have features in common with Leslie matrices and the relationships between these two types of matrices will be explained a little later. Companion matrices are called companion because they are closely associated with polynomials of the form $\lambda^n-c_i\lambda^{n-1}-c_2\lambda^{n-2}-\cdots-c_n$. Given such a polynomial it is easy to write down its companion matrix. This companionship is not just one-way since the characteristic polynomial of $A$ is, up to sign, the polynomial to which $A$ is companion. This last remark can be verified easily by writing down

the determinant and expanding it via its last column. That is,

$$ch_A(\lambda) = det(A - \lambda I) = \begin{vmatrix} c_1 - \lambda & c_2 & \dots & \dots & c_n \\ 1 & -\lambda & & & \\ & 1 & \ddots & & \\ & & \ddots & \ddots & \\ & & & 1 & -\lambda \end{vmatrix}$$

$$= -\lambda \begin{vmatrix} c_1 - \lambda & c_2 & \dots & \dots & c_{n-1} \\ 1 & -\lambda & & & \\ & 1 & \ddots & & \\ & & \ddots & \ddots & \\ & & & 1 & -\lambda \end{vmatrix} - (-1)^n c_n \begin{vmatrix} 1 & -\lambda & & & \\ & 1 & -\lambda & & \\ & & \ddots & \ddots & \\ & & & \ddots & -\lambda \\ & & & & 1 \end{vmatrix} =$$

$$\vdots$$

$$= -\lambda(-1)^{n-1}(\lambda^{n-1} - c_1\lambda^{n-2} - \cdots - c_{n-1}) - (-1)^n c_n$$
$$= (-1)^n[\lambda^n - c_1\lambda^{n-1} - \cdots - c_{n-1}\lambda - c_n]\,,$$

(the ellipses indicate an inductive argument). In what follows we will blithely ignore the $(-1)^n$, and simply say that $ch_A(\lambda) = \lambda^n - c_1\lambda^{n-1} - \cdots - c_n$. In fact, we will usually drop the subscript $A$ and just say $ch(\lambda) = \lambda^n - c_1\lambda^{n-1} - \cdots - c_n$.

The Cayley-Hamilton Theorem (see Appendix C) which says that every matrix satisfies its characteristic polynomial and gives the relation $ch(A) = A^n - c_1A^{n-1} - c_2A^{n-2} - \cdots - c_nI = 0$ between a polynomial and its companion matrix. We will use this relation to compute powers of the matrix $A$. From the relation,

$$A^n = c_1A^{n-1} + c_2A^{n-2} + \cdots + c_nI\,, \tag{6.1}$$

so

$$A^{n+1} = c_1A^n + c_2A^{n-1} + \cdots + c_nA\,,$$

and using (6.1),

$$A^{n+1} = (c_1^2 + c_2)A^{n-1} + (c_1c_2 + c_3)A^{n-2} + \dots + (c_1c_{n-1} + c_n)A + c_1c_nI\,.$$

Extending this example, all powers of $A$ can be expressed as linear combinations of the zero$^{th}$ through $n-1^{st}$ power of $A$, i.e., for any $k$ there are coefficients $\alpha_1(k), \dots, \alpha_n(k)$ so that

$$A^k = \alpha_1(k)A^{n-1} + \alpha_2(k)A^{n-2} + \cdots + \alpha_n(k)I\,.$$

This formula says that the problem of computing the powers of a matrix can be reduced to computing $n$ scalar functions. Further, the formulas for computing these functions have the following simple form:

$$\begin{aligned}
\alpha_1(k+1) &= c_1 \cdot \alpha_1(k) + \alpha_2(k) \\
\alpha_2(k+1) &= c_2 \cdot \alpha_1(k) + \alpha_3(k) \\
&\vdots \\
\alpha_{n-1}(k+1) &= c_{n-1} \cdot \alpha_1(k) + \alpha_n(k) \\
\alpha_n(k+1) &= c_n \cdot \alpha_1(k)
\end{aligned}$$

This is a set of $n$ coupled difference equations. These equations can be separated by converting to an $n^{th}$-order difference equation. For example,

$$\alpha_1(k+1) = c_1\alpha_1(k) + c_2\alpha_1(k-1) + \cdots + c_n\alpha_1(k-n+1)\,.$$

If one knew the solution to this difference equation, then the other scalar functions could be computed by

$$\begin{aligned}
\alpha_2(k+1) &= c_2\alpha_1(k) + c_3\alpha_1(k-1) + \cdots + c_n\alpha_1(k-n+2) \\
&\vdots \\
\alpha_n(k+1) &= c_n\alpha_1(k)\,.
\end{aligned}$$

If one desired, these could also be replaced by $n$ uncoupled difference equations

$$\begin{aligned}
\alpha_1(k+1) &= c_1\alpha_1(k) + c_2\alpha_1(k-1) + \cdots + c_n\alpha_1(k-n+1) \\
\alpha_2(k+1) &= c_1\alpha_2(k) + c_2\alpha_2(k-1) + \cdots + c_n\alpha_2(k-n+1) \\
&\vdots \\
\alpha_n(k+1) &= c_1\alpha_n(k) + c_2\alpha_n(k-1) + \cdots + c_n\alpha_n(k-n+1)\,.
\end{aligned}$$

Notice that these are really the same difference equations but each copy of the equation will, in general, have a different solution because the initial conditions will be different for each copy.

Although the above formulas give computationally efficient methods to compute powers of a matrix, we would like a single method which gives a good estimate of the asymptotic behavior of the sequence of powers of a matrix. That is, what will $A^k$ look like when $k$ is large? In general the answer is easy, either all the elements in $A^k$ will be large—that is, grow exponentially in magnitude as a function of $k$—or all elements of $A^k$ will be small— that is, decrease exponentially in magnitude as a function of $k$. Of course, there will be $A$'s which are exceptions to this general case. In fact, what we would like to do is to change $A$ so that the overall growth or

decline trend is normalized out, and then see how $A^k$ will behave. To do so we would like to assume that there is a single number $\lambda_0$, so that $\lambda_0$ is the largest number (in magnitude) such that $A-\lambda_0 I$ is singular and then study the behavior of $(A/\lambda_0)^k$. We expect the growth trend to be normalized out, so that $(A/\lambda_0)^k$ may have a limit and we want to compute this limit. It will be important to determine or at least estimate $\lambda_0$, and we will return to this estimation later.

First, let us look for an **eigenmatrix** $P_\lambda$ whose form is preserved but whose magnitude is multiplied by $\lambda$ when $P_\lambda$ is multiplied by $A$. That is, we want to solve the equation

$$AP_\lambda = \lambda P_\lambda \,. \tag{6.2}$$

Since we are interested in computing powers of $A$, we assume that $P_\lambda$ can be represented as a polynomial in $A$. So assuming

$$P_\lambda = p_0 A^{n-1} + p_1 A^{n-2} + \cdots + p_{n-1} I \,,$$

then

$$\begin{aligned} \lambda P_\lambda = AP_\lambda &= p_0 A^n + p_1 A^{n-1} + \cdots + p_{n-1} A \\ &= (p_0 c_1 + p_1) A^{n-1} + (p_0 c_2 + p_2) A^{n-2} + \ldots \\ &\quad + (p_0 c_{n-1} + p_{n-1}) A + p_0 c_n I \,. \end{aligned} \tag{6.3}$$

Equating the coefficients of corresponding powers of $A$ on each side of (6.3) gives

$$\begin{aligned} p_0 c_n &= p_{n-1}\lambda \\ p_0 c_{n-1} + p_{n-1} &= p_{n-2}\lambda \\ &\vdots \\ p_0 c_2 + p_2 &= p_1 \lambda \\ p_0 c_1 + p_1 &= p_0 \lambda \,. \end{aligned} \tag{6.4}$$

Assuming $\lambda \neq 0$, these equations can be solved as:

$$\begin{aligned} p_{n-1} &= \frac{p_0 c_n}{\lambda} \\ p_{n-2} &= \frac{p_0 c_{n-1}}{\lambda} + \frac{p_{n-1}}{\lambda} = p_0 \left(\frac{c_{n-1}}{\lambda} + \frac{c_n}{\lambda^2}\right) \\ &\vdots \\ p_1 &= \frac{p_0 c_2}{\lambda} + \frac{p_2}{\lambda} = p_0 \left(\frac{c_2}{\lambda} + \frac{c_3}{\lambda^2} + \cdots + \frac{c_n}{\lambda^{n-1}}\right) . \end{aligned} \tag{6.5}$$

The last equation and (6.4) give

$$p_0 c_1 + p_1 = p_0 \left(c_1 + \frac{c_2}{\lambda} + \cdots + \frac{c_n}{\lambda^{n-1}}\right) = p_0 \lambda \,.$$

This is a valid equation for all values of $p_0$ if $\lambda^n = c_1\lambda^{n-1}+\cdots+c_n$, which is true for every $\lambda$ which is a root of the characteristic polynomial. Of course if $\lambda$ is not a root of the characteristic polynomial, then this last equation implies that $p_0 = 0$, and hence by (6.5) that $p_i = 0$ for all $i$.

The eigenmatrices $P_\lambda$ can also be expressed in terms of the values of the polynomials $g_i(\lambda)$, defined as

$$\begin{aligned}
g_1(\lambda) &= \lambda - c_1 \\
g_2(\lambda) &= \lambda^2 - c_1\lambda - c_2 = \lambda g_1(\lambda) - c_2 \\
&\vdots \\
g_i(\lambda) &= \lambda^i - c_1\lambda^{i-1} - \cdots - c_i = \lambda g_{i-1}(\lambda) - c_i \\
g_{n-1}(\lambda) &= \lambda^{n-1} - c_1\lambda^{n-2} - \cdots - c_{n-1} \\
g_n(\lambda) &= \lambda^n - c_1\lambda^{n-1} - \cdots - c_{n-1}\lambda - c_n = ch(\lambda)
\end{aligned}$$

which form the sequence of polynomials from Horner's method (see Section 5.4). So up to a scalar multiplier $P_\lambda$ is

$$P_\lambda = A^{n-1} + g_1(\lambda)A^{n-2} + g_2(\lambda)A^{n-2} + \cdots + g_{n-1}(\lambda)I\,, \tag{6.6}$$

which follows by using $ch(\lambda) = 0$ and rearranging the formulas from (6.5).

In the simple case when $ch_A(\lambda)$ has $n$ distinct roots $\lambda_1, \lambda_2, \ldots, \lambda_n$, we will show it is easy to represent powers of $A$ in terms of the $P_\lambda$. Assume that there are $n$ scalars $\alpha_1, \alpha_2, \ldots, \alpha_n$ so that

$$I = \sum_{i=1}^{n} \alpha_i P_{\lambda_i}\,, \tag{6.7}$$

then

$$A^k = A^k \sum_{i=1}^{n} \alpha_i P_{\lambda_i} = \sum_{i=1}^{n} \alpha_i A^k P_{\lambda_i} = \sum_{i=1}^{n} \alpha_i \lambda_i^k P_{\lambda_i}\,.$$

Further, if $\lambda_1$ is a strictly dominant eigenvalue then

$$\lim_{k\to\infty} \frac{A^k}{\lambda_1^k} = \lim_{k\to\infty} \sum_{i=1}^{n} \alpha_i \left(\frac{\lambda_i}{\lambda_1}\right)^k P_{\lambda_i} = \alpha_1 P_{\lambda_1}\,.$$

We are left with calculating $\alpha_1$. Luckily, this is not difficult. Since $AP_{\lambda_i} = \lambda_i P_{\lambda_i}$, $(A - \lambda_i I)P_{\lambda_i} = 0$ and for any scalars $a, b$,

$$\begin{aligned}
&(A - \lambda_i I)(A - \lambda_j I)(aP_{\lambda_i} + bP_{\lambda_j}) \\
&\quad = a(A - \lambda_j I)(A - \lambda_i I)P_{\lambda_i} + b(A - \lambda_i I)(A - \lambda_j I)P_{\lambda_j} \\
&\quad = a(A - \lambda_j I)0 + b(A - \lambda_i I)0 = 0\,.
\end{aligned}$$

So the matrix polynomial $(A-\lambda_2 I)(A-\lambda_3 I)\cdot\cdots\cdot(A-\lambda_n I)$ annihilates every linear combination of $P_{\lambda_2},\ldots,P_{\lambda_n}$, that is,

$$(A-\lambda_2 I)\ldots(A-\lambda_n I)\sum_{i=2}^{n} b_i P_{\lambda_i} = 0\,.$$

Now the matrix polynomial $(A-\lambda_2 I)\ldots(A-\lambda_n I)$ can be written as $\dfrac{ch(A)}{A-\lambda_1 I}$ where the bar means polynomial division since the polynomial in $A$ in the denominator exactly divides the polynomial in $A$ in the numerator. Next, consider the action of $(A-\lambda_j I)$ on $P_{\lambda_1}$. We see

$$(A-\lambda_j I)P_{\lambda_1} = AP_{\lambda_1}-\lambda_j P_{\lambda_1} = \lambda_1 P_{\lambda_1}-\lambda_j P_{\lambda_1} = (\lambda_1-\lambda_j)P_{\lambda_1}\,.$$

So

$$(A-\lambda_i I)(A-\lambda_j I)P_{\lambda_1} = (\lambda_1-\lambda_i)(\lambda_1-\lambda_j)P_{\lambda_1}$$

and

$$\begin{aligned}\frac{ch(A)}{A-\lambda_1 I}P_{\lambda_1} &= (\lambda_1-\lambda_2)(\lambda_1-\lambda_3)\ldots(\lambda_1-\lambda_n)P_{\lambda_1}\\ &= \lim_{\lambda\to\lambda_1}\frac{ch(\lambda)}{\lambda-\lambda_1}P_{\lambda_1}\\ &= \lim_{\lambda\to\lambda_1}\frac{ch(\lambda)-ch(\lambda_1)}{\lambda-\lambda_1}P_{\lambda_1}\\ &= ch'(\lambda_1)P_{\lambda_1}\,.\end{aligned}$$

Actually, in this calculation we made use of no special property of the eigenvalue $\lambda_1$, so

$$\frac{ch(A)}{A-\lambda_i I}P_{\lambda_i} = ch'(\lambda_i)P_{\lambda_i}$$

holds for each eigenvalue $\lambda_i$.

We now want to use this result on the equation

$$I = \sum_{i=1}^{n}\alpha_i P_{\lambda_i}$$

to compute the $\alpha_i$'s. So

$$\frac{ch(A)}{A-\lambda_j I}\,I = \sum_{i=1}^{n}\alpha_i\,\frac{ch(A)}{A-\lambda_j I}\,P_{\lambda_i} \tag{6.8}$$

$$= \alpha_j\,ch'(\lambda_j)P_{\lambda_j} \tag{6.9}$$

where

$$P_{\lambda_j} = A^{n-1}+g_1(\lambda_j)A^{n-2}+\cdots+g_{n-1}(\lambda_j)I\,.$$

Since the coefficient of $A^{n-1}$ in $\frac{ch(A)}{A-\lambda_j I}$ is 1, equating the coefficients we have $\alpha_j = 1/ch'(\lambda_j)$. In fact, we also have $P_{\lambda_j} = \frac{ch(A)}{A-\lambda_j I}$, which one can easily verify is the same matrix polynomial as the one we computed in (6.6). These computations demonstrate the following theorem.

**Theorem 6.4.1.** *If $A$ is a companion matrix with $n$ distinct eigenvalues and $\lambda_1$ is a strictly dominant eigenvalue then*

$$\lim_{k\to\infty} \frac{A^k}{\lambda_1^k} = \frac{1}{ch'(\lambda_1)} P_{\lambda_1} \quad \textit{where} \quad P_{\lambda_1} = \frac{ch(A)}{A-\lambda_1 I}\,.$$

A few comments on this theorem are necessary. First, if the eigenvalues are distinct isn't there only one largest eigenvalue? No, even if we assume that eigenvalues are distinct, there may still be several eigenvalues with the same magnitude. Consider

$$A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

Here, $ch_A(\lambda) = \lambda^2 - 1$, and the two eigenvalues, $\lambda_1 = 1$ and $\lambda_2 = -1$, have the same magnitude. In this case, it is easy to see that

$$A^K = \begin{cases} I & \text{if } K \text{ is even} \\ A & \text{if } K \text{ is odd} \end{cases}.$$

So, $A^K/\lambda_1^K$ has no limit. Taking absolute values will not help because

$$\left|A^K/\lambda_1^K\right| = \left|A^K/\lambda_2^K\right| = \begin{cases} I & \text{if } K \text{ is even} \\ A & \text{if } K \text{ is odd} \end{cases}.$$

It would be possible to get a convergence theorem if absolute values were replaced by *matrix norms*. Recall that absolute value takes a complex number which may be negative or have a non-zero imaginary part and returns a positive real number, the magnitude of the complex number. In a similar fashion, it is possible to define a norm which takes a matrix and returns a positive real number, the magnitude of the matrix. We will not pursue matrix norms at this point. Second, why does $\lambda_1$ have to be distinct from the other eigenvalues? Because the limit contains $1/ch'(\lambda_1)$, and if $\lambda_1$ were a multiple eigenvalue, $ch'(\lambda_1)$ would be 0, and the claimed limit would not be defined. Also, there would be an inconsistency in (6.8) and the scalars in (6.7) could not be found.

There is a logical hole in what we have just done. We assumed that solving for $P_\lambda$ could be carried out by equating the coefficients of $A^j$ for each $j$ on either side of the equation $AP_\lambda = \lambda P_\lambda$. (See equation (6.2).) This is tantamount to assuming that the powers of $A$ are *linearly independent*; that is, if

$$b_1 A^{n-1} + b_2 A^{n-2} + \cdots + b_n I = 0\,,$$

then $b_1 = b_2 = \cdots = b_n = 0$. This is equivalent to the minimal polynomial being equal to the characteristic polynomial. For general matrices this assumption is not valid. For example, if $A = I$, then $1 \cdot A + (-1) \cdot I = 0$ and the powers of this $A$ are not linearly independent. Fortunately, we have an extra assumption about the matrices we are using. We have assumed that they are in *companion form*, that is,

$$A = \begin{bmatrix} c_1 & c_2 & \dots & c_n \\ 1 & 0 & \dots & 0 \\ & \ddots & & \\ 0 & & 1 & 0 \end{bmatrix}$$

with $c_n \neq 0$.

We now show that if $A$ is in companion form then $A^{n-1}$, $A^{n-2}, \dots, I$ are linearly independent. Assume that we have $b_1, b_2, \dots, b_n$ so that

$$b_1 A^{n-1} + b_2 A^{n-2} + \cdots + b_n I = 0\,. \tag{6.10}$$

We want to show that $A$ being in companion form forces each $b_i$ to be 0. This equation (6.10) is a matrix equation and for it to hold as a matrix equation it must hold for each column. In particular, $A^i\, e_n$ will be the last column of $A^i$ and then (6.10) implies that

$$b_1 A^{n-1}\, e_n + b_2 A^{n-2}\, e_n + \cdots + b_n I\, e_n = 0\,. \tag{6.11}$$

We claim that $A^i\, e_n$ has the form

$$\begin{pmatrix} * \\ * \\ \vdots \\ * \\ c_n \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix} \quad \left.\begin{matrix} \\ \\ \\ \\ \\ \\ \\ \\ \end{matrix}\right\} \; i \qquad \left.\begin{matrix} \\ \\ \\ \\ \\ \\ \\ \\ \end{matrix}\right\} \; n-i\,;$$

that is, the $i^{\text{th}}$ entry is $c_n$ and the last $n-i$ entries are 0. Clearly, this claim is valid for $i = 1$, since the last column of $A$ is simply $\begin{pmatrix} c_n \\ 0 \\ \vdots \\ 0 \end{pmatrix}$. Assuming that the last column has the claimed form for $A^i$, then the last column of

$A^{i+1}$ is given by

$$A^{i+1}\, e_n = A\, A^i\, e_n = \begin{bmatrix} c_1 & c_2 & \dots & c_n \\ 1 & 0 & \dots & 0 \\ & \ddots & & \\ 0 & & 1 & 0 \end{bmatrix} \begin{pmatrix} — \\ \vdots \\ — \\ c_n \\ 0 \\ \vdots \\ 0 \end{pmatrix} = \begin{pmatrix} — \\ \vdots \\ — \\ — \\ c_n \\ 0 \\ \vdots \\ 0 \end{pmatrix}$$

as claimed. The claim holds for $i = 1$ through $i = n-1$. For $i = 0$, we want the last column of $I$ which is obviously $e_n$. For (6.11) to be valid, $b_n$ must be 0 since all of the other vectors being added have a 0 as the last component. Hence, (6.11) becomes $b_1 A^{n-1}\, e_n + \cdots + b_{n-1} A\, e_n = 0$, and similarly we can see that $b_{n-1} = 0$. Continuing this argument, $b_1 = b_2 = \ldots = b_n = 0$. So the first $n$ powers of a companion matrix are linearly independent and equating coefficients of these powers was a valid operation and there is no logical hole in the proof of Theorem 6.4.1.

### *6.4.1 Matrices with Repeated Eigenvalues*

Unfortunately, even companion matrices can fail to have distinct eigenvalues, and since we only get one $P_\lambda$ for each $\lambda$, it may not be possible to represent $I$ as a linear combination of the $P_\lambda$'s. But the last theorem suggests the natural generalization. If $\lambda_1$ is an eigenvalue of multiplicity $m$, then $ch(A)$ is divisible by $(A - \lambda_1 I)^m$. So instead of having a single $P_{\lambda_1}$ given by $\dfrac{ch(A)}{A - \lambda_1 I}$, we can have instead $m$ matrices $P_{\lambda_1}^{(1)}, P_{\lambda_1}^{(2)}, \ldots, P_{\lambda_1}^{(m)}$ so that

$$P_{\lambda_1}^{(j)} = \frac{ch(A)}{(A - \lambda_1 I)^j}\,.$$

This form has the pleasant consequence that

$$(A - \lambda_1 I) P_{\lambda_1}^{(j)} = P_{\lambda_1}^{(j-1)} \text{ for } j > 1$$

and, of course, $(A - \lambda_1 I) P_{\lambda_1}^{(1)} = 0$.

As before, to study the behavior of powers of $A$ we look at an expansion of $I$. To keep the notation simple, we will write the $P_\lambda$'s without superscripts, and the expansion is:

$$I = \sum_{i=1}^{n} \alpha_i P_{\lambda_i}\,,$$

which looks like the expansion we used before. The extra complication is that the formula for the $\alpha_i$'s is slightly more complicated. If $\lambda_j$ is a root of

multiplicity 1, consider

$$\frac{ch(A)}{A-\lambda_j I}I = \frac{ch(A)}{A-\lambda_j I}\sum \alpha_i P_{\lambda_i} = \alpha_j \frac{ch(A)}{A-\lambda_j I}P_{\lambda_j} = \alpha_j ch'(\lambda_j)P_{\lambda_j}\,,$$

and so $\alpha_j = \frac{1}{ch'(\lambda_j)}$ since $P_{\lambda_j} = \frac{ch(A)}{A-\lambda_j I}$. If $\lambda_i$ is a root of multiplicity $m$, consider

$$\begin{aligned}\frac{ch(A)}{(A-\lambda_i I)^m}I =& \frac{ch(A)}{(A-\lambda_i I)^m}\sum \alpha_j P_{\lambda_j} \\ =& \frac{ch(A)}{(A-\lambda_i I)^m}[\alpha_1 P^{(1)} + \alpha_2 P^{(2)} + \cdots + \alpha_m P^{(m)}]\end{aligned}$$

where $P^{(1)}, \ldots, P^{(m)}$ are the eigenmatrices associated with $\lambda_i$. Of course, the eigenmatrices associated with other eigenvectors are annihilated. Let

$$r(A) = \frac{ch(A)}{(A-\lambda_i I)^m}\,.$$

We have

$$P^{(m)} = r(A)[\alpha_1 P^{(1)} + \alpha_2 P^{(2)} + \cdots + \alpha_m P^{(m)}]\,.$$

Now

$$\begin{aligned}AP^{(1)} &= \lambda_i P^{(1)} \\ AP^{(2)} &= \lambda_i P^{(2)} + P^{(1)}\,, \qquad A^2P^{(2)} = \lambda_i^2 P^{(2)} + 2\lambda_i P^{(1)}\,.\end{aligned}$$

So

$$\begin{aligned}A^l P^{(2)} &= \lambda_i^l P^{(2)} + l\cdot\lambda_i^{l-1}P^{(1)}\,, \quad \text{and} \\ A^l P^{(j)} &= \lambda_i^l P^{(j)} + \binom{l}{1}\lambda_i^{l-1}P^{(j-1)} + \binom{l}{2}\lambda_i^{l-2}P^{(j-2)} + \cdots\end{aligned}$$

Hence

$$\begin{aligned}P^{(m)} = &\alpha_1 r(\lambda_i)P^{(1)} \\ &+\alpha_2[r(\lambda_i)P^{(2)} + r'(\lambda_i)P^{(1)}] \\ &\vdots \\ &+\alpha_m[r(\lambda_i)P^{(m)} + r'(\lambda_i)P^{(m-1)} + r''(\lambda_i)P^{(m-2)} + \cdots]\end{aligned}$$

Equating the coefficients of the $P^{(i)}$'s gives

$$\begin{aligned}
1 &= \alpha_m r(\lambda_i) \\
0 &= \alpha_m r'(\lambda_i) + \alpha_{m-1} r(\lambda_i) \\
0 &= \alpha_m r''(\lambda_i) + \alpha_{m-1} r'(\lambda_i) + \alpha_{m-2} r(\lambda_i) \\
&\vdots \\
0 &= \alpha_m r^{m-1}(\lambda_i) + \alpha_{m-1} r^{m-2}(\lambda_i) + \cdots \alpha_1 r(\lambda_i)\,.
\end{aligned}$$

Since $r(\lambda_i) \neq 0$, this is a triangular set of $m$ linear equations in $m$ unknowns, with a unique solution and this solution can be found by back substitution. We can compute powers of $A$ by

$$A^k = \sum_{i=1}^{n} \alpha_i A^k P_{\lambda_i}$$

but we may have that $AP^{(2)}_{\lambda_i} = \lambda_i P^{(2)}_{\lambda_i} + P^{(1)}_{\lambda_i}$ if there are eigenvalues with multiplicity greater than 1. But these "extra terms" can only grow as a polynomial in $k$ times $\lambda_i^k$. Thus, if we assume that $\lambda_1$ is the eigenvalue of greatest magnitude and that $\lambda_1$ is <u>not</u> a multiple root, then $\lim_{k\to\infty} \frac{A^k}{\lambda_i^k}$ will not have any extra terms because $(\frac{\lambda_i}{\lambda_1})^k$ times any polynomial in $k$ will still go to 0 as $k$ increases. Thus, we are able to state an improved form of Theorem 6.4.1.

**Theorem 6.4.2.** *If $A$ is a companion matrix and $\lambda_1$ the eigenvalue with largest magnitude is a simple root of the characteristic polynomial of $A$, then*

$$\lim_{k\to\infty} \frac{A^k}{\lambda_1^k} = \frac{1}{ch'(\lambda_1)} P_{\lambda_1} \text{ where } P_{\lambda_1} = \frac{ch(A)}{A - \lambda_1 I}\,,$$

*and this limiting matrix has rank 1. That is, there is some vector $X$ so that the limiting matrix can be written as*

$$(X \quad X \quad \cdots \quad X) \begin{bmatrix} b_1 & 0 & \cdots & 0 \\ & b_2 & & \\ & & \ddots & \\ 0 & \cdots & 0 & b_n \end{bmatrix},$$

*or said another way, every column of the limiting matrix is a scalar multiple of any other column of the limiting matrix.*

*Proof.* Let $Z$ be the limiting matrix. Clearly, $Z$ satisfies the equation $AZ = \lambda_1 Z$, but then each column $Z_i$ of $Z$ also satisfies $AZ_i = \lambda_1 Z_i$ and so

$$\begin{aligned} c_1 Z_{i\cdot 1} + c_2 Z_{i\cdot 2} + \cdots + c_n Z_{i\cdot n} &= \lambda_1 Z_{i\cdot 1} \\ Z_{i\cdot 1} &= \lambda_1 Z_{i\cdot 2} \\ \vdots \qquad & \quad \vdots \\ Z_{i\cdot n-1} &= \lambda_1 Z_{i\cdot n} \end{aligned}$$

where $Z_{i\cdot j}$ means the $j^{\text{th}}$ component of the vector $Z_i$. If $Z_{i\cdot n}$ were known then $Z_{i\cdot n-1}, \ldots, Z_{i\cdot 1}$ could be computed. The top equation is redundant and merely says that $\lambda_1$ satisfies the characteristic polynomial. Thus, up to one unknown multiplier, $Z_{i\cdot n}$, this system of equations has a unique solution and the theorem follows. □

## 6.5 Nonnegative Companion Matrices

Although companion matrices capture the *structure* of Leslie matrices, they do not take into account the Leslie assumptions of nonnegativity. In this subsection we will consider the extra properties which follow from the assumption that all entries of a companion matrix are nonnegative.

The characteristic polynomial of a Leslie matrix or a nonnegative companion matrix can be written in the form:

$$ch(\lambda) = \lambda^n - \sum c_i \lambda^{n-i}$$

where each $c_i \geq 0$ and $c_n > 0$. As shown in Section 5.1, such nonnegative polynomials are of two types – the primitive type which has $\gcd\{i|c_i > 0\} = 1$, and the periodic type which has $g = \gcd\{i|c_i > 0\} > 1$.

A primitive polynomial has exactly one positive real root $\lambda_0$, and this root is dominant in the sense that $\lambda_0 > |\lambda_i|$ for any other root $\lambda_i$. A periodic polynomial with period $g$ only has nonzero coefficients at positions which are multiples of $g$, that is, $\lambda$ only appears in the polynomial as $\lambda^g$. For a periodic nonnegative companion matrix, the characteristic polynomial can be viewed as a polynomial $p(x)$ where $x = \lambda^g$ and $p(x)$ is a primitive polynomial with a unique positive real root $x_0$. The positive root of $ch(\lambda)$ is the positive solution to $\lambda^g = x_0$, i.e., $\lambda_0 = x_0^{1/g}$. This $\lambda_0$ is not strictly dominant because $\lambda^g = x_0$ has $g$ roots all of the same magnitude. These equal magnitude roots will give rise to oscillations and asymptotic periodic behavior of period $g$. We will consider these possibilities in the next subsection. Oscillations are also possible in companion matrices without the nonnegative assumption. We will not consider such matrices in detail, but see Exercise 6.19 where you will show that a companion matrix which

is not nonnegative can have periodic behavior without its characteristic polynomial being periodic.

Since a nonnegative companion matrix with a primitive characteristic polynomial has a dominant eigenvalue we can prove a stronger version of Theorem 6.4.2.

**Theorem 6.5.1.** *If $A$ is a nonnegative companion matrix with a primitive characteristic polynomial, then $A$ has a strictly dominant positive eigenvalue $\lambda_0$ and*

$$\lim_{K\to\infty} \frac{A^K}{\lambda_0^K} = \frac{1}{ch'(\lambda_0)} P_{\lambda_0} \quad \textit{where} \quad P_{\lambda_0} = \frac{ch(A)}{A - \lambda_0 I}$$

*and this limiting matrix is a strictly positive matrix with rank* 1.

*Proof.* Nonnegativity implies that $\lambda_0$ is the dominant eigenvalue and from Exercise 5.2 we know that $ch'(\lambda_0) > 0$. Theorem 6.4.2 implies convergence. We are left with showing that that the limiting matrix is strictly positive. By the division algorithm $ch(\lambda)/(\lambda - \lambda_0) = \lambda^{n-1} + g_1\lambda^{n-2} + \cdots + g_{n-1}$ where

$$\begin{aligned} g_1 &= \lambda_0 - c_1 \\ g_2 &= \lambda_0^2 - c_1\lambda_0 - c_2 \\ &\vdots \\ g_{n-1} &= \lambda_0^{n-1} - c_1\lambda_0^{n-2} - \cdots - c_{n-1} \quad = \quad c_n/\lambda_0. \end{aligned}$$

Clearly $g_{n-1} > 0$ and this implies that $g_{n-2}$ through $g_1$ are also positive. (Also, refer to Exercise 5.20.) So $P_{\lambda_0} = A^{n-1} + g_1A^{n-2} + \cdots + g_{n-1}I$ which is a positive sum of nonnegative matrices. But it could happen that every power of $A$ is not strictly positive. If we look at the graph $G_A$ corresponding to $A$ (see Section 7.2.3 for more details on the correspondence between a nonnegative matrix and its graph), then $A_{ij}^l$, the $(i,j)$ entry of the $l^{th}$ power of $A$, will be positive when there is a path from from $v_j$ to $v_i$ of length $l$ in $G_A$. The graph for a nonnegative companion matrix will contain at least the edges in the following diagram




so there will be a path between any pair of vertices and the length of this path is between 0 and $n-1$. Hence, $A_{ij}^l$ will be positive for some $0 \le l \le n-1$, and the positive sum of the powers of $A$ will be strictly positive. □

### *6.5.1 Periodic Nonnegative Companion Matrices*

Now that we understand the convergence properties of *primitive* companion matrices, we would like to consider nonnegative companion matrices which are periodic in the sense that their characteristic polynomials can be written in the form

$$ch(\lambda) = (\lambda^g)^{\frac{n}{g}} - c_g(\lambda^g)^{\frac{n-g}{g}} - c_{2g}(\lambda^g)^{\frac{n-2g}{g}} \cdots - c_{g\ n/g}.$$

That is, these polynomials only have nonzero coefficients at positions which are a multiple of $g$, and they can be viewed as polynomials in $\lambda^g$ rather than as polynomials in $\lambda$. We want to "reduce" the periodic case to the primitive case. By reduce here, we mean that we would like to represent or think about a nonprimitive matrix as several primitive matrices, apply the theorems about primitive matrices, and then put the results together to obtain an analysis of nonprimitive matrices. This idea of reduction is a central unifying concept in all of mathematics, and is also the basis for divide-and-conquer algorithms [89] in computer science. (See also Section 9.4.)

Assume we have a matrix $L$ with $g = 2$. This $g$ is, of course, the greatest common divisor of the cycle lengths in the graph of the matrix. This quantity is also called the **index of imprimitivity**, but it will be easier to simply call it $g$. We can associate a graph $G(L)$ with the matrix

$$L = \begin{bmatrix} 0 & c_2 & 0 & \dots & 0 & c_n \\ 1 & 0 & 0 & \dots & 0 & 0 \\ & & \dots & & & \\ 0 & 0 & 0 & \dots & 1 & 0 \end{bmatrix}.$$

There is an edge from $v_n$ to $v_1$ and a cycle of length $n$ passing through this edge. The assumption that $g = 2$ implies that $n$ is even. So the graph looks like



There may be some other edges from $v_1$ but they will all go to vertices in the second column; that is, vertices with an even index because $g = 2$. Now, let's form a new graph whose edges are the paths of length 2 in $G(L)$:




The old edges from $v_1$ to $v_i$ now appear as edges from $v_1$ to $v_{i-1}$ and as edges from $v_2$ to $v_i$, but there are *no* edges from an even vertex to an odd vertex, and there are *no* edges from an odd vertex to an even vertex. Notice that the original connected graph has been split into two graphs with no edges between them. Further notice that two new graphs are identical (isomorphic). By the usual reasoning about adjacency matrices, $L^2$ is the matrix which corresponds to this graph with two parts. So if we permute the columns of $L^2$ so that the odd positions appear first and we permute the rows in the same way, we will obtain a matrix of the form

$$\begin{bmatrix} A & 0 \\ 0 & A \end{bmatrix} \quad \text{where} \quad A = \begin{bmatrix} f_2 & f_4 & \cdots & f_n \\ 1 & & & 0 \\ & \ddots & & \vdots \\ & & 1 & 0 \end{bmatrix}.$$

So pleasantly enough, $A$ is a companion matrix, and in fact a *primitive* companion matrix because using length two paths divides the cycle lengths by 2, and converts $g = 2$ for $L$ into $g = 1$ for $A$.

At this point, we should recall a few facts about permutation matrices. A **permutation matrix**, $\Pi$, has exactly one 1 in each row and each column and 0's elsewhere. If $A$ is a matrix and $\Pi$ is a permutation matrix, then $A\Pi$ is a matrix with the columns of $A$ permuted. The position of the 1 in the first column of $\Pi$ determines which column of $A$ will appear as the first column of $A\Pi$. Similarly, $\Pi A$ is a matrix with the rows permuted. The position of the 1 in the first row of $\Pi$ will determine which row of $A$ will appear as the first row of $\Pi A$. Now, consider $\Pi^T$, the **transpose** of $\Pi$, the matrix whose rows are the columns of $\Pi$. So $\Pi^T A \Pi$ would permute the columns of $A$ and permute the rows of $A$ in the same way the columns were permuted. Of course, this is exactly the kind of operation we want to carry out on nonprimitive matrices. Finally, consider $\Pi^T \Pi$. The single 1 in the $i^{\text{th}}$ row of $\Pi^T$ matches the single 1 in the $i^{\text{th}}$ column of $\Pi$, and since

there are no other 1's in the same row as this 1, the product matrix will have in its $i^{\text{th}}$ row a 1 only at position $i$. Hence, $\Pi^T\Pi = I$, the identity matrix, that is, the transpose of a permutation matrix is the inverse of the permutation matrix.

For our example matrix $L$ the appropriate $\Pi$ permutes the odd columns in front of the even columns, and

$$\Pi^T L^2 \Pi = \begin{bmatrix} A & 0 \\ 0 & A \end{bmatrix}$$
$$\text{and } L^2 = \Pi \begin{bmatrix} A & 0 \\ 0 & A \end{bmatrix} \Pi^T .$$

Now consider $(\Pi^T L^2 \Pi)(\Pi^T L^2 \Pi) = \Pi^T L^4 \Pi$ since $\Pi^T \Pi = I$. Thus,

$$\Pi^T L^{2K} \Pi = \begin{bmatrix} A^K & 0 \\ 0 & A^K \end{bmatrix}.$$

Of course, $A$ satisfies the hypothesis of the convergence Theorem 6.5.1, so

$$\lim_{K\to\infty} \frac{A^K}{\mu_0^K} = \frac{1}{ch'_A(\mu_0)} \frac{ch_A(A)}{A - \mu_0 I}$$

where $ch_A(\lambda)$ is the characteristic polynomial of $A$ and $\mu_0$ is the unique positive root of this polynomial. Most conveniently, $ch_A(\lambda^2) = ch_L(\lambda)$; that is, replacing $\lambda^2$ by $\lambda$ gives the characteristic polynomial for $A$. This also means that the dominant eigenvalue $\mu_0$ for $A$ is the square of the dominant eigenvalue $\lambda_0$ for $L$. Hence,

$$\lim_{K\to\infty} \frac{L^{2K}}{\lambda_0^{2K}} = \Pi \begin{bmatrix} \lim_{K\to\infty} \frac{A^K}{\mu_0^K} & 0 \\ 0 & \lim_{K\to\infty} \frac{A^K}{\mu_0^K} \end{bmatrix} \Pi^T$$
$$= \frac{1}{ch'_A(\mu_0)} \Pi \left\{ \frac{ch_A(A)}{A - \mu_0 I} \begin{bmatrix} I & 0 \\ 0 & I \end{bmatrix} \right\} \Pi^T$$

where $I$ is an $\frac{n}{2} \times \frac{n}{2}$ identity matrix.

While the above formula gives an answer, the matrix without the $\Pi$'s might be more intuitive:

$$\lim_{K\to\infty} \frac{L^{2K}}{\lambda_0^{2K}} = \begin{bmatrix} \alpha_1 & 0 & \alpha_2 & 0 & \cdots & \alpha_{n/2} & 0 \\ 0 & \alpha_1 & 0 & \alpha_2 & \cdots & 0 & \alpha_{n/2} \\ \alpha_1 B_2 & 0 & \alpha_2 B_2 & 0 & \cdots & \alpha_{n/2} B_2 & 0 \\ 0 & \alpha_1 B_2 & 0 & \alpha_2 B_2 & \cdots & 0 & \alpha_{n/2} B_2 \\ \vdots & \vdots & \vdots & \vdots & & \vdots & \vdots \\ \alpha_1 B_{n/2} & 0 & \alpha_2 B_{n/2} & 0 & \cdots & \alpha_{n/2} B_{n/2} & 0 \\ 0 & \alpha_1 B_{n/2} & 0 & \alpha_2 B_{n/2} & \cdots & 0 & \alpha_{n/2} B_{n/2} \end{bmatrix}$$

Here, we are trying to display the facts that the limiting matrix for $A$ has rank one and that the permutations $\Pi$ and $\Pi^T$ spread out the two copies of the limiting $A$ matrix.

Let us call this limiting matrix $L_\infty$ and look at the action of $L$ on $L_\infty$. Clearly, either by the derivation or direct calculation

$$\frac{L^2}{\lambda_0^2} L_\infty = L_\infty \, .$$

But what about $\frac{L}{\lambda_0} L_\infty$? This operation results in a matrix which is in a sense complementary to $L_\infty$. That is, the produced matrix, let us call it $L_w$, has 0's where $L_\infty$ has positive elements and $L_w$ has positive elements where $L_\infty$ has 0's. Unfortunately, the values are not quite so nice. In $L_\infty$, each column is either a multiple of the first column or a multiple of the first column shifted down one component. In $L_w$, the odd position columns all start with 0 and are multiples of the first column of $L_w$. The even position columns will all be multiples of the $2^{nd}$ column of $L_w$ and this second column will start with a positive entry, but it is not necessary that the second column is a multiple of a shifted copy of the first column.

The limiting behavior for general $g$ follows immediately from our example with $g = 2$.

**Theorem 6.5.2.** *For a nonnegative companion matrix $L$ with period $g$,*

$$\lim_{K \to \infty} \frac{L^{gK}}{\lambda_0^{gK}} = L_0 = \frac{1}{ch'_A(\lambda_0^g)} \Pi \left\{ \frac{ch_A(A)}{A - \lambda_0^g I} \begin{bmatrix} I & & 0 \\ & \ddots & \\ 0 & & I \end{bmatrix} \right\} \Pi^T$$

*where*

$$A = \begin{bmatrix} f_g & f_{2g} & \cdots & f_n \\ 1 & & & 0 \\ & \ddots & & \vdots \\ & & 1 & 0 \end{bmatrix} \, , \; ch_A(\lambda) = \lambda^{\frac{n}{g}} - f_g \lambda^{\frac{n}{g}-1} - \cdots - f_n \, ,$$

*and $\Pi$ is the permutation matrix which moves the columns with indices $\equiv 1 \bmod g$ before those with indices $\equiv 2 \bmod g, \ldots,$ before those with indices $\equiv 0 \bmod g$.*

a) *$L_0$ has rank $g$;*

b) *If $i \equiv j \pmod g$, the $i^{th}$ column of $L_0$ has positive elements in positions $j$, $j+g$, $j+2g$, $\cdots$ and 0's elsewhere;*

c) *Every column of $L_0$ is a shifted multiple of column 1.*

$$\text{Let } L_r = \left( \frac{L}{\lambda_0} \right)^r L_0 \qquad\qquad 0 \le r < g \, .$$

d) $L_0, L_1, \ldots, L_{g-1}$ *are linearly independent.*

e) *If* $i + r \equiv j \pmod{g}$, *the* $i^{th}$ *column of* $L_r$ *has positive elements in positions* $j$, $j+g$, $j+2g$, $\cdots$ *and* $0$*'s elsewhere.*

f) $\left(\frac{L}{\lambda_0}\right)^g L_r = L_r$.

## 6.6 Back to Leslie Matrices

We have used companion matrices rather than Leslie matrices up to this point because the survival rates in the Leslie matrices give a slight extra complication. Let $B$ be a Leslie matrix

$$B = \begin{bmatrix} f_1 & f_2 & \cdots & f_n \\ s_1 & & & 0 \\ & \ddots & & \vdots \\ & & s_{n-1} & 0 \end{bmatrix},$$

then $B$ can be converted to the companion matrix $L$ via the diagonal matrix $S$ by

$$L = S^{-1}BS \quad \text{where}$$

$$S = \begin{bmatrix} 1 & & & & \\ & s_1 & & & \\ & & s_1 s_2 & & \\ & & & \ddots & \\ & & & & s_1 \ldots s_{n-1} \end{bmatrix},$$

$$S^{-1} = \begin{bmatrix} 1 & & & & \\ & 1/s_1 & & & \\ & & 1/s_1 s_2 & & \\ & & & \ddots & \\ & & & & 1/s_1 \ldots s_{n-1} \end{bmatrix}$$

$$\text{and } L = \begin{bmatrix} f_1 & s_1 f_2 & s_1 s_2 f_3 & \cdots & s_1 \ldots s_{n-1} f_n \\ 1 & & & & 0 \\ & 1 & & & \vdots \\ & & \ddots & & \vdots \\ & & & 1 & 0 \end{bmatrix}.$$

This claim can be easily verified by direct calculation. Of course, we also have $B = SLS^{-1}$, and $B^K = SL^K S^{-1}$. Hence, the asymptotic behavior of $B$ can be directly calculated from the asymptotic behavior of the companion matrix $L$. Each entry of $B^K$ can be obtained from the corresponding

entry of $L^K$ by multiplying and dividing by products of the survival rates. Specifically, the $i,j$ element is obtained by multiplying by $s_1 \dots s_{i-1}$ and dividing by $s_1 \dots s_{j-1}$. So the $i,j^{\text{th}}$ multiplier is

$$\frac{s_1 \dots s_{i-1}}{s_1 \dots s_{j-1}} = \begin{cases} s_j \dots s_{i-1} & \text{if } i > j \\ 1 & \text{if } i = j \\ 1/s_i \dots s_{j-1} & \text{if } i < j\,. \end{cases}$$

If $B$ is a *primitive* Leslie matrix then the corresponding companion matrix $L$ will also be primitive and there will be an analog to Theorem 6.5.1. But $S$ and $S^{-1}$ can be factored across powers and limits and then applied to the limiting matrix. Since the limiting matrix is a polynomial in the companion matrix, applying $S$ and $S^{-1}$ converts this to a polynomial in the Leslie matrix. Hence the convergence theorem for *primitive* Leslie matrices is identical to the convergence theorem for primitive companion matrices:

**Theorem 6.6.1.** *If $A$ is a primitive Leslie matrix, then $A$ has a strictly dominant positive eigenvalue $\lambda_0$ and*

$$\lim_{K\to\infty} \frac{A^K}{\lambda_0^K} = \frac{1}{ch'(\lambda_0)} P_{\lambda_0} \quad \textit{where} \quad P_{\lambda_0} = \frac{ch(A)}{A - \lambda_0 I}$$

*and this limiting matrix is a strictly positive matrix with rank* 1.

### *6.6.1 Periodic Leslie Matrices*

For periodic Leslie matrices we can obtain a convergence theorem by taking Theorem 6.5.2 and multiplying the companion matrices by $S$ and $S^{-1}$. In a moment we will restate that theorem, this time for Leslie matrices rather than companion matrices. An obvious question to ask is why did we bother to analyze companion matrices rather than looking directly at Leslie matrices? The answer is that if we took a Leslie matrix with $g > 1$, took the $g^{\text{th}}$ power, and then permuted rows and columns the resulting matrix would look like

$$\begin{bmatrix} A_1 & & & \\ & A_2 & & \\ & & \ddots & \\ & & & A_g \end{bmatrix}.$$

The matrices $A_1, \dots, A_g$ would be Leslie matrices, but they would not necessarily be identical. The problem area is the survival rates. The survival rates for these matrices can be different. While the $A$ matrices are similar this is somewhat messy to show. By using companion matrices, the corresponding $A$ matrices are identical, because their subdiagonals will only contain 1's.

**Theorem 6.6.2.** *For a Leslie matrix $L$ with index of imprimitivity $g$,*

$$\lim_{K\to\infty}\frac{L^{gK}}{\lambda_0^{gK}} = L_0 = \frac{1}{ch'_A(\lambda_0^g)} S\Pi \left\{ \frac{ch_A(A)}{A-\lambda_0^g I} \begin{bmatrix} I & & 0 \\ & \ddots & \\ 0 & & I \end{bmatrix} \right\} \Pi^T S^{-1}$$

*where*

$$A = \begin{bmatrix} s_1\dots s_{g-1}f_g & s_1\dots s_{2g-1}f_{2g} & \cdots & s_1\dots s_{n-1}f_n \\ 1 & & & 0 \\ & & & \\ & \ddots & & \vdots \\ & & 1 & 0 \end{bmatrix}$$

$$ch_A(\lambda) = \lambda^{\frac{n}{g}} - s_1\dots s_{g-1}f_g\lambda^{\frac{n}{g}-1} - \cdots - s_1\dots s_{n-1}f_n$$

$$ch_A(\lambda_0^g) = 0$$

$$S = diag(1, s_1, s_1s_2, \dots, s_1\dots s_{n-1}),$$

*and $\Pi$ is the appropriate permutation matrix.*

a) *$L_0$ has rank $g$;*

b) *If $i \equiv j \pmod g$, the $i^{th}$ column of $L_0$ has positive entries in positions $j$, $j+g$, $j+2g, \dots$;*

c) *Every column of $L_0$ is a shifted multiple of column 1.*

$$\text{Let } L_r = \left(\frac{L}{\lambda_0}\right)^r L_0\,, \qquad 0 \le r < g.$$

d) *$L_0, L_1, \dots, L_{g-1}$ are linearly independent.*

e) *If $i + r \equiv j \pmod g$, the $i^{th}$ column of $L_r$ has positive elements in positions $j$, $j+g$, $j+2g, \dots$;*

f) $\left(\frac{L}{\lambda_0}\right)^g L_r = L_r$.

Figures 6.1 and 6.2 are from the Census Bureau. The two sides of these graphs show the predicted number of males and females in the United States. Both of these are projections, that is, they are calculated estimates based on a model and on data gathered at an earlier date. Figure 6.1 shows the estimate for 2000, and figure 6.2 shows the estimate for 2050. The age classes start with the youngest on the bottom and proceed to the oldest on the top. In this form, we might expect the graphs to have a pyramid form rather than the inverted pyramid form expected when the oldest age class is on the bottom. Figure 6.1 shows that the U. S.




FIGURE 6.1. Projected U. S. Population in 2000.




FIGURE 6.2. Projected U. S. Population in 2050.

population is not in equibrilium because the "baby boom" is still passing through the population. Figure 6.2 shows that in 50 years, the prediction is that the population will reach a distribution with roughly a pyramid shape. We should mention that the Census Bureau uses models which are more complicated than the simple Leslie model. For example, their models include immigration. In spite of this, an approach to a pyramid shape is still predicted. A conclusion like this, which does not depend on the details of the mathematical model, is called **robust**. Scientists generally have more confidence in robust predictions because they know that they have omitted many details in creating a model.

### *6.6.2 Averaging*

As we have just seen, the asymptotic behavior of a periodic Leslie matrix is more complicated than the straightforward convergence result for primitive matrices. Here, we want to look at the "average" behavior of a periodic matrix over its period, and show that we can obtain a convergence theorem similar to that for primitive matrices.

**Theorem 6.6.3 (Averaging Theorem).** *If $L$ is an $n \times n$ is a Leslie matrix (or a nonnegative companion matrix) with period $g > 1$, then*

$$\lim_{t\to\infty} \frac{1}{g}\sum_{i=1}^{g} \frac{L^{t+i}}{\lambda_0^{t+i}} = \frac{1}{ch'(\lambda_0)}P_{\lambda_0} = \frac{1}{ch'(\lambda_0)}\frac{ch(L)}{L-\lambda_0 I}$$

$$\begin{aligned} \textit{where } P_{\lambda_0} =& L^{n-1} + \lambda_0 L^{n-2} + \cdots + \lambda_0^{g-1}L^{n-g} + (\lambda_0^g - c_g)L^{n-g-1} \\ &+ \cdots + (\lambda_0^{n-1} - c_g\lambda_0^{n-g-1} - \dots)I \end{aligned}$$

*which is a strictly positive matrix of rank* 1.

*Proof.* As we saw in Theorem 5.1.4, $L$ has $g$ roots of maximum modulus, namely, $\lambda_0, \omega\lambda_0, \dots \omega^{g-1}\lambda_0$ where $\omega$ is a principal $g^{th}$ root of unity. For each of these simple roots there is an eigenmatrix $P_j$ so that $LP_j = \lambda_0\omega^j P_j$. As in previous arguments, we can expand $I$ in terms of $L$'s eigenmatrices which are again linearly independent. Applying $(L/\lambda_0)^t$ and taking the limit, all terms except those corresponding to maximal modulus eigenvalues will

disappear. So we need only look at

$$\begin{aligned}
&\frac{1}{g}\sum_{i=0}^{g-1}\left(\frac{L}{\lambda_0}\right)^i\sum_{j=0}^{g-1}\alpha_j P_j \\
&\quad=\frac{1}{g}\sum_{j=0}^{g-1}\alpha_j\sum_{i=0}^{g-1}\left(\frac{L}{\lambda_0}\right)^i P_j \\
&\quad=\frac{1}{g}\sum_{j=0}^{g-1}\alpha_j\sum_{i=0}^{g-1}\omega^{ij}P_j \\
&\quad=\frac{1}{g}\sum_{j=0}^{g-1}\alpha_j P_j\sum_{i=0}^{g-1}\omega^{ij}.
\end{aligned}$$

Now since $\omega$ is a principal $g^{th}$ root of unity.

$$\sum_{i=0}^{g-1}\omega^{ij}=\begin{cases}0 & \text{if} \quad j\not\equiv 0 \pmod g\\ g & \text{if} \quad j\equiv 0 \pmod g\end{cases}$$

and the above sum becomes

$$\frac{1}{g}\,\alpha_0 P_0 g \quad = \quad \alpha_0 P_0\,,$$

and by the argument leading up to Theorem 6.4.1 $\alpha_0 = 1/ch'(\lambda_0)$. The argument for Theorem 6.4.2 shows that the limiting matrix has rank 1, and the argument for Theorem 6.5.1 shows that the limiting matrix is strictly positive. □

## 6.7 The Limiting Effect of $L$ on Nonnegative Vectors

Now that we know the limiting form of Leslie matrices, we are in a position to ask about the long term predictions made by the Leslie model. Specifically, if one measures a population and represents this population by the nonnegative vector $X_0$, and also measures the fertility rates and survival rates and uses these measurements to construct the Leslie matrix $L$, then under the assumption that these rates remain constant one can predict that after $k$ time intervals the population should be $L^k X_0$. Of course, it is relatively easy to calculate $L^k X_0$ even for very large $k$ by using a computer. On the other hand, one hopes that the theory developed in the earlier sections of this chapter will lead to predictions that do not require much computation.

Leslie's Convergence Theorem for nonnegative vectors (Theorem 6.2.1) follows immediately from Theorem 6.6.1 on the convergence of primitive matrices. As long as $X_0$ is nonnegative with at least one positive component, $X_t$ will converge to a positive multiple of the stable age distribution. The convergence will be in the absolute error sense if all eigenvalues, except perhaps $\lambda_0$, are less than 1 in absolute value. In general, convergence will only be in the relative error sense. As long as at least one component of $X_0$ is positive, then $L^k X_0$ converges to a positive multiple of the unique positive eigenvector which corresponds to $\lambda_0$. Further, if $\lambda_0 > 1$, the positive eigenvector will have decreasing components and thus display the inverted pyramid form. Each age class in this eigenvector will be less than $1/\lambda_0$ times the previous age class. If $\lambda_0 = 1$, then the components of the eigenvector are nonincreasing, but one would need the assumption that the survival rates are less than 1 to get decreasing age classes. If $\lambda_0 < 1$, then the form of the eigenvector will depend on the ratios between the survival rates and $\lambda_0$. In the "usual" case, one assumes $\lambda_0 > 1$ but that other eigenvalues may have absolute values greater than 1 and concludes that $L^k X_0$ will converge in a relative error sense to an inverted pyramid form.

Theorem 6.5.1 also shows that there is convergence even if $X_0$ is not nonnegative, but the convergence may be to a negative or zero multiple of the stable age distribution. The coefficient $\gamma$ calculated in the Leslie Convergence Theorem will be the appropriate multiplier.

We now want to consider what happens when we have a nonprimitive Leslie matrix. Will oscillations be possible? Will oscillations be necessary? To see what will happen, let us apply $L_0$ from Theorem 6.6.2 to $X_0$. First, $S^{-1}$ is applied to $X_0$ and it changes the values by dividing them by positive quantities. So a positive component of $X_0$ is not changed into a 0 component, and a 0 component is not changed into a positive component. Next, $\Pi^T$ acts by gathering components into $g$ subvectors. Each subvector will consist of those components whose indices were equal modulo $g$ in the original $X_0$. Because of the decomposed structure of the matrix, each of the subvectors is now treated separately. In fact, each of the subvectors is now acted on by the matrix

$$\frac{ch_A(A)}{ch'_A(\lambda_0^g)(A - \lambda_0^g I)},$$

which is the limiting form for the primitive Leslie matrix $A$. So each subvector which contains a nonzero component is taken to a multiple of the unique positive eigenvector of $A$. Next, the permutation matrix $\Pi$ undoes the permutation caused by $\Pi^T$. Finally, the diagonal matrix $S$ multiplies each component by the appropriate product of survival rates.

There are several immediate conclusions from this computation. As long as there is no $i \in \{1, \ldots, g\}$ so that $X_i$, $X_{i+g}$, $X_{i+2g}$, $X_{i+\lfloor \frac{n}{g} \rfloor g}$ are all 0, $L_0 X_0$ will be strictly positive. So as few as $g$ components of $X_0$ need to be

positive to force $L_0X_0$ to be positive. On the other hand, if there is such an $i$ then $L_0X_0$ will have 0's in the locations whose indices equal $i$ mod $g$.

While it is easy to compute the asymptotic period of a Leslie matrix, it is a little more difficult to compute the asymptotic period of the vector $X_t$ because this vector's period will depend on the numerical values of its components and not only on whether or not a component is 0. For example, if $L = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$ then $L^tX$ will have period 3 for most vectors $X$, but $L^tX$ will have period 1 if all the components of $X$ are the same. For instance,

$$L^3 \begin{pmatrix} 3 \\ 2 \\ 1 \end{pmatrix} = L^2 \begin{pmatrix} 1 \\ 3 \\ 2 \end{pmatrix} = L \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix} = \begin{pmatrix} 3 \\ 2 \\ 1 \end{pmatrix} \qquad \text{but} \quad L \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}.$$

For this example, $X_0 = (x_1, x_2, x_3)^T$ will lead to a cycle of period 3 iff at least one of $(1, \omega, \omega^2)X_0 = x_1 + \omega x_2 + \omega^2 x_3$ and $(1, \omega^2, \omega)X_0 = x_1 + \omega^2 x_2 + \omega x_3$ is nonzero, where $\omega$ is a primitive $3^{rd}$ root of unity.

More generally, using the partial sums of the characteristic polynomial from Horner's Method,

$$\begin{aligned} h_0(\lambda) &= 1 \\ h_1(\lambda) &= \lambda - f_1 \\ h_2(\lambda) &= \lambda^2 - f_1\lambda - f_2 s_1 \\ &\vdots \\ h_{n-1}(\lambda) &= \lambda^{n-1} - f_1\lambda^{n-2} - f_2 s_1 \lambda^{n-3} - \ \ldots \ - f_{n-1} s_1 \ldots s_1 , \end{aligned}$$

define the row vectors $R_j$ for $j = 0, \ldots, g-1$ by

$$\begin{aligned} R_0 &= \left(1, \frac{h_1(\lambda_0)}{s_1}, \frac{h_2(\lambda_0)}{s_1 s_2}, \ldots, \frac{h_{n-1}(\lambda_0)}{s_1 \ldots s_{n-1}}\right) \\ R_j &= \left(1, \frac{h_1(\lambda_0 \omega^j)}{s_1}, \frac{h_2(\lambda_0 \omega^j)}{s_1 s_2}, \ldots, \frac{h_{n-1}(\lambda_0 \omega^j)}{s_1 \ldots s_{n-1}}\right). \end{aligned}$$

It is easy to check that $R_jL = \lambda_0 \omega^j R_j$ and that $R_jL^g = \lambda_0^g R_j$. Now, $R_jL^tX_0/\lambda_0^t = (R_j \cdot X_0)\omega^{jt}$ and this is a periodic function of $t$. Let us define $per(j)$ as the least $p > 0$ so that $\omega^{jp} = 1$. In particular, if $j = 0$, $per(j) = 1$ and the above function has period 1, that is, it is a constant function. Notice also, that $per(j) \leq g$ and $per(j)$ will divide $g$. Asympotically $L^tX_0/\lambda_0^t$ is a sum of periodic functions, and its period is the least common multiple of the periods of the functions in the sum. These considerations lead to the following theorem.

**Theorem 6.7.1.** *If $L$ is a Leslie matrix with period $g$ then asymptotically $X_t/\lambda_0^t$ will be a periodic function of $t$ with period $lcm\{per(j) | (R_j \cdot X_0) \neq 0\}$.*

*This period will be a divisor of $g$. (Here, $per(j)$ is the least $p > 0$ so that $\omega^{jp} = 1$ where $\omega$ is a principal $g^{th}$ root of unity.)*

### *6.7.1 The Period of Total Population*

Now that we know that $X_t/\lambda_0^t$ will become periodic, we would like to know if this periodicity will be visible if one observes only the total population rather than each component of the population vector. Here the **total population** is the sum of the components of $X_t$, and can be written as the inner product $TOTAL = (E, X_t)$ where $E$ is the vector of all 1's.

Since we are interested in the asymptotic period, let $\overline{X}_t = \sum_{j=0}^{g-1} \alpha_j \omega^{jt} C_j$, where $\lambda_0$ is the positive real eigenvalue of $L$, $\omega$ is a principal $g^{\text{th}}$ root of unity, and for $\lambda_j = \lambda_0 \omega^j$

$$C_j = (\lambda_j^{n-1}\,,\, \lambda_j^{n-2} s_1\,,\, \lambda_j^{n-3} s_1 s_2\,,\, \ldots\,,\, s_1 \cdots s_{n-1})^T\,.$$

Note that $\overline{X}_t$ will be a periodic function of $t$ and $X_t/\lambda_0^t$ will approach $\overline{X}_t$. The period of $\overline{X}_t$ will divide $g$ and will depend on which of the $\alpha$'s are nonzero, namely, $per(\overline{X}_t) = lcm\{per(j) \mid \alpha_j \neq 0\}$. Similarly,

$$per(TOTAL) = per((\overline{X}_t, C_j)) = lcm\{per(j) \mid \alpha_j(\overline{X}_t, C_j) \neq 0\}$$

and

$$per(\overline{X}_t) = per(TOTAL) \text{ for every } X_0 \iff (\overline{X}_t, C_j) \neq 0 \text{ for all } j = 0, \ldots g-1\,.$$

To see when this will occur, let us define the polynomial $e(\lambda)$ by

$$e(\lambda) = \sum_{i=1}^{n} s_1 \cdots s_{i-1} \lambda^{n-i} = \lambda^{n-1} + s_1 \lambda^{n-2} + \ldots + s_1 \cdots s_{n-1}$$

so that $e(\lambda_j) = (E, C_j)$. To show that $(E, C_j) \neq 0$, we only need to show that $e(\lambda)$ does not have $\lambda_j$ as a root. Since each $\lambda_j$ has modulus $\lambda_0$, if we can show that $e(\lambda)$ has no root of modulus $\lambda_0$ then we can conclude that no $\lambda_j$ is a root of $e(\lambda)$. For this, we use the fairly standard trick of multiplying $e(\lambda)$ by $\lambda - \lambda_0$ and find that

$$\begin{aligned} e(\lambda)(\lambda - \lambda_0) =& \lambda^n - (\lambda_0 - s_1)\lambda^{n-1} - (\lambda_0 - s_2) s_1 \lambda^{n-1} \\ & - \ldots - (\lambda_0 - s_{n-1}) s_1 \cdots s_{n-2} \lambda - s_1 \cdots s_{n-1} \lambda_0\,, \end{aligned}$$

which is a primitive nonnegative polynomial when $\lambda_0 > s_i$ for all $i = 1, \ldots n-1$. Under this assumption, by Corollary 5.1.5 the polynomial $p(\lambda) = e(\lambda)(\lambda - \lambda_0)$ has a unique positive root which is strictly dominant. Since $\lambda_0 > 0$ is a root of $p(\lambda)$, $\lambda_0$ must be this root. Hence, all of the other roots of $p(\lambda)$, which are exactly the roots of $e(\lambda)$, are strictly less than $\lambda_0$ in complex modulus. From this we obtain the following theorem.

**Theorem 6.7.2.** *For a Leslie model the period of total population equals the period of the population vector if*

$$\lambda_0 > \max\{s_1, s_2, \ldots, s_{n-1}\}.$$

For Leslie models we assume that all $s_i \leq 1$, and in most applications the further restriction $\lambda_0 > 1$ applies. So, this theorem suggests that we should see the same period of oscillation in both the population vector and the total population. Because this theorem only gives a *sufficient* condition for the periods to be equal, they may be the same even if the condition of the theorem is not satisfied. The following result gives some other sufficient conditions.

**Corollary 6.7.3.** *Other sufficient conditions for the period of total population to equal the period of the population vector are:*

*(a)* $\lambda_0 \geq \max\{s_1, s_2, \ldots, s_{n-1}\}$ *and* $\gcd(\{j|\lambda_0 > s_j\} \cup \{g\}) = 1$

*(b)* $\lambda_0 < \min\{s_1, s_2, \ldots, s_{n-1}\}$

*(c)* $\lambda_0 \leq \min\{s_1, s_2, \ldots, s_{n-1}\}$ *and* $\gcd(\{j|\lambda_0 < s_j\} \cup \{g\}) = 1$

In actual practice even the total population is difficult to observe, but a weighted total population might be observed. For example, the probability of observing an organism might be correlated with its size and age. So older age classes might be more heavily represented in a sample than younger age classes. Such observations can be modeled by replacing $E$ with an arbitrary nonnegative weight vector $W$. By agruments similar to the above one can show that $W$ must satisfy a very restrictive set of equations to give a period different from that of the population vector. Hence, it is reasonable to expect that the asymptotic periods of the population vector and the weighted population vector will be identical.

## 6.8 Exercises

**Ex 6.1.** Give an inductive proof that Euclid's algorithm correctly computes the greatest common divisor. You should assume that $a$ and $b$ are nonnegative integers. You can use the smaller of the two numbers as your induction variable.

**Ex 6.2.** Show that if $a \leq n$ and $b \leq n$, then Euclid's algorithm will compute the gcd in time $O(\log n)$. (HINT: Assume that $a$ is the smaller of the two numbers. Consider the situation when

$$\begin{aligned}\gcd(a, b) &= \gcd(b \bmod a, a) \\ &= \gcd(a \bmod [b \bmod a], b \bmod a)\end{aligned}$$

and show that $a \bmod [b \bmod a] \leq \frac{a}{2}$.)

**Ex 6.3.** Show that the bound in the previous exercise is worst case optimal by giving an infinite sequence of pairs of integers so that Euclid's algorithm uses $\Omega(\log b)$ steps on the pair $(a, b)$. (HINT: Think Fibonacci.)

**Ex 6.4.** Investigate the use of absolute values with powers of a matrix by considering

$$A = \begin{bmatrix} -1/2 & 1/2 \\ 1 & 0 \end{bmatrix}$$

and showing which of the following have limits: $A^k$, $|A^k|$, $A^k/\lambda_1^k$, $|A^k/\lambda_1^k|$, $|A^k|/\lambda_1^k$, $A^k/|\lambda_1^k|$, $|A^k|/|\lambda_1^k|$. How close is $|A^8/\lambda_1^8|$ to its limit?

**Ex 6.5.** For the Leslie matrix,

$$L = \begin{bmatrix} 0 & 2 & 0 & 0 & 5 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix},$$

determine the least $m_0$ so that $L^{m_0} \gg 0$. Compare your $m_0$ to the upper bound computed from the formula in Theorem 6.1.2.

**Ex 6.6.** For the Leslie matrix,

$$L = \begin{bmatrix} 0 & 2 & 5 & 3 & 4 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 1/2 & 0 & 0 & 0 \\ 0 & 0 & 1/2 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix},$$

determine the least $m_0$ so that $L^{m_0} \gg 0$. Compare your $m_0$ to the upper bound from the formula in Theorem 6.1.2. How does this example differ from the example in Exercise 6.5?

**Ex 6.7.** For the Leslie matrix,

$$L = \begin{bmatrix} 0 & 0 & 0 & 3 & 5 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 1/2 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1/2 & 0 \end{bmatrix},$$

determine the least $m_0$ so that $L^{m_0} \gg 0$. Compare your $m_0$ to the upper bound given by Theorem 6.1.2

**Ex 6.8.** Show that if $L$ is a Leslie matrix with only $f_n > 0$ and $f_{n-1} > 0$, then the least $m_0$ so that $L^{m_0} \gg 0$ is $m_0 = (n-1)^2 + 1$.

**Ex 6.9.** Create a Leslie matrix, $L$, so that for all $i$, $f_i \cdot f_{i+1} = 0$, but $\gcd\{i \mid f_i > 0\} = 1$. Compute the least power $m_0$ so that $L^{m_0} \gg 0$.

**Ex 6.10.** Show that the assumption that $\lambda_1$ is unique is necessary for convergence by considering

$$A = \begin{bmatrix} 2 & -1 \\ 1 & 0 \end{bmatrix}$$

and showing that $\lambda_0 = 1$ is a *double* eigenvalue of $A$. Further show that

$$A^K = \begin{bmatrix} K+1 & -K \\ K & -(K-1) \end{bmatrix} = K \begin{bmatrix} 2 & -1 \\ 1 & 0 \end{bmatrix} - (K-1) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

and so $A^K/\lambda_1^K$ does not have a finite limit.

**Ex 6.11.** Let

$$F = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix},$$

be the Fibonacci matrix.

(a) Compute $ch_F(\lambda)$.
(b) Find the eigenvalue of largest magnitude, $\lambda_1$.
(c) Find $P_{\lambda_1} = ch_F(\lambda)/(F - \lambda_1 I)$.
(d) Use the formula from Theorem 6.4.1 to compute

$$\lim_{k\to\infty} \frac{F^k}{\lambda_1^k}.$$

(e) Use $\lambda_1 + \lambda_2 = 1$ and $\lambda_1\lambda_2 = -1$ to eliminate $\lambda_2$ from your formula.
(f) The limiting matrix should be

$$\frac{1}{\sqrt{5}} \begin{bmatrix} \lambda_1 & 1 \\ 1 & \frac{1}{\lambda_1} \end{bmatrix}.$$

Is this consistent with what you know about the asymptotic form for the Fibonacci numbers?
(g) Calculate the rank of the limiting matrix.
(h) Find a form for the matrix that clearly displays the rank. (Notice that you can multiply each row of a matrix by a different scalar by pre-multiplying the matrix by a diagonal matrix with the scalar multipliers down the diagonal. Similarly, the columns of a matrix can be scaled by post-multiplying by a diagonal matrix.)

**Ex 6.12.** Consider running the Fibonacci sequence in reverse. Use the inverse of the Fibonacci matrix.

(a) Show that the inverse Fibonacci matrix can be transformed by permuting rows and columns to the companion form:

$$\begin{bmatrix} -1 & 1 \\ 1 & 0 \end{bmatrix}.$$

(b) Show that the eigenvalue with largest magnitude is negative.
(c) Let $A$ be the matrix from (a), and let $\lambda_1$ be the eigenvalue of largest magnitude. Does

$$\lim_{k\to\infty} \frac{A^k}{\lambda_1^k}$$

exist?
(d) Compare

$$\lim_{k\to\infty} \frac{A^k}{\lambda_1^k} \quad \text{to} \quad \lim_{k\to\infty} \frac{F^k}{\lambda_1^k}.$$

(e) What does (d) tell you about the growth in magnitude of the Fibonacci numbers with negative indices?

**Ex 6.13.** The **trace** of a square matrix $A$ is defined to be the sum of the entries on its main diagonal. If $G(A)$ is the graph associated with a nonnegative matrix $A$, show that the trace of $A^k$ is nonzero iff there is a cycle of length $k$ in the graph $G(A)$.

**Ex 6.14.** Assume that you have an $O(n^3)$ algorithm which computes the characteristic polynomial of an $n \times n$ matrix $A$. Assume that $A$ is a nonnegative matrix. Let the characteristic polynomial be $\lambda^n + C_1\lambda^{n-1} + \cdots + C_{n-1}\lambda + C_n$. Use the following lemma to construct an $O(n^3)$ algorithm which determines whether or not $A$ is primitive.

**Leverier's Lemma.** *Let $C_1, \ldots, C_n$ be the coefficients of the characteristic polynomial of the matrix $A$. For each $i$, let $S_i$ be the trace of $A^i$. Then,*

$$\begin{bmatrix} 1 & 0 & 0 & \cdots & 0 \\ S_1 & 2 & 0 & \cdots & 0 \\ S_2 & S_1 & 3 & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & 0 \\ S_{n-1} & \cdots & S_2 & S_1 & n \end{bmatrix} \begin{pmatrix} C_1 \\ C_2 \\ \vdots \\ \vdots \\ C_n \end{pmatrix} = - \begin{pmatrix} S_1 \\ S_2 \\ \vdots \\ \vdots \\ S_n \end{pmatrix}.$$

**Ex 6.15.** Prove the Cayley-Hamilton Theorem for companion matrices. That is, show that a companion matrix satisfies its characteristic polynomial. Further show that this is the lowest degree polynomial satisfied by the matrix.

**Ex 6.16.** Consider the Leslie matrix

$$\begin{bmatrix} 1 & 2 \\ 1 & 0 \end{bmatrix}.$$

Find the eigenvectors and the stable age distribution. Does every initial population converge to the stable age distribution? Consider both absolute error and relative error, and show that certain vectors converge in one sense but not in the other.

**Ex 6.17.** Show by example that there are Leslie matrices, $L$, so that that the entries of the limiting matrix $\lim_{k\to\infty} L^k/\lambda_0^k$ are *not* rational functions of the the entries in $L$.

**Ex 6.18.** Many of the results demonstrated for Leslie matrices also hold for positive matrices, i.e., matrices in which every entry is positive. Consider the two positive matrices:

$$A = \begin{bmatrix} 6 & 5 \\ 10 & 11 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 3 & 2 \\ 1 & 2 \end{bmatrix}.$$

Show by calculation that for each matrix, nonnegative initial vectors converge toward the positive eigenvector of the matrix. Is this convergence in the absolute value or the relative value sense? Does either or both of these matrices display the inverted pyramid form expected for Leslie matrices?

**Ex 6.19.** Show that the matrix

$$A = \begin{bmatrix} 1 & -1 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$$

has period 4, but that its characteristic polynomial is *not* periodic. Further, show that for most vectors $X$, $A^K\ X$ has period 4. Show that there are vectors $X$ so that $A^K\ X$ has period 1. Show that there are no vectors $X$ so that $A^K\ X$ has period 2.

**Ex 6.20.** Prove the results about the period of the total population vector given in Corollary 6.7.3.

**Ex 6.21.** Consider the Leslie matrix

$$L = \begin{bmatrix} 0 & f_2 \\ s_1 & 0 \end{bmatrix} \qquad \text{with} \qquad s_1 f_2 = 1.$$

Find conditions on $X_0$ so that $L^t\ X_0$ will oscillate with period 2. Show that there is a weight vector $W$ so that $W\ L^t\ X_0$ will always have period 1 even if $L^t\ X_0$ oscillates with period 2.

# 7
# Matrix Difference Equations

As we saw in Chapter 6 on the Leslie model, elements of a sequence can be vectors instead of the usual real or complex scalars. In this chapter we will consider linear difference equations whose solutions are sequences of vectors. Such equations are often called matrix difference equations because these equations can be written using matrix and vector notation. We will look at homogeneous equations and at the special case in which the matrix is nonnegative. We will discuss the behavior of primitive matrices and also look at Markov chains. We will give an efficient algorithm to determine whether or not a matrix is primitive. After that we will look at inhomogeneous matrix difference equations and see how to reduce them to one-dimensional difference equations.

## 7.1 Homogeneous Matrix Equations

Let us return to the simplest difference equation: $x_{t+1} = ax_t$. When we considered such equations before, $x_t$ was a sequence of (usually) real numbers, and $a$ was a real constant. We now want to allow the elements of the sequences to be vectors, and let the multiplier be a matrix. It is usual to mark these changes by using capital letters for vectors and matrices instead of the lower case letters we use for scalars. So, our homogeneous matrix difference equation is written as

$$X_{t+1} = MX_t \tag{7.1}$$

(We are using $M$ to remind us that this letter stands for a matrix.) For this equation to make sense, the sizes of the vectors and matrices must agree. So, we let the $X$'s be $k \times 1$ vectors, that is, column vectors, and let $M$ be a $k \times k$ matrix. In printing it is more convenient to write row vectors, so we also use $X^T$( the transpose of $X$) as the row vector corresponding to the column vector $X$. To write the components of a vector we use

$$X = (x_1, x_2, \ldots, x_k)^T.$$

As you may notice we use subscripts for double duty, both to indicate a component of a vector, and to indicate the position of a vector in a sequence. When there is a possibility of confusion, we will use function notation for the sequence, for example,

$$X(t) = (x_1(t), x_2(t), \ldots, x_k(t))^T$$

and

$$X(t+1) = (x_1(t+1), x_2(t+1), \ldots, x_k(t+1))^T.$$

So, (7.1)can be written in component form as

$$\begin{aligned} x_1(t+1) &= m_{11}x_1(t) + m_{12}x_2(t) + \cdots + m_{1k}x_k(t) \\ x_2(t+1) &= m_{21}x_1(t) + m_{22}x_2(t) + \cdots + m_{2k}x_k(t) \\ &\cdots \\ x_k(t+1) &= m_{k1}x_1(t) + m_{k2}x_2(t) + \cdots + m_{kk}x_k(t). \end{aligned}$$

As you can see in this form there is one equation for each component and the next value for each component can depend on the values of *all* the components. While the component form makes the equations appear complicated, the matrix form makes the equation appear simple. In fact, in matrix form the solution is also simple. It is:

$$X_t = M^t X_0. \tag{7.2}$$

Unfortunately, this solution doesn't tell us very much, rather it reduces the problem of solving a difference equation to the problem of computing powers of a matrix.

There are at least two obvious ways to compute the solution to (7.2). One way is to start with $X_0$, compute $X_1$ by $MX_0$, compute $X_2$ by $MX_1$, and so forth. This procedure will take about $tk^2$ arithmetic operations to compute $X_t$ because $k^2$ operations would be used to compute each vector matrix product. The other technique is to compute $M^t$ and then compute the one matrix vector product $M^t\, X_0$. Using the classical matrix multiplication method, computing the product of two matrices uses about $k^3$ operations. (But see Chapter 9 for some discussion of other matrix product algorithms.) Thus it might seem to take $t\,k^3$ operations to compute $M^t$, but **repeated**

**squaring** can save many of these operations. For example, $M^{16}$ can be computed by computing $M^2$, then squaring to get $M^4$, then squaring $M^4$ to get $M^8$, and squaring $M^8$ to get $M^{16}$. In general, $M^t$ can be computed using at most $2\log t$ (log is base 2 logarithm) matrix products by using repeated squaring. So as soon as $2\log t$ is less than $t$, repeated squaring will use fewer arithmetic operations. Further, for big enough $t$, $2k^3 \log t < k^2 t$, and so the matrix powering by repeated squaring will use fewer operations than the matrix vector product method.

Another method for computing $M^t$ comes from the following lemma (see Appendix C).

**Lemma 7.1.1.** *If $M$ is a $k \times k$ matrix then there is a polynomial $p$ of degree at most $k$ so that $p(M) = 0$.*

How can we use this polynomial to compute $M^n$? We have

$$M^k = c_1 M^{k-1} + \ \ldots \ + c_{k-1} M + c_k I$$

and obviously

$$M^n = c_1 M^{n-1} + \ \ldots \ + c_{k-1} M^{n-(k-1)} + c_k M^{n-k}.$$

Now, we can interpret this formula as a difference equation with the obvious initial conditions $M^0 = I$, $M^1 = M$, $M^2 = M^2$, $\ldots$, $M^{k-1} = M^{k-1}$. From Chapter 2 we know that the solution to this difference equation can be written in the form

$$M^n = \sum_{i=1}^{k} A_i x_i(n)$$

where the $x_i(n)$ are $k$ linearly independent solutions of $x(n) = c_1 x(n-1) + \ldots + c_{k-1} x(n-(k-1)) + c_k x(n-k)$. What are the $A_i$'s? To make sense of this equation, the $A_i$'s must be matrices and further they must be linear combinations of $I, M, \ldots, M^{k-1}$.

The following example may help to explain the foregoing argument. For the Fibonacci numbers, we can represent the difference equation in matrix form as:

$$\begin{pmatrix} f_n \\ f_{n-1} \end{pmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} \begin{pmatrix} f_{n-1} \\ f_{n-2} \end{pmatrix}$$

and treat this as a matrix difference equation. We want to compute powers of

$$\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}.$$

Here it is easy to check that $M^2 = M + I$ and so $M^n = M^{n-1} + M^{n-2}$. Since $\lambda^2 - \lambda - 1$ has two distinct roots, $\lambda_1$ and $\lambda_2$, we can write the solution of this difference equation for $M^n$ as

$$M^n = A_1 \lambda_1^n + A_2 \lambda_2^n$$

with initial conditions

$$I = M^0 = A_1 + A_2$$
$$M = M^1 = A_1\lambda_1 + A_2\lambda_2$$

which can be solved to give

$$A_1 = \frac{1}{\lambda_1 - \lambda_2}(M - \lambda_2 I)$$
$$A_2 = \frac{-1}{\lambda_1 - \lambda_2}(M - \lambda_1 I)$$

so

$$\begin{aligned} M^n &= \frac{1}{\lambda_1 - \lambda_2}[\lambda_1^n(M - \lambda_2 I) - \lambda_2^n(M - \lambda_1 I)] \\ &= \frac{\lambda_1^n - \lambda_2^n}{\lambda_1 - \lambda_2}M + \frac{\lambda_1^{n-1} - \lambda_2^{n-1}}{\lambda_1 - \lambda_2}I \\ &= f_n M + f_{n-1} I. \end{aligned}$$

As another example of this method, let's compute

$$M^n = \begin{bmatrix} -4 & 9 \\ -4 & 8 \end{bmatrix}^n.$$

Here, $M^2 = 4M - 4I$. So we want any two linearly independent solutions of

$$x(n) = 4x(n-1) - 4x(n-2).$$

Let's take $x_1(n) = n2^n$ and $x_2(n) = (n-1)2^{n-1}$. It's easy to verify that these are two linearly independent solutions to our difference equation. We want to find matrices $A_1$ and $A_2$ so that

$$M^n = n2^n A_1 \quad + \quad (n-1)2^{n-1}A_2.$$

For $n = 1$, we have

$$M^n = \quad M = \quad 12^1 A_1 \quad + \quad (1-1)2^{1-1}A_2 \quad = \quad 2A_1,$$

which gives $A_1 = 1/2M$. For $n = 2$, we have

$$M^n = M^2 = 4M - 4I = 2 \cdot 2^2 A_1 + 2 \cdot 2^1 A_2 = 4M + 2A_2,$$

which gives $A_2 = -2I$. So we have

$$M^n = n2^{n-1}A_1 \quad - \quad (n-1)2^n A_2 \tag{7.3}$$
$$= \begin{bmatrix} (-3n+1)2^n & 9n2n-1 \\ -n2^{n+1} & (3n+1)2^n \end{bmatrix}. \tag{7.4}$$

A different way to compute $M^n$ is to use the **Jordan Canonical Form** stated in the following theorem (see Appendix C).

**Theorem 7.1.2.** *For any complex $k \times k$ matrix $M$, there is a nonsingular matrix $P$ so that*

$$P^{-1}MP = \begin{bmatrix} J_1 & & 0 \\ & \ddots & \\ 0 & & J_j \end{bmatrix}$$

*where each Jordan block $J_i$ has the form* $\begin{bmatrix} \lambda_i & 1 & 0 & & \dots & 0 \\ 0 & \lambda_i & 1 & 0 & \dots & 0 \\ & & \ddots & \ddots & & \\ & & & & & 1 \\ & & & & & \lambda_i \end{bmatrix}$.
*In the simplest case each $J_i$ is $1 \times 1$ and $P^{-1}MP$ is a diagonal matrix.*

We will not prove this theorem here, instead we will focus on how this result can be used to compute $M^n$.

**Lemma 7.1.3.** *If $J$ is a Jordan block then its $n^{th}$ power is*

$$J^n = \begin{bmatrix} \lambda^n & \binom{n}{1}\lambda^{n-1} & \binom{n}{2}\lambda^{n-2} & \dots & \binom{n}{r}\lambda^{n-r} \\ 0 & \lambda^n & \binom{n}{1}\lambda^{n-1} & \dots & \\ & & \ddots & & \vdots \\ & & & & \lambda^n \end{bmatrix}$$

*that is, its $i^{th}$ row-$j^{th}$ column entry is $(J^n)_{ij} = \lambda^{n-(j-i)}\binom{n}{j-i}$ for $j \geq i$ and $(J^n)_{ij} = 0$ for $j < i$.*

Here $\binom{n}{r}$ is the binomial coefficient $\dfrac{n!}{r!(n-r)!}$. So if $D$ is a matrix with Jordan blocks on its diagonal then

$$D^n = \begin{bmatrix} J_1^n & & & \\ & J_2^n & & \\ & & \ddots & \\ & & & J_r^n \end{bmatrix}$$

and from the lemma we can write down the exact form of $D^n$ and by the theorem, $P^{-1}MP = D$, and rearranging $M = PDP^{-1}$. So to compute $M^n$ we can compute $(PDP^{-1})^n$. Luckily, this is easy because

$$(PDP^{-1})^2 = PDP^{-1}PDP^{-1} = PD^2P^{-1}$$

and

$$(PDP^{-1})^n = PD^nP^{-1}.$$

From above we know how to compute $D^n$. Notice that $P$ and $P^{-1}$ do *not* depend on $n$. So all of the dependence on $n$ is in the powers of the Jordan

blocks, and thus the entries of $M^n$ are linear combinations of the entries of the powers of the Jordan blocks.

For the Fibonacci example, the matrix $M$ is $\begin{bmatrix}1 & 1\\ 1 & 0\end{bmatrix}$. The elements of the Jordan blocks are found by finding the solutions $\lambda_i$'s to $\det(M-\lambda I)=0$. These are the familar values $\lambda_1=(1+\sqrt{5})/2$ and $\lambda_2=(1-\sqrt{5})/2$. The columns of the $P$ matrix are found by solving the equations:

$M\begin{pmatrix}P_{11}\\ P_{21}\end{pmatrix}=\lambda_1\begin{pmatrix}P_{11}\\ P_{21}\end{pmatrix}$ and $M\begin{pmatrix}P_{12}\\ P_{22}\end{pmatrix}=\lambda_2\begin{pmatrix}P_{12}\\ P_{22}\end{pmatrix}$.

which gives $\begin{pmatrix}P_{11}\\ P_{21}\end{pmatrix}=\begin{pmatrix}\lambda_1\\ 1\end{pmatrix}$ and $\begin{pmatrix}P_{12}\\ P_{22}\end{pmatrix}=\begin{pmatrix}\lambda_2\\ 1\end{pmatrix}$.

So

$$\begin{aligned}M^n &= PD^nP^{-1}\\ &= \begin{bmatrix}\lambda_1 & \lambda_2\\ 1 & 1\end{bmatrix}\begin{bmatrix}\lambda_1^n & 0\\ 0 & \lambda_2^n\end{bmatrix}\begin{bmatrix}\frac{1}{\lambda_1-\lambda_2} & \frac{-\lambda_2}{\lambda_1-\lambda_2}\\ \frac{-1}{\lambda_1-\lambda_2} & \frac{\lambda_1}{\lambda_1-\lambda_2}\end{bmatrix}\\ &= \frac{1}{\lambda_1-\lambda_2}\begin{bmatrix}\lambda_1^{n+1}-\lambda_2^{n+1} & \lambda_1^n-\lambda_2^n\\ \lambda_1^n-\lambda_2^n & \lambda_1^{n-1}-\lambda_2^{n-1}\end{bmatrix}=\begin{bmatrix}f_{n+1} & f_n\\ f_n & f_{n-1}\end{bmatrix}.\end{aligned}$$

(Note that we have used $\lambda_1\lambda_2=-1$ to simplify the expression.) It is easy to check that this method of calculating $M^n$ gives the same values as the previous method, although the expressions may look a little different.

A comment is in order. These two methods rely on different polynomials. The first method uses the **minimal polynomial**, that is, the lowest degree polynomial which maps the matrix $M$ to the 0 matrix, while the Jordan method uses the **characteristic polynomial** which is obtained by computing $\det(M-\lambda I)$. The characteristic polynomial for a $k\times k$ matrix always has degree $k$, while the minimal polynomial has degree at most $k$. The minimal polynomial is a divisor (a factor) of the characteristic polynomial, that is, there exists a polynomial $q_M(\lambda)$, so that, $ch_M(\lambda)=\min_M(\lambda)q_M(\lambda)$. The distinction between these two polynomials only arises when there are at least two Jordan blocks with the same value of $\lambda$. A easy example may clarify this. Consider the $k\times k$ identity matrix $I$ which is already in Jordan form with $k$ Jordan blocks and each block has $\lambda=1$. The characteristic polynomial here is $(\lambda-1)^k$, but the minimal polynomial is $(\lambda-1)$. Clearly, $I$ satisfies the minimal polynomial because $I-I=0$. Here the minimal polynomial is a *proper* divisor of the characteristic polynomial.

Our two methods for computing $M^n$ will also allow us to estimate the growth of $M^n$. The Jordan form is most useful for this bounding. If $J$ is a Jordan block of size $r$ and eigenvalue $\lambda$, then there is a entry of $J^n$ which is $\binom{n}{r}\lambda^{n-r}$. We can concisely represent this growth as $\Theta(n^r|\lambda|^{n-r})$. The Jordan form of a matrix consists of diagonal Jordan blocks, so each block is raised to the $n^{th}$ power independently of the other blocks. If $D$ is the Jordan form then $D^n$ has an entry which grows as $\Theta(n^r|\lambda_1|^{n-r})$ where $|\lambda_1|$

is the maximum of the absolute values of the eigenvalues and $r$ is the size of the largest Jordan block which contains an eigenvalue of absolute value $|\lambda_1|$. Clearly from our second calculation method, $M^n$ consists of entries which are linear combinations of the entries of $D^n$. So no entry of $M^n$ can grow faster than the fastest growing entry in $D^n$. Unfortunately, this does not insure that $M^n$ has an entry which grows like the largest entry in $D^n$ because there may be cancellations which occur in the linear combinations. On the other hand, the largest entry in $D^n$ is also a linear combination of some of the entries of $M^n$, so at least one of the entries in this sum must be as large as the largest entry in $D^n$. This observation gives:

**Lemma 7.1.4.** *The largest entry in $M^n$ obeys $\max|(M^n)_{ij}| = \Theta(n^{r_1}|\lambda_1|^n)$ where $|\lambda_1|$ is the maximum eigenvalue in absolute value and $r_1$ is the size of the largest Jordan block which has an eigenvalue of absolute value $|\lambda_1|$.*

**Lemma 7.1.5.** *Every entry in $M^n$ can be bounded from above but not necessarily from below, so every entry satisfies $|(M^n)_{ij}| = O(n^{r_1}|\lambda_1|^n)$.*

Let us return to solving equation (7.1). As we know the solution is: $X_t = M^t X_0$. What can we say about the growth of $X_t$? The actual growth will depend on $X_0$. In essence, $X_0$ can pick out some or all of the Jordan blocks. So detailed knowledge will be necessary to get an actual growth rate. But $X_t$ can not grow faster than $M^t$, and we have:

**Theorem 7.1.6.** *The growth of a solution to a matrix difference equation can be bounded from above so that $|x_i(t)|$ , $\max|x_i(t)|$, and $|\sum \alpha_i x_i(t)|$ are all $O(n^{r_1}|\lambda_1|^n)$ where $|\lambda_1|$ is the maximum eigenvalue in absolute value and $r_1$ is the size of the largest Jordan block which has an eigenvalue of absolute value $|\lambda_1|$.*

## 7.2 Nonnegative Matrix Equations

### *7.2.1 Perron-Frobenius Theory*

The Perron-Frobenius Theorem simplifies working with nonnegative matrix difference equations. Provided the matrix is *primitive*, the equations are as simple as the Leslie equations of Chapter 6 and not much more complicated than scalar equations.

**Definition 7.2.1.** A matrix $M$ is **nonnegative** if every element $m_{i,j}$ is $\geq 0$. A matrix $M$ is **primitive** if $M$ is nonnegative and there is a positive integer $t$ so that every element in $M^t$ is strictly positive, which we will write as $M^t \gg 0$.

**Theorem 7.2.1 (Perron-Frobenius).** *If $M$ is a primitive matrix then*
*(a) $M$ has a maximum positive real eigenvalue $\lambda_0$*

*(b)* $\lambda_0$ *has multiplicity 1, (it is a simple root of the characteristic polynomial)*
*(c) for every other eigenvalue* $\lambda_i$, $\lambda_0 > |\lambda_i|$ *(it is strictly dominant)*
*(d)* $\min_i \sum_j m_{i,j} < \lambda_0 < \max_i \sum_j m_{i,j}$
$\min_j \sum_i m_{i,j} < \lambda_0 < \max_j \sum_i m_{i,j}$
*(e) the row and column eigenvectors associated with* $\lambda_0$ *are strictly positive, and are unique up to scaling*
*(f) the sequence* $M^t$ *is asymptotically one-dimensional, its columns converge to the column eigenvector associated with* $\lambda_0$*, and its rows converge to the row eigenvector associated with* $\lambda_0$*.*
*(g)* $\lambda_0 = \max |Mx|/|x|$ *where* $|x| = \sqrt{\sum x_i^2}$*.*

We will not prove this theorem; proofs appear in many places including [8, Chapter 2] and [127, Chapter 1].

### *7.2.2 Applications to Markov Chains*

One of the most important applications of nonnegative matrices is Markov chains. Such chains are used as models in biological, physical, and social sciences. The idea of a **Markov chain** is simple: a system is modeled as a finite set of states and a set of probablities that describe how one state transitions to other states. If the states are assumed to have certain initial probabilities, then one asks how this probability distribution evolves in time and what its asymptotic distribution will be.

We illustrate this with a Markov chain which has three states called 1, 2, and 3. The probability of being in state $i$ is $p_i$ and the probability of transitioning among the states is given by

$$\hat{P} = \begin{bmatrix} 1/2 & 1/3 & 0 \\ 1/2 & 1/3 & 1/2 \\ 0 & 1/3 & 1/2 \end{bmatrix} P = MP$$

where $P$ is the current vector of probabilities and $\hat{P}$ is the next vector of probabilities. These transitions can also be represented by the following labeled graph:



We want to find the long term behavior of this chain, i.e., to calculate $\lim_{t\to\infty} M^t P(0)$. We expect that this behavior should be independent of the initial distribution $P(0)$, and so we want to find $\lim_{t\to\infty} M^t$.

It is a simple matter to compute the characteristic polynomial for $M$ and find $ch_M(\lambda) = \lambda^3 - \frac{8}{6}\lambda^2 + \frac{1}{4}\lambda + \frac{1}{12}$. By the Cayley-Hamilton Theorem, $M$ satisfies its characteristic polynomial, so

$$M^3 = \frac{1}{12}(16M^2 - 3M - I)$$

and, in general

$$M^t = \frac{1}{12}(16M^{t-1} - 3M^{t-2} - M^{t-3}).$$

This is a linear homogeneous difference equation with characteristic polynomial

$$\lambda^3 - \frac{1}{12}(16\lambda^2 - 3\lambda - 1)$$

which has roots 1, $1/2$, $-1/6$.

Therefore the solution $M^t$ can be expressed as a linear combination of powers of these roots. So we express $M^t$ as

$$M^t = C_1(1)^t + C_2(1/2)^t + C_3(-1/6)^t$$

where $C_1$, $C_2$, $C_3$ depend on the initial conditions. Since the initial conditions are $M^0 = I$, $M^1 = M$, $M^2 = M^2$, we expect $C_1$, $C_2$, $C_3$ to be matrices which are linear combinations of $I$, $M$, and $M^2$. Writing these out in expanded form, we have

$$\begin{aligned} M^t =& (c_{12}M^2 + c_{11}M + c_{10}I)(1)^t \\ &+ (c_{22}M^2 + c_{21}M + c_{20}I)(1/2)^t \\ &+ (c_{32}M^2 + c_{31}M + c_{30}I)(-1/6)^t \end{aligned}$$

where the $c_{ij}$'s are scalars. At first glance we have a problem because there are 9 unknown coefficients and we have only 3 initial condition equations. But, if $I$, $M$, and $M^2$ are linearly independent, we really have 9 equations. For example, the first initial condition gives

$$\begin{aligned} I = M^0 =& 0 \cdot M^2 + 0 \cdot M + 1 \cdot I \\ =& (c_{12}M^2 + c_{11}M + c_{10}I)(1)^0 \\ &+ (c_{22}M^2 + c_{21}M + c_{20}I)(1/2)^0 \\ &+ (c_{32}M^2 + c_{31}M + c_{30}I)(-1/6)^0 \end{aligned}$$

and equating the coefficients of $M^2$ gives

$$0 = c_{12} + c_{22} + c_{32},$$

equating the coefficients of $M$ gives

$$0 = c_{11} + c_{21} + c_{31},$$

equating the coefficients of $I$ gives

$$1 = c_{10} + c_{20} + c_{30}.$$

Therefore one equation in matrices becomes, in this example, three equations in scalars. For this example, $I, M, M^2$ are linearly independent, and it is easy to solve for the coefficients and find:

$$\begin{aligned} M^t =& \frac{1}{7}(12M^2 - 4M - I) \\ & - \left(\frac{1}{2}\right)^{t+1}(6M^2 - 5M - I) \\ & + \frac{1}{14}\left(\frac{-1}{6}\right)^t(18M^2 + -27M + 9I). \end{aligned}$$

Taking limits drives both $(\frac{1}{2})^{t+1}$ and $(\frac{-1}{6})^t$ to 0, and so

$$\lim_{t\to\infty} M^t = \frac{1}{7}(12M^2 - 4M - I).$$

Writing out this matrix gives

$$\lim_{t\to\infty} M^t = \frac{1}{7}\begin{bmatrix} 2 & 2 & 2 \\ 3 & 3 & 3 \\ 2 & 2 & 2 \end{bmatrix}.$$

Several important results can now be seen:

(a) the limiting $M^t$ is a matrix of rank 1,
(b) regardless of the initial probability distribution, the limiting distribution is $(2/7, 3/7, 2/7)^T = (p_1, p_2, p_3)^T$,
(c) $\lambda_0 = 1$ is a strictly dominant positive eigenvalue of $M$,
(d) the column eigenvector corresponding to 1 is $(2/7, 3/7, 2/7)^T$,
(e) the row eigenvector corresponding to 1 is $(1, 1, 1)$.

It is easy to see that $M$ is primitive. For instance, $M^2 \gg 0$. So the Perron-Frobenius Theorem applies and many of these results could have been obtained without calculation.

### *7.2.3 Graphs and Matrices*

Many problems about nonnegative matrices can be solved by translating the problem to a problem about graphs. Graphs are objects which consist of vertices and edges between some of these vertices. Small graphs can be represented by drawing diagrams which can be visually inspected for various properties. Often these visual operations can be codified as algorithms

which can be used to determine properties of graphs which are too large to be conveniently drawn. In many cases, these graph based algorithms can determine matrix properties more quickly than algorithms based on matrix operations.

**Definition 7.2.2.** A **(directed) graph** $G = (V, E)$ consists of a finite set $V$ of vertices and an edge set $E$ where $E \subseteq V \times V$. A **(directed) edge** $(v_i, v_j)$ goes from vertex $v_i$ to vertex $v_j$. A nonnegative $n \times n$ matrix $M$ has a **corresponding graph** $G(M)$ in which $V = \{v_1, v_2, \ldots, v_n\}$ and $(v_j, v_i)$ is in $E$ exactly when $M_{i,j} \neq 0$. (While $(v_j, v_i)$ corresponding to $M_{i,j}$ may seem backwards, it is the natural definition when matrix $M$ times vector $X$ is computed as $MX$.) A matrix $M$ with all the non-zero entries replaced by 1's is called the **adjacency matrix** for $G(M)$.

EXAMPLES:

$$M_1 = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \qquad G(M_1) = \qquad v_1 \longleftarrow v_2$$

$$M_2 = \begin{bmatrix} 1 & 0 \\ 1 & 2 \end{bmatrix} \qquad G(M_2) = \qquad \circlearrowleft v_1 \longrightarrow v_2 \circlearrowright$$

This correspondence between a nonnegative matrix and a graph gives a correspondence between matrix multiplication and paths in the graph. A **path** from $v_i$ to $v_j$ is a sequence of vertices $v_i, v_{i1}, v_{i2}, \ldots, v_{ir}, v_j$ which starts at $v_i$, ends at $v_j$, and for each vertex in the sequence there is an edge to the next vertex in the sequence. The **path length** is the number of edges in the path, which is one less than the number of vertices in the path. For example, in $G(M_1)$ there is a path of length 1 from $v_2$ to $v_1$. In $G(M_2)$ there is a path $v_1, v_1, v_2, v_2$ of length 3 from $v_1$ to $v_2$. Notice also that $G(M_1)$ has no paths of length 2 or greater. Our paths allow a vertex or an edge to be repeated even several times. Paths which do not have repeated vertices are called **simple paths**.

Let us consider matrix multiplication. The row times column rule tells us that if $C = A \cdot B$ then

$$c_{ij} = \sum_k a_{ik} b_{kj}.$$

Specializing this formula to the special case when both $A$ and $B$ are the same nonnegative matrix $M$ gives

$$c_{ij} = \sum_k m_{ik} m_{kj}.$$

When will $c_{ij}$ be nonzero? By our assumption that $M$ is nonnegative, each summand is the product of two nonnegative numbers and hence is nonnegative. So the only way for $c_{ij}$ to be 0 is for every summand to be 0. (Otherwise there will be at least one $k$ so that both $m_{ik}$ and $m_{kj}$ are both nonzero and $c_{ij}$ will be nonzero.) In our graph interpretation this means that there is an edge $v_i \leftarrow v_j$ in $G(C)$ exactly when there at least one $k$ so that there is an edge $v_i \leftarrow v_k$ and an edge $v_k \leftarrow v_j$ in $G(M)$. Said another way, there is an edge $v_i \leftarrow v_j$ in $G(C)$ exactly when there is a path of length 2 from $v_j$ to $v_i$ in $G(M)$.

This observation tells us that the powers of the matrix $M$ contain information about the existence or non-existence of paths in $G(M)$. An even stronger relationship can be shown if we assume that $M$ is a **0-1 matrix**, that is a nonnegative matrix in which all positive entries are 1. For a 0-1 matrix, $M$, $\sum_k m_{ik}m_{kj}$ will count the number of paths of length 2 from $v_j$ to $v_i$ in $G(M)$. This result can be generalized as the following lemma states.

**Lemma 7.2.2.** *Let $M$ be a 0-1 matrix. Then the $(i,j)^{th}$ entry of the $L^{th}$ power of $M$ counts the number of paths of length $L$ from $v_j$ to $v_i$ in the associated graph $G(M)$.*

Notice that *paths* rather than *simple paths* are counted. For example, for the graph




we have

$$M = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \qquad M^3 = \begin{bmatrix} 0 & 2 & 0 \\ 2 & 0 & 2 \\ 0 & 2 & 0 \end{bmatrix}$$

and $M^3$ counts the paths of length 3. In particular, $(M^3)_{21} = 2$ says that there are two paths of length 3 from $v_1$ to $v_2$. These (non-simple) paths are

$$v_1 \longrightarrow v_2 \longrightarrow v_1 \longrightarrow v_2$$

and

$$v_1 \longrightarrow v_2 \longrightarrow v_3 \longrightarrow v_2.$$

Notice that we can naturally replace each positive entry with a 1 to convert a nonnegative matrix to a 0-1 matrix. To count paths as in the above lemma, we would use natural number arithmetic, but to study the existence or non-existence of paths we could use **Boolean OR** as addition and **Boolean AND** as multiplication. For the remainder of this section our matrices will be **Boolean**.

Other matrix properties also correspond to graph properties. A graph is **strongly connected** iff for all pairs of vertices $v_i, v_j$ there is a path from $v_i$ to $v_j$. Of course, this statement contains *all*, so it implies that there is also a path from $v_j$ to $v_i$, but there is no assurance that the path from $v_i$ to $v_j$ has the same length as the path from $v_j$ to $v_i$. For example, in the strongly connected graph

$$\begin{array}{ccc} v_1 & \longrightarrow & v_2 \\ \uparrow & & \downarrow \\ v_4 & \longleftarrow & v_3 \end{array}$$

there is a path of length 1 from $v_1$ to $v_2$, but the length of any path from $v_2$ to $v_1$ is at least 3.

What matrix property corresponds to graph strong connectedness? Since, as we have seen, $(M^L)_{ij} > 0$ iff there is a path of length $L$ from $v_j$ to $v_i$, then the existence of paths between every pair of vertices corresponds to

$$\sum_{L \geq 0} M^L \gg 0.$$

(Recall that for the matrix $A$, $A \gg 0$ means that every element of $A$ is positive.) This formula suggests an infinite amount of work, but if there is a path from $v_i$ to $v_j$, then there is a path of length at most $(n-1)$ from $v_i$ to $v_j$, and this formula can be amended to

$$\sum_{L=0}^{n-1} M^L \gg 0.$$

This form suggests a calculation using $n-2$ matrix multiplications and $n-1$ matrix additions. Can this calculation be carried out with fewer matrix operations? Yes, but we need to look at the calculation slightly differently. One way to recast the calculation is to notice that if $c_0, c_1, \ldots, c_{n-1}$ are positive numbers, then

$$\sum_{L=0}^{n-1} M^L \gg 0 \quad \text{iff} \quad \sum_{L=0}^{n-1} c_L M^L \gg 0.$$

So we want to choose these $c_L$'s to make this calculation easier. Now recall that

$$(M+I)^{n-1} = \sum_{L=0}^{n-1} \binom{n-1}{L} M^L I^{n-1-L} = \sum_{L=0}^{n-1} \binom{n-1}{L} M^L.$$

(The Binomial Theorem holds here because $I$ commutes with $M$.) Choosing $c_L = \binom{n-1}{L}$ means that

$$\sum_{L=0}^{n-1} M^L \gg 0 \quad \text{iff} \quad (M+I)^{n-1} \gg 0.$$

Of course, $(M+I)^{n-1}$ can be computed by repeated matrix squarings and some other matrix multiplies. To further simplify this calculation, notice that $(M+I)^{n-1} \gg 0$ iff $(M+I)^n \gg 0$. Instead of $n-1$ one could use $2^r$ where $r = \lceil \log_2(n-1) \rceil$, that is, $2^r$ is the least power of 2 which is greater than or equal to $n-1$. So, $r$ matrix squarings would suffice. Thus, whether or not a graph is strongly connected can be decided in time $O(T(n)\log n)$ where $T(n)$ is the time to compute the product of two $n \times n$ matrices. (We'll discuss matrix multiplication algorithms more fully in Chapter 9.) The best currently known value for $T(n)$ is $O(n^\alpha)$ where $\alpha > 2$. At best this method decides strong connectedness in $O(n^\alpha \log n)$.

A matrix may have many zero entries and these zeros correspond to edges which do *not* exist in the graph. A more compact representation might avoid representing these non-edges and only represent the edges that actually exist. If we think of a vertex $v$, the edges from $v$ tell us which vertices can be reached from $v$ in one step. So, we could represent a graph by a collection of sets with one set for each vertex, where the set for a vertex $w$ would contain the vertices which can be reached in one step from $w$. In fact, the matrix of the graph *is* such a representation. That is, the column corresponding to the vertex $w$ is a **bit vector representation** of the set of vertices that can be reached in one step from $w$. A bit vector represents a subset of $\{1, \ldots, n\}$ by a 0-1 vector with $n$ components. Namely, if $j$ is in the subset then the $j^{th}$ bit in the vector is 1, and if $j$ is not in the subset then the $j^{th}$ bit in the vector is 0. This matrix representation uses $n^2$ bits. Is a more compact representation possible? Yes, if there are not too many edges. This new representation is called the **next node representation**. For each vertex $v$ there is a **list** and each item in the list is a vertex which can be reached from $v$ in one step, that is, there is an edge from $v$ to each vertex on $v$'s list. There is a separate next node list for each vertex. Since the vertices have labels from $\{1, \ldots, n\}$ the name of each vertex can be represented in $O(\log n)$ bits. Because there will be one item in this array of lists for each edge in the graph, the whole structure can be represented in $O(E \log n)$ bits where $E$ is the number of edges in the graph. So the next node representation will be smaller than the matrix representation as long as $E < n^2/\log n$.

We hope to use this next node representation to determine strong connectedness more quickly. Since strong connectedness means that there is path between every pair of vertices, then for any vertex $v$, if the graph is strongly connected there is a path from $v$ to every vertex and also a path from every vertex to $v$. Conversely if there is a vertex $v$ so that there is a path from $v$ to every vertex and also a path from every vertex to $v$, then the graph is strongly connected because for any pair of vertices, say $w$ and $z$, there is a path from $w$ to $v$ and a path from $v$ to $z$, so following these two paths in order gives a path from $w$ to $z$. These observations suggest a strong connectedness algorithm. Pick an arbitrary vertex $v$ and do a search to find all the vertices which can be reached from $v$, and then do a "backward"

search to find all the vertices that can reach $v$. By "reach" we mean that there is path in the graph that can be followed from the starting vertex to the vertex to be "reached".

With the next node representation these searches are easy. One starts at a vertex $v$, and puts all of the next nodes of $v$ into a structure. Then one takes a vertex, say $w$, from the structure and puts into the structure all of $w$'s next nodes which have not previously been put into the structure. This search halts when the structure is empty. The search is a success if each vertex has been put into the structure. It is easy to keep track of which vertices have been in the structure by using a bit array.

This sort of search can be done in $O(E)$ time if we assume that checking and putting a vertex into the structure are unit operations. For the forward search, the next node representation suffices and we can find the next node to place in the structure by following a pointer and then checking the bit array to determine whether this node has already been put in the structure. For the backward search, a previous node representation is needed. Of course, this previous node representation is just the next node representation for the graph in which every edge $v_i \longrightarrow v_j$ has been turned around into the edge $v_i \longleftarrow v_j$. This can be computed quickly by running through the next node list for, say $v$, and putting $v$ on the previous node list for each $w$ which is on the next node list of $v$.

At the end of these two searches one has to check that *all* vertices have been found. A bit vector representation for found vertices will make these checks easy. Overall, each search costs $O(E)$ and there are two $O(n)$ checks. So, assuming that $E > n$, strong connectedness can be determined in $O(E)$ time.

In the next section we will show that the same sort of graph thinking can lead to an efficient algorithm to determine whether or not a matrix is primitive.

### *7.2.4 Comments on Nonprimitivity*

We are interested in *primitive* matrices because then the Perron-Frobenius theorem applies and the difference equations associated with such matrices will have asymptotic one-dimensional behavior when the initial conditions are nonnegative. This is the same result we found for primitive Leslie matrices in Chapter 6. We were also able to show that if a Leslie matrix is not primitive then it is periodic with period $g = \gcd\{i | f_i > 0\}$. For example, if $A$ is a $6 \times 6$ with all survival rates equal to 1, and with $f_2$ and $f_6$ the only positive fertility rates, then $A^6 - f_2 A^2 - f_6 I = 0$ and this polynomial has period 2. The graph corresponding to $A$ is:




while the graph corresponding to $A^2$ is:




These graphs tell us that the population decomposes into two separate populations, one consisting of the odd numbered components and one consisting of the even numbered components. If we use the original matrix, these two populations change places at each step, that is, the odds become evens and the evens become odds. On the other hand if we use the square of the original matrix, then the populations remain separate; the odds stay as odds and the evens stay as evens. Using the squared matrix corresponds to looking at the populations at every two time units instead of looking at them at every time step. For our example, squaring the original $6 \times 6$ matrix and taking only the components in odd rows and odd columns gives the $3 \times 3$ matrix:

$$\begin{bmatrix} f_2 & 0 & f_6 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}.$$

The same $3 \times 3$ matrix is obtained by taking only the even rows and columns. These matrices are identical because the survival rates are all 1. If nonequal survival rates are used these matrices would be:

$$\begin{bmatrix} s_1 f_2 & 0 & s_5 f_6 \\ s_1 s_2 & 0 & 0 \\ 0 & s_3 s_4 & 0 \end{bmatrix} \text{ and } \begin{bmatrix} s_1 f_2 & 0 & s_1 f_6 \\ s_2 s_3 & 0 & 0 \\ 0 & s_4 s_5 & 0 \end{bmatrix}.$$

These matrices look slightly different, but they have the same characteristic polynomial, $\lambda^2(s_1 f_2 - \lambda)$.

In general, for strongly connected nonnegative matrices, if a matrix, $M$, is not primitive, then for the greatest common divisor of the cycle lengths $g$, the matrix $M^g$ will decompose into $g$ primitive matrices. This decomposition will not be as regular as the decomposition for Leslie matrices. In particular, the number of vertices in each component need not be the same, and the decomposition does not have to be periodic across the indices of the matrix. For example, the graph




is strongly connected with $g = 3$. The third power of the corresponding matrix decomposes using the following three sets of indices $\{1, 5\}, \{2, 6\}, \{3, 4, 7\}$. The graphs for this decomposition are




and the corresponding decomposed matrices are

$$\begin{bmatrix} + & + \\ + & 0 \end{bmatrix} \quad \begin{bmatrix} + & + \\ + & 0 \end{bmatrix} \quad \begin{bmatrix} + & 0 & + \\ + & + & 0 \\ 0 & + & 0 \end{bmatrix}.$$

Unfortunately nonnegative matrices can also fail to be primitive in another way. They can also fail to be strongly connected. For Leslie matrices this is no great problem. If an $n \times n$ Leslie matrix has $f_k$ with $k < n$ as its last positive fertility rate, then one can analyze the $k \times k$ submatrix with the first $k$ rows and the first $k$ columns. If this $k \times k$ matrix is primitive, then its powers will converge as specified by the theorems for Leslie matrices in

Chapter 6 and the $k \times k$ submatrix of the $n \times n$ matrix will converge in the same manner, but the last $n-k$ columns of the $n \times n$ matrix will converge in $n-k$ steps to columns consisting solely of 0's. For the solution vector, $X$, the first $k$ components will converge as usual for a $k \times k$ Leslie system. The last $n-k$ components of $X$ will converge like the first $k$ components, but these last components will also be multiplied by appropriate products of survival rates.

The periodic Leslie case is similar. If $g$ is the period, using the $g^{th}$ power of the Leslie matrix will decompose the system into $g$ Leslie systems with $k \times k$ primitive submatrices. These systems will behave as just described.

The case for general nonnegative matrices is, unfortunately, more complicated because there are a myriad of ways in which primitive and periodic blocks may be connected. Let us discuss some of these possibilities in terms of the graphs associated with the matrices. Of course, this short description is insuffient to give more than a flavor of the possible complications. In a simple situation, the graph may be disconnected, that is, there may be a partition of the vertex set into several subsets so that there is *no* path from a vertex in one subset to a vertex in another subset. In this situation, the disconnected components can be analyzed separately. But, a strange thing can happen. One of the components could have an oscillation of period $p$, while another could have an oscillation of period $q$. Viewed together as a single system, there would be an oscillation of period $lcm(p,q)$. So, for example, if $p$ were 2 and $q$ were 3, there would be an oscillation of period 6. This multiplication of periods makes it possible for a system to have a period much greater than the system's dimension. This is very different from *strongly connected* systems in which the period of any oscillation must divide the dimension.

Let us now consider the connected case in which there is a path between every pair of vertices if we ignore the direction of the edges. For example, $v_1 \longrightarrow v_2$ is connected since there is a directed path from $v_1$ to $v_2$ and a path from $v_2$ to $v_1$ where we ignore the direction of the edge. Of course, this example is only connected and not strongly connected. To analyze connected graphs, we make use of an equivalence relation which says that two vertices $v_i$ and $v_j$ are equivalent iff there is a directed path from $v_i$ and $v_j$ and a directed path from $v_j$ and $v_i$. The partition induced by this equivalence relation breaks the graph into **strongly connected blocks**, that is, the vertices in a block and the edges within a block comprise a strongly connected graph. We construct a new graph which has as its vertices these strongly connected blocks and which has an edge from block $B_i$ to block $B_j$ iff there is a vertex in $B_i$ which has an edge to a vertex in $B_j$. This new graph is called a **DAG**, a **directed acyclic graph**, because it has no directed cycles (any directed cycles are inside a block) and the edges between blocks have direction. Two special kinds of blocks need to be singled out. These are the sources and sinks. A **source** is a block with no in-coming edges. Conversely, a **sink** is a block with no out-going edges. In analyzing

a system, the blocks which are important are determined from the initial conditions. A block is **active** if the initial condition for at least one component in the block is positive. (Remember that we are only dealing with nonnegative systems with nonnegative initial conditions.) A block is **eventually active** if either the block is initially active or there is a directed path to the block from an initially active block. Of particular interest are the eventually active sinks. In analogy with fluid flow, we would expect the sinks to contain all of the asymptotic behavior. That is, if we consider the initial conditions to be an initial distribution of fluid in various containers corresponding to blocks, then we expect the fluid to flow down hill and end up in the sinks. Certainly, if a block consists of a single vertex with an out-going edge, we expect the fluid to flow out of this chamber. Of course, there could be an in-coming flow which would keep fluid in this chamber, but we expect that all fluid will eventually flow out. When a block has cycles, we expect the fluid to flow around these cycles, but if there is an out-going edge we expect some of the fluid to flow out, and even the part of the fluid that is recirculating should eventually hit the out-going edge and flow out. So, at least asymptotically we expect the eventual system behavior to be determined by the eventually active sinks. As before, this set of sinks will behave as a disconnected system (unless there is only one eventually active sink) and we can analyze the behavior as we did for disconnected systems. There is only one thing wrong with this picture; it is only valid in restricted situations. This analogy to fluid flow only makes sense if fluid is neither created nor destroyed.

If we assume that our system is a Markov chain then the fluid flow analogy will make sense because the total probability always remains equal to 1. Even in the Markov case we have passed over some difficulties. The asymptotic probability distribution within a sink depends only on the submatrix for the sink if that submatrix is primitive, but the probability of being in a particular sink depends on the initial conditions over the whole system. Further, if the sink is periodic then the maximum period is determined by the sink's submatrix, but the actual period will depend on the initial conditions across the whole matrix. If the system is not a Markov chain, then blocks can have eigenvalues larger than 1 which would correspond to creation of fluid, and some vertices may lack out-edges and self-loops which would correspond to the destruction of fluid. The analysis of such systems would have to take into account both the graphical properties of the matrix and the actual numerical values in the matrix. So, we will leave such systems with the comment that their analyses are complicated.

## 7.3 Algorithms for Primitivity

In this section we will investigate algorithms for determining whether or not a nonnegative matrix is primitive. The most straightforward algorithm is based on the observation that if a power of a nonnegative matrix is positive then all higher powers are also positive. We show that if a power $A$ is positive, then the $(n-1)^2+1^{st}$ power of $A$ is positive.

**Theorem 7.3.1.** *If $A$ is a primitive nonnegative $n \times n$ matrix and $m_0$ is the least nonnegative integer $m$ so that $A^m \gg 0$ then $m_0$ obeys*

$$m_0 \leq (n-2)l + n \leq (n-1)^2 + 1$$

*where $l$ is the length of the shortest cycle in the graph corresponding to $A$.*

*Proof.* If $n = 1$ then $A = [a_{11}]$ with $a_{11} \neq 0$. So, the corresponding graph consists of a single vertex with a self-loop. For such a graph, $l = 1$. Since $A^0 = [1]$, the formulas are correct. We next note that for $n > 1$, $l = n$ means that $A$ is a permutation matrix which could not be primitive. Therefore, $n > 1$ and $l \leq n-1$.

Consider a cycle of length $l$. Since the graph is strongly connected and $l < n$, there is a vertex $v$ on this cycle which has an edge to a vertex not on this cycle. Let $S_i$ be the set of all vertices which can be reached from the vertex $v$ in exactly $il+1$ steps. Clearly $|S_0| \geq 2$ and $S_0 \subseteq S_1 \subseteq \ldots \subseteq S_n$. Also, whenever $S_{i+1} = S_i$ then $S_{i+2} = S_{i+1} = S_i$. Therefore, either $|S_i| = n$ or

$$2 \leq |S_0| < |S_1| < \ldots < |S_i|\,.$$

This implies $|S_{n-2}| = n$, and every vertex is reachable from $v$ in $(n-2)l+1$ steps. If $y$ is any vertex, the strong connectedness implies the existence of an in-coming edge from some $x$ to $y$. Since $x$ is reachable from $v$ in $(n-2)l+1$ steps, $y$ is reachable from $v$ in $(n-2)l+2$ steps. From this argument we see that for every $j \geq 1$ every $y$ is reachable from $v$ in $(n-2)l+j$ steps. To go from any vertex $x$ to any vertex $y$, one can go from $x$ to $v$ and then go from $v$ to $y$. But there is a path of length at most $n-j \leq n-1$ from $x$ to $v$ which means one can go from $x$ to $y$ in $n-j+(n-2)l+j$ steps. This gives the first upper bound. The second upper bound follows from $l \leq n-1$. □

So the simple algorithm is to compute $A^{(n-1)^2+1}$ and check to see whether or not all entries in this computed matrix are positive. The analysis of this algorithm is also simple. Matrix multiplies will be the most time-consuming step, so let $M(n)$ be the time to compute the product of two $n \times n$ matrices. If we compute the power of $A$ by:

```
POWER := A
FOR I := 1 TO (n-1)^2 DO
    POWER := A × POWER
```

then our algorithm will use $O(n^2)$ matrix multiplies and its time complexity will be $O(n^2 M(n))$. But most of these matrix multiplies can be avoided by **fast exponentiation** or **repeated squaring**, that is, by multiplying POWER by POWER rather than by $A$.

```
POWER := A
I := 1
WHILE I < (n − 1)^2 + 1 DO
      I := 2 × I
      POWER := POWER × POWER
```

Clearly the power of $A$ will be doubled on each execution of the WHILE loop. So if the loop is executed $j$ times, then POWER will contain $A^{2^j}$ and $I$ will contain $2^j$. From the loop condition, we have $2^j \geq (n-1)^2 + 1$ and $2^j < 2(n-1)^2$. So $2\log(n-1) + 1 > j > 2\log(n-1)$ and the repeated squaring method has time complexity $O(\log n M(n))$, which is superior to $O(n^2 M(n))$.

At this point we should say something about $M(n)$, the time complexity of multiplying two $n \times n$ matrices. The classical row times column algorithm uses $\Theta(n^3)$ arithmetic operations (additions and multiplications). Is this the best possible? Well, that depends on what kind of elements are in your matrices and on what kinds of operations you are allowed to use. For example, when we are trying to determine if a power of a nonnegative matrix is positive, it might be reasonable to limit the matrix entries to 1, standing for positive, and 0, standing for zero. When we multiply these entries in the usual way, we get 1 iff both factors are 1, the same operation as Boolean "AND." The sum of two entries is positive unless both addends are 0. Since we want to represent all positives by 1, this addition is the same as Boolean "OR." Now if we restrict the operations on matrix elements to "AND" and "OR," and require that our computer program have no branching, then one can show that any such program must have time complexity $\Omega(n^3)$ and the classical method is the best possible. (See [98, V. 2, pp. 159–168].) If one allows branching, then a faster algorithm is possible. The Four Russians' algorithm will compute the logical product of two matrices in $O(n^3/\log n)$ time. (See Aho, Hopcroft, Ullman [2].)

This may require a small explanation. So far the operations OR and AND, or addition and multiplication, are monotone in the sense that $X \geq Y \Longrightarrow f(X) \geq f(Y)$. In general, $X$ and $Y$ will be vectors. Assume that $\geq$ is defined for the scalars, then $X \geq Y$ iff $\forall i\ x_i \geq y_i$. That is, we extend the ordering from scalars to vectors by requiring that the scalar ordering hold on each component of the vectors. Notice that even if the scalar ordering is a total ordering, the vector ordering may only be (and usually is) a partial ordering.

For example, $\{0,1\}$ is totally ordered by $\geq$ defined as

$$0 \geq 0, \quad 1 \geq 0, \quad \text{and } 1 \geq 1\,,$$

but extending this $\geq$ to vectors gives

$$\begin{matrix} 0 \\ 0 \end{matrix} \geq \begin{matrix} 0 \\ 0 \end{matrix} \qquad \begin{matrix} 0 \\ 1 \end{matrix} \geq \begin{matrix} 0 \\ 0 \end{matrix} \qquad \begin{matrix} 1 \\ 0 \end{matrix} \geq \begin{matrix} 0 \\ 0 \end{matrix} \qquad \begin{matrix} 1 \\ 1 \end{matrix} \geq \begin{matrix} 0 \\ 1 \end{matrix} \qquad \begin{matrix} 1 \\ 0 \end{matrix} \geq \begin{matrix} 1 \\ 0 \end{matrix},$$

which does not specify any relationship between

$$\begin{matrix} 0 \\ 1 \end{matrix} \quad \text{and} \quad \begin{matrix} 1 \\ 0 \end{matrix}.$$

Similarly, when the usual ordering on the reals is extended to vectors, we get

$$\begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \geq \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \quad \text{iff} \quad x_1 \geq y_1 \text{ and } x_2 \geq y_2\,,$$

and we get no relationship if $x_1 > y_1$ and $y_2 > x_2$.

When we discuss the usual sort of operations, $X$ and $Y$ are vectors with two components and $f(X)$ and $f(Y)$ are scalars, so we could rewrite the monotonicity condition as

$$x_1 \geq y_1 \text{ and } x_2 \geq y_2 \Longrightarrow f(x_1, x_2) \geq f(y_1, y_2)\,.$$

For example,

$$x_1 \geq y_1 \text{ and } x_2 \geq y_2 \Longrightarrow (x_1 \text{ OR } x_2) \geq (y_1 \text{ OR } y_2)\,.$$

Now the question is: Will using nonmonotonic operations allow matrix multiplication to be calculated more quickly? The answer is "YES," as shown by Strassen [137] in 1969. (We will return to **Strassen's method** in Chapter 9.) He showed that, by using subtraction of matrices, he could produce a divide-and-conquer algorithm which used only 7 half-size multiplies, rather than the 8 half-size multiplies used by a divide- and-conquer algorithm which uses only addititon and multiplication. Strassen's algorithm has time complexity $\Theta(n^{\log 7})$. A number of researchers have found very clever ways to further reduce the complexity of matrix multiplication. At present, the best algorithm has time complexity $\Theta(n^{\alpha})$ where $\alpha$ is less than 2.3.

These faster matrix multiplication methods could be used to produce a faster algorithm for determining if a matrix is primitive. Of course, to use the methods requiring subtraction, computation could be carried out in the integers. While this would add some complexity because the size of the integers could increase, it can be shown that this increase is not very significant. Are these "faster" methods practical? Since complexity analysis is only asymptotic, it is very possible that a "faster" method is slower than

standard methods for all reasonably sized problems. For Strassen's method, $n > 1000$ is needed to be competitive with the classical method. For other nonmonotonic methods, much larger values of $n$ seem to be needed for there to be any speed-up over the classical method. On the other hand, the Four Russians' method may be competitive for reasonable values of $n$ (say, $n \sim 100$) if some of the operations are implemented as bit vector operations.

After all this talk about matrix multiplication, might there be other methods for testing primitivity which don't use matrix multiplication? Yes, there are methods which are based on calculation of graph properties. The basic technique used in these methods is **depth-first search**. In this technique, one explores all edges in a graph by following each edge which leads to an unused vertex and backing up when there are no edges to unused vertices. The complexity of depth-first search is $O(E)$ where $E$ is the number of edges in the graph. In our matrix application, $E$ will be the number of positive entries in the matrix and so $O(E)$ will be $O(n^2)$.

To determine if a graph (directed graph) is strongly connected one can do two depth-first searches, one on the graph and one on the reversed graph. (In the reversed graph each directed edge is turned around so that the tail of the original edge becomes the head of the new edge, and the head of the original edge becomes the tail of the new edge.) The graph is strongly connected iff both these searches find all the vertices. So strong connectedness can be tested in $O(E)$. Of course breadth-first search could be used here in place of depth-first search.

Recall that nonnegative matrix, $M$, is **primitive** if and only if there is a positive integer $k$ so that $M^k \gg 0$, that is, so that every element of $M^k$ is strictly positive. If $M^k \gg 0$ the corresponding graph has at least one path of length $k$ between every pair of vertices. In particular, for every vertex there is a cycle of length $k$ from that vertex back to that vertex. Further, since $M^{k+1}$ is also strictly positive there is a cycle of length $k+1$ around each vertex. These observations lead to the following theorem.

**Theorem 7.3.2.** *A nonnegative matrix $M$ is primitive iff the corresponding graph is strongly connected and has two relatively prime cycles.*

*Proof.* Since $k$ and $k+1$ are relatively prime, the above observations show the *only if* part. For the *if* part, if there are two relatively prime cycles, then every positive integer greater than, say $B$, can be represented as a positive linear combination of the lengths of these two cycles. (See Exercise 5.5.) Since the graph is strongly connected there are paths between every pair of vertices. If the graph has $n$ vertices, then take $k = 3(n-1) + B$ and we claim that there is a path of length $k$ between any pair of vertices. A path from $v$ to $w$ can be constructed by going from $v$ to $x_1$ on the first cycle, then around this cycle as many times as necessary, then from $x_1$ to $x_2$ on the second cycle, then around the second cycle as many times as necessary,

and finally from $x_2$ to $w$. The paths from $v$ to $x_1$, $x_1$ to $x_2$, and $x_2$ to $w$ have total length at most $3(n-1)$. And going around the two cycles the appropriate number of times will result in a path of length $3(n-1)+B$ because any number greater than or equal to $B$ can be created by sums of the two cycle lengths. This proves there is a path of length $k$ between every pair of vertices, and $M^k \gg 0$. □

It is, perhaps, easy to misread this theorem. It does not say that there are two *simple* cycles whose lengths are relatively prime. For example, one can construct (refer to Exercise 7.2) a strongly connected graph on 20 vertices with only three simple cycles which have lengths $6, 15$, and 20, and although this graph does *not* have two relatively prime simple cycles, the corresponding matrix is still primitive because one can produce a cycle of length 21 by going around the 6 cycle and the 15 cycle. The correct corollary is:

**Corollary 7.3.3.** *A nonnegative matrix is primitive iff the corresponding graph is strongly connected and the greatest common divisor of the lengths of its simple cycles is* 1.

These relatively prime cycles may be relatively long. So instead of trying to find them, our strategy is to find the lengths of small cycles and then to determine if the greatest common divisor of these lengths is 1. Ideally, we would find the lengths of all simple cycles, i.e. those with no repeated vertices. Unfortunately while this is possible, it seems computationally difficult. In particular, determining if an $n$ vertex graph has a simple cycle of length $n$ is the famous **Hamiltonian circuit problem** (refer to [59]) which is known to be $NP$-complete. So, we will look for the lengths of short cycles which may have repeated vertices.

Our procedure starts by picking an arbitrary vertex $v$, and then doing a backward search and a forward search from $v$. As seen above, this pair of searches can check that the graph is strongly connected because each vertex will appear in both the forward search and in the backward search exactly when the graph is strongly connected.

We now need a procedure for finding cycle lengths. Consider our search to be breadth-first, so for each vertex we can talk of a successor set of vertices and a predecessor set of vertices. For each vertex $w$, let $P(w)$ be the length of the shortest path from $w$ to $v$. Since there is a path of length 0 from $v$ to $v$, we let $P(v) = 0$. This assignment can be calculated by doing a backward breadth-first search from $v$. Such a search can be done in $O(E)$ where $E$ is the number of edges in the graph. For the forward search, define $S_0 = \{v\}$ and calculate inductively

$$S_{i+1} = Succ(S_i) - \textbf{ Already Visited Vertices}$$

and for each $w \in Succ(S_i)$ put $P(w)+i+1$ into $C$ the set of cycle lengths. Here, of course, we are using $Succ(X)$ to mean the set of vertices which

can be reached by a path of length 1 from a vertex in the set $X$. We remove the **Already Visited Vertices** so that edges will not be used repeatedly and so this forward search can be accomplished in $O(E)$ time.

We now need to argue that $\gcd(C)$ is the greatest common divisor of all cycle lengths.

**Definition 7.3.1.** If $C$ is a set of natural nembers then the $\pm$ **closure** of $C$ is the smallest set $C^{\pm}$ which contains $C$ and is closed under addition and subtraction, that is, if $a \in C^{\pm}$ and $b \in C^{\pm}$, then $a+b \in C^{\pm}$, if $a \in C^{\pm}$ and $b \in C^{\pm}$, then $|a-b| \in C^{\pm}$.

**Lemma 7.3.4.** *Let $g$ be the greatest common divisor of a set $C$ of natural numbers, then $g$ is also the greatest common divisor of the set $C^{\pm}$. Also, $C^{\pm} = g\mathbb{N}$, that is $C^{\pm}$ consists of all multiplies of $g$.*

*Proof.* The elements of $C^{\pm}$ can be assigned types indicating the number of operations needed to create the element from the elements of $C$. Elements of $C$ are type 0 elements of $C^{\pm}$, and $x$ is an element of type $K$ if $x = a+b$ or $x = |a-b|$ where one of $a$ and $b$ has type $K-1$ and the other has type at most $K-1$. By hypothesis, $g$ is the greatest common divisor of elements of type 0. Assume that $g$ is the greatest common divisor of elements of type at most $K-1$. If $g$ divides both $a$ and $b$, then $g$ also divides their sum and their difference and hence $g$ divides all elements of type at most $K$, and since adding elements to a set can not increase the greatest common divisor, $g$ is the greatest common divisor of elements of type at most $K$. Since every element of $C^{\pm}$ has finite type, $g$ is the greatest common divisor of $C^{\pm}$. Why is $C^{\pm} = g\mathbb{N}$? By Euclid's algorithm $g \in C^{\pm}$. By closure under addition all multiplies of $g$ are in $C^{\pm}$. By closure under subtraction, 0 is in $C^{\pm}$. And since $g$ is the greatest common divisor of $C^{\pm}$, all elements of $C^{\pm}$ are multiplies of $g$. □

We proceed by showing that all cycles lengths are in $C^{\pm}$. If we can show this then we don't need to construct $C^{\pm}$, but instead we can find the greatest common divisor of $C$, and that will be the greatest common divisor of all cycle lengths.

**Lemma 7.3.5.** *If $C$ is the set of numbers found by the primitivity algorithm then the length of every cycle is in $C^{\pm}$.*

*Proof.* Consider any cycle, simple or non-simple. Let $v_1, v_2, \ldots, v_L$ be the vertices in this cycle of length $L$. For every $v_i$ in this cycle there will be an edge $(v_i, v_{i+1 \pmod L})$. Let $P(v_i)$ be the position at which $v_i$ is found in the backward search, and let $Q(v_i)$ be the position at which $v_i$ is first found in the forward search, then both

$$P(v_i) + Q(v_i) \quad \text{and} \quad P(v_{i+1}) + Q(v_i) \; + \; 1$$

are put into $C$ because from the edge $(v_i, v_{i+1 \pmod L})$, $v_{i+1}$ is found on the step after $v_i$. Since $C^{\pm}$ is closed under addition and subtraction, the following is in $C^{\pm}$:

$$\sum_{i=1}^{L}[P(v_{i+1}) + Q(v_i) \ + \ 1] \quad - \quad \sum_{i=1}^{L}[P(v_i) + Q(v_i)].$$

But this sum is $L$, the cycle length, and every cycle length is in $C^{\pm}$. □

**Theorem 7.3.6.** *If $C$ is the set of numbers found by the primitivity algorithm then* $\gcd(C)$ *is the greatest common divisor of the cycle lengths of the input strongly connected graph. In particular, the strongly connected graph is primitive if and only if* $\gcd(C) = 1$.

*Proof.* By Lemma 7.3.4, $C^{\pm}$ contains the length of every cycle in the graph. But, by Lemma 7.3.5, $\gcd(C) = \gcd(C^{\pm})$, and so $\gcd(C)$ is the greatest common divisor of the cycle lengths of the graph. By Theorem 7.3.2 a strongly connected graph is primitive iff the greatest common divisor of its cycle lengths is 1. □

To finish up this algorithm, we need to look at the complexity of finding the GCD of a set of numbers. The obvious way to compute GCD is to start with with the two smallest numbers, use Euclid's algorithm to find their GCD, replace these elements with their GCD, and apply this procedure recursively to the remaining set. If there are $n$ elements in the set, there will be $O(n)$ calls to Euclid. If the elements are less than $2n$ (as they will be in our application), each call to Euclid could use $O(n)$ divisions. In total this procedure could take time $O(n \log n)$. This is an overestimate. On each division, either the division is exact and the larger number is eliminated, or one of the numbers is decreased. In fact, if the division is not exact the smaller number is decreased by a factor of least 2 in every two divisions. Since the smallest number only has $\log n$ bits. It will be reduced to 1, if there are $2 \log n$ divisions which are not exact. Hence, the GCD of a set of $n$ integers from $\{1, \ldots, 2n\}$ can be computed in $O(n)$ time. Coupled with the above this gives:

**Theorem 7.3.7.** *Whether or not an $n$ vertex graph is primitive can be determined in $\Theta(E + n)$ where $E$ is the number of edges in the graph.*

**Theorem 7.3.8.** *Whether or not an $n \times n$ nonnegative matrix is primitive can be determined in $\Theta(n^2)$. Except for an initial scan of the matrix, the algorithm runs in $\Theta(E)$ where $E$ is the number of nonzero entries in the matrix.*

Let us finish this section with a couple of comments. If a graph has many edges, then the graph is almost surely primitive. With some effort one could calculate a constant $\gamma$ and show that if $E > \gamma n^2$, then the graph

is primitive. It can also be shown that there is a constant $\alpha$, so that if each vertex in the graph has degree $> \alpha \log n$, then if the edges are assigned at random the graph is almost surely primitive [11]. In graphs that arise in modeling, the structure of the graph may allow a fast proof of primitivity without going through our algorithm. The point is that our complexity estimates may be overestimates. Finally, the algorithm lets one pick $v$. Which $v$ should you pick? A reasonable choice is to choose the $v$ with the largest number of edges going out from $v$.

## 7.4 Matrix Difference Equations with Input

In the previous sections of this chapter we have considered matrix difference equations of the form

$$X_{t+1} = MX_t.$$

These are called *homogeneous* because multiplying each component of $X_t$ by $\alpha$ will result in each component of $X_{t+1}$ being multiplied by $\alpha$. The solutions to such equations also satisfy the *additive* property in that if $X_{t+1} = MX_t$ and $W_{t+1} = MW_t$ then letting $Z_t = X_t + W_t$ gives $Z_{t+1} = MZ_t$. While this additive property holds for the difference equation, it does *not* hold for the initial conditions because if $X_0 = C$ and $W_0 = C$ then $Z_0 = X_0 + W_0 = 2C \neq C$ if $C \neq 0$. Homogeneous difference equations are used to model systems with *no* input. The current state of such a system depends only on the previous state of the system. In contrast, many other systems are better described by inhomogeneous equations because the state of the system will depend on the input as well as on the internal workings. For example, the states of a sewage system will depend on what is fed into the system as well as on the internal processing within the system.

We will describe inhomogeneous systems by the matrix difference equation:

$$X_{t+1} = MX_t + Y_t. \tag{7.5}$$

But, in many fields more complicated systems models are used. For example, control engineers often use a pair of equations:

$$X_{t+1} = MX_t + BY_t$$
$$Z_{t+1} = CX_t + DY_t$$

where $X_t$ is the internal state, $Y_t$ is the input, and $Z_{t+1}$ is the output, and ask such questions as: What input sequence will produce a desired output sequence? Of course, even more complicated models including nonlinear models are used in a variety of fields. Here we will restrict ourselves to equations in the form of equation (7.5).

Our first result is a trivial observation, but it may seem surprising.

**Theorem 7.4.1.** *Let $X_{t+1} = MX_t + Y_t$, then the input sequence $Y_t$ can be chosen so that the solution $< X_t >$ is any desired sequence.*

*Proof.* If $X_0$ is the initial condition and $X_1$ is the next desired value then taking $Y_0 = -MX_0 + X_1$ gives $X_1 = MX_0 + Y_0 = MX_0 - MX_0 + X_1 = X_1$. Similarly, if $X_t$ was the last value and $X_{t+1}$ is the next desired value then taking $Y_t = -MX_t + X_{t+1}$ gives $X_{t+1}$ the desired value. □

This result may seem surprising because linear systems are so simple that they should have very limited possibilities, and so it's surprising that every desired sequence can be produced from a linear system. On the other hand, the result is trivial because it says that if you can produce any desired input sequence, i.e. any $Y_t$, then you can pass your constructed sequence, which contains all the complexity, through a linear system and obtain whatever you would like. The linear system is *not* adding any complexity or flexibility, you had that when you were constructing the input sequence.

A major virtue of linear systems is that we can write down their solutions. Of course, this form of solution is formal and doesn't really tell us much about how the solution behaves. So after we give the solution, we will then have to add some hypotheses about the structure of the matrices and the growth of the input sequence to be able to get some bounds on the behavior of the solution.

The solution to

$$X_{t+1} = MX_t + Y_t \tag{7.6}$$

with initial condition $X_0$ is

$$X_t = M^t X_0 + \sum_{i=0}^{t-1} M^{t-1-i} Y_i. \tag{7.7}$$

Notice that this is formally the same as the solution to the first order one-dimensional difference equation

$$x_{t+1} = ax_t + y_t,$$

but the meanings of the symbols are different. $X_t$ is now a vector rather than a scalar. $X_0$ is a vector of initial conditions rather than a single scalar initial condition. $Y_t$ is a sequence of vectors rather than scalars. The multiplier is the matrix $M$ rather than the scalar $a$. Another change is that the order of operands is now important. While scalar multiplication is commutative, that is, order doesn't matter, order matters for vectors and matrices. For example, a $k \times k$ matrix times a $k \times 1$ vector must be multiplied in the order matrix times vector.

### *7.4.1 Reduction to One Dimension*

In general multi-dimensional systems are hard to deal with. But linear systems have such nice additive properties that we can hope that multi-dimensional linear systems are not much more complicated than one-dimensional linear systems. We will show that a multi-dimensional linear system can be decomposed into a set of one-dimensional linear systems.

Let us start with special case that often arises in practice.

**Theorem 7.4.2.** *The $k$-dimensional linear difference equation*

$$X_{t+1} = MX_t + Y_t$$

*can be decomposed into the set of one-dimensional difference equations*

$$\begin{aligned}
\hat{x}_1(t+1) &= \lambda_1 \hat{x}_1(t) + \hat{y}_1(t) \\
\hat{x}_2(t+1) &= \lambda_2 \hat{x}_2(t) + \hat{y}_2(t) \\
&\vdots \\
\hat{x}_k(t+1) &= \lambda_k \hat{x}_k(t) + \hat{y}_k(t)
\end{aligned}$$

*if the matrix $M$ has $k$ linearly independent eigenvectors. The $\lambda_1, \ldots, \lambda_k$ are the eigenvalues corresponding to these eigenvectors, and the $\hat{x}$'s and $\hat{y}$'s are linear combinations of the components of $X$ and $Y$ respectively.*

*Proof.* If $Z$ is an eigenvector of $M$, then $MZ = \lambda Z$ where $\lambda$ is the corresponding eigenvalue. If $M$ has $k$ linearly independent eigenvectors $Z_1, Z_2, \ldots, Z_k$ with corresponding eigenvalues $\lambda_1, \lambda_2, \ldots, \lambda_k$ then

$$M[Z_1 Z_2 \ldots Z_k] = [Z_1 Z_2 \ldots Z_k] \begin{bmatrix} \lambda_1 & & & & \\ & \cdot & & & \\ & & \cdot & & \\ & & & \cdot & \\ & & & & \lambda_k \end{bmatrix}.$$

Here $[Z_1 Z_2 \ldots Z_k]$ is the $k \times k$ matrix which has the $i^{th}$ eigenvector as its $i^{th}$ column. The equality follows because post-multiplying a matrix by a diagonal matrix results in each component of the $i^{th}$ column being multiplied by the $i^{th}$ element on the diagonal of the diagonal matrix. Putting this in matrix form, we have

$$M\mathcal{Z} = \mathcal{Z}\Lambda$$

where $\mathcal{Z}$ is the matrix of eigenvectors and $\Lambda$ is the diagonal matrix of eigenvalues. If the eigenvectors are linearly independent, then $\mathcal{Z}$ is an invertible matrix and

$$\mathcal{Z}^{-1} M \mathcal{Z} = \Lambda \quad \text{and} \quad M = \mathcal{Z}\Lambda\mathcal{Z}^{-1}.$$

Replacing $M$ in the matrix equation gives

$$X_{t+1} = \mathcal{Z}\Lambda\mathcal{Z}^{-1}X_t + Y_t$$

and multiplying by $\mathcal{Z}^{-1}$ gives

$$(\mathcal{Z}^{-1}X_{t+1}) = (\Lambda\mathcal{Z}^{-1}X_t) + (\mathcal{Z}^{-1}Y_t).$$

So setting $\hat{X} = (\mathcal{Z}^{-1}X)$ and $\hat{Y}_t = (\mathcal{Z}^{-1}Y_t)$ gives

$$\hat{X}_{t+1} = \Lambda\hat{X}_t + \hat{Y}_t.$$

Because $\Lambda$ is diagonal, each component of this equation is independent of the other components, and so we can write the $k$-dimensional difference equation as a set of $k$ difference equations each of dimension 1. □

Once the matrix difference equation has been reduced to this one-dimensional form, the methods of Chapters 3 and 4 can be used to find the solutions to the 1-dimensional equations, and then assembling these component solutions into a vector and multiplying by the matrix $\mathcal{Z}$ will give the solution $X_t$ to the matrix difference equation.

This reduction can also be described in terms of the eigenvectors and generealized eigenvectors of $M$. As in section 2.3, a matrix has both row and column eigenvectors. A row vector $R_i$ is an eigenvector of $M$ when $R_i\, M \,=\, \lambda_i\, R_i$ where $\lambda_i$ is the corresponding eigenvalue. Starting with the equation

$$X_{t+1} \;=\; MX_t \;+\; Y_t, \tag{7.8}$$

we can multiply by $R_i$ to obtain

$$\begin{aligned} R_iX_{t+1} \;&=\; R_iMX_t \;+\; R_iY_t \\ &=\; \lambda_i\;(R_i, X_t) \;+\; (R_i, Y_t). \end{aligned}$$

Since $(R_i, X_t)$ is one dimensional, let us call it $r_i(t)$, and then we have the one dimensional difference equation

$$r_i(t+1) \;=\; \lambda_i\; r_i(t) \;+\; (R_i, Y_t)$$

which has the solution

$$r_i(t) \;=\; \lambda_i^t\; r_i(0) \;+\; \sum_{j=0}^{t-1} \lambda_i^{t-1-j} (R_i, Y_t).$$

If $M$ has a complete set of eigenvectors, then the solution $X_t$ to (7.8) can be written as as a linear combination of the solutions to a set of one dimensional linear equations. In fact if, $V_1, \ldots, V_k$ are the column eigenvectors of $M$ then

$$X_t \;=\; \sum_{i=1}^{k} r_i(t)\; V_i$$

where each $r_i(t)$ is the solution to one of the one dimensional equations and the $V_i$ are normalized so that $(R_i, V_i) = 1$.

This special situation often arises in modeling physical systems because the laws of physics often lead to real symmetric matrices and such matrices always have a full set of linearly independent eigenvectors. When a matrix does not have a full set of eigenvectors, the above method fails, but we can still fall back on the Jordan form for the matrix. (See Section 7.1 and Appendix C.) Using the Jordan form and following the above construction will lead to a set of difference equations of the form

$$\hat{X}_{t+1} = \begin{bmatrix} \lambda & 1 & 0 & \dots & 0 \\ 0 & \lambda & 1 & \dots & 0 \\ & & . & & \\ & & & . & 1 \\ & & & & \lambda \end{bmatrix} \hat{X}_t + \hat{Y}_t,$$

that is, every matrix difference equation decomposes into a set of matrix difference equations in which the matrices are Jordan blocks. These Jordan blocks can be transformed into companion blocks (see Exercise 7.13). Even more generally, all matrices (even over fields other than the reals or the complexes) are similar to blockwise companion matrices.

**Theorem 7.4.3 (Rational Canonical Form).** *Any square matrix, $M$, over any field, is similar to*

$$\begin{bmatrix} C_{m_1} & & & \\ & C_{m_2} & & \\ & & \ddots & \\ & & & C_{m_k} \end{bmatrix}$$

*where each $C$ is a companion matrix and has the associated polynomial $m_i(x)$, so that $m_i(x)$ divides $m_{i+1}(x)$. Here each $C$ has the form*

$$\begin{bmatrix} c_1 & & \dots & & c_r \\ 1 & & & & 0 \\ & 1 & & & \\ & & \ddots & & \\ & & & 1 & 0 \end{bmatrix}.$$

*Of course different $C$'s may have different $r$'s and different $c_i$'s.*

This is a very powerful theorem because it applies to any field and because all of the operations used to convert a matrix to rational canonical form are rational operations in the field. This is in sharp contrast with the Jordan form which requires one to find the roots of polynomials. In general, this is difficult because some polynomials will *not* have roots in the

field. For this reason, one may start with integer matrices and have to go to complex matrices to obtain the Jordan form. Further, the roots may not be expressible by an algebraic formula. Even though the rational canonical form exists, it may not be easy to find. Algorithms for this problem are known (for example, refer to Harrison [65]), but none is straightforward enough to be included here. Another difficulty of rational form is uniqueness. A matrix may be similar to a variety of rational forms. (See Exercise 7.12.) The above theorem picks one out by a divisibility condition on the polynomials associated with the companion matrices. Of course, one can choose a different set of relations on the polynomials to obtain a different "rational canonical form".

To solve matrix difference equations, all we need is one decomposition into rational form. If such a decomposition is found, then we have a set of matrix difference equations in which the matrices are in companion form. To see how a companion matrix difference equation can be decomposed into one dimensional difference equations, let us consider an example with a $3 \times 3$ companion matrix. The component equations are:

$$\begin{aligned} x_1(t+1) =& c_1 x_1(t) + c_2 x_2(t) + c_3 x_3(t) + y_1(t) \\ x_2(t+1) =& x_1(t) + y_2(t) \\ x_3(t+1) =& x_2(t) + y_3(t). \end{aligned}$$

Rewriting from the bottom these become:

$$\begin{aligned} x_3(t) =& x_2(t-1) + y_3(t-1) \\ x_2(t) =& x_1(t-1) + y_2(t-1) \\ x_2(t-1) =& x_1(t-2) + y_2(t-2). \end{aligned}$$

So,

$$x_3(t) = x_1(t-2) + y_2(t-2) + y_3(t-1)$$

and

$$\begin{aligned} x_1(t+1) =& c_1 x_1(t) + c_2[x_1(t-1) + y_2(t-1)] \\ & + c_3[x_1(t-2) + y_2(t-2) + y_3(t-1)] + y_1(t) \\ =& c_1 x_1(t) + c_2 x_1(t-1) + c_3 x_1(t-2) \\ & + c_2 y_2(t-1) + c_3 y_2(t-2) + c_3 y_3(t-1) + y_1(t). \end{aligned}$$

In this form, we can see that $x_1(t)$ is the solution of a $3^{rd}$-order one-dimensional difference equation. The other components are just shifted versions of the sequence for $x_1$ with some of the input sequence added. For example,

$$x_3(t) = x_1(t-2) + y_2(t-2) + y_3(t-1)$$

which we can regard as a trivial difference equation which has its output, $x_3(t)$, equal to its input because $x_3$ does not appear on the right hand side of the equation.

With this result we have arrived at the theorem:

**Theorem 7.4.4.** *Any $k$-dimensional matrix difference equation can be reduced to a one-dimensional difference equation of order at most $k$.*

### *7.4.2 Reduction to Homogeneous Form*

The matrix difference equation

$$X_{t+1} = MX_t + Y_t \tag{7.9}$$

can be reduced to homogeneous form if the input is well-behaved. Specifically, if there is a homogeneous matrix difference equation $Z_{t+1} = AZ_t$ and $Y_t = PZ_t$, then (7.9) can be written as $\mathcal{X}_{t+1} = \mathcal{M}\,\mathcal{X}_t$, that is

$$\mathcal{X}_{t+1} = \begin{pmatrix} X_{t+1} \\ Z_{t+1} \end{pmatrix} = \begin{bmatrix} M & P \\ 0 & A \end{bmatrix} \begin{pmatrix} X_t \\ Z_t \end{pmatrix}.$$

This allows one to write the solution to (7.9) as

$$X_t = Q\ \mathcal{M}^t \begin{pmatrix} X_0 \\ Z_0 \end{pmatrix}$$

where $Q$ is the projection matrix which returns the first $k$ components of its input vector. This reduction replaces an inhomogeneous equation by a homogeneous equation at the cost of increaasing the size of the homogeneous equation.

Since $\mathcal{M}$ has special form, one can hope that computing powers of $\mathcal{M}$ may be easier than computing powers of a general matrix. If the matrices $M$ and $A$ do *not* have any eigenvalues in common, then an easier computation is possible. If $V_i$ is an eigenvector or generalized eigenvector of $M$ then $(V_i^T, 0^T)$ will be a corresponding eigenvector or generalized eigenvector of $\mathcal{M}$. Similarly, if $U_j$ is an eigenvector or generalized eigenvector of $A$ corresponding to the eigenvector $\lambda_j$, and if $\lambda_j$ is not an eigenvalue of $M$ then $(\widehat{V}_i^T, U_j^T)$ will be a corresponding eigenvector or generalized eigenvector of $\mathcal{M}$. The assumption that $\lambda_j$ is not an eigenvalue of $M$ is needed to assure that $(M-\lambda_j I)^m$ is a nonsingular matrix, and hence that the linear equation for $\widehat{V}_i$ has a solution. These considerations give:

**Theorem 7.4.5.** *If $M$ and $A$ have no eigenvalues in common, then the solution to*

$$X_{t+1} = MX_t + Y_t$$

*can be written ion the form*

$$X_t = \sum_{i=1}^{k} \alpha_i t^{m_i} \lambda_i^t V_i + \sum_{j=1}^{k_2} \beta_j t^{m_j} \gamma_j^t U_j$$

## 7.5 Exercises

**Ex 7.1.** Let

$$M = \begin{bmatrix} -4 & 9 \\ -4 & 8 \end{bmatrix}.$$

Use the two solutions $x_1(n) = n2^{n-1}$ and $x_2(n) = 2^n$ to find an expression for $M^n$. Compare your result to that found in equation 7.4

**Ex 7.2.** Construct the graph on 20 vertices which has only simple cycles of lengths 6, 15, and 20. Find the least $k$ so that there is a path of length $k$ between every pair of vertices. Construct the corresponding matrix, $M$ , show that $M^k \gg 0$ and that $M^\ell$ is not strictly positive if $\ell < k$.

**Ex 7.3.** Let $A$ be a $10 \times 10$ Leslie matrix in which $f_6$ is the last positive fertility rate. Assume that the initial population is $x_1 = x_2 = x_3 = x_4 = x_5 = x_6 = 0, x_7 = 100, x_8 = 100, x_9 = 90, x_{10} = 10$. Find the asymptotic population vector for $X_{t+1} = AX_t$.

**Ex 7.4.** Let $A = \begin{bmatrix} 0 & 0 & 2 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$. Let $X_0 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$. Calculate the asymptotic population vector for $X_{t+1} = AX_t$. How does this calculation reflect the remarks about nonprimitive matrices?

**Ex 7.5.** Let $g$ be the greatest common divisor of the cycle lengths for a nonnegative strongly connected matrix. Is $M^g$ is primitive? Does $M^g$ decompose into a set of primitive submatrices?

**Ex 7.6.** Given the following graph, determine $g$, the greatest common divisor of the cycle lengths, and find the decomposition of the graph into $g$ disconnected subgraphs.




**Ex 7.7.** Let $X_{t+1} = MX_t$ with $M = \begin{bmatrix} 0 & 0 & 0 \\ 1/3 & 0 & 1 \\ 2/3 & 1 & 0 \end{bmatrix}$. Find the graph for this system. Find the DAG for this graph. Is the system a Markov chain? Is it periodic? If $X_0 = (x_1(0), x_2(0), x_3(0))^T$ find the solution. Find initial conditions which lead to an asymptotic fixed point.

**Ex 7.8.** Let $X_{t+1} = MX_t$ with $M = \begin{bmatrix} 0 & 0 & + & 0 \\ + & 0 & 0 & 0 \\ 0 & + & 0 & + \\ + & 0 & 0 & 0 \end{bmatrix}$. Draw the graph and DAG for this system. Find the period, $g$, of the system and show how $M^g$ decomposes into subsystems.

**Ex 7.9.** Let $X_{t+1} = MX_t$ with $M = \begin{bmatrix} 0 & 2 & 0 \\ 1 & 0 & 0 \\ 1 & 0 & 1 \end{bmatrix}$. Find the graph and DAG for this system. Let $X_0 = (x_1(0), x_2(0), x_3(0))^T$ and find

$$\lim_{t\to\infty} \frac{X_{2t}}{2^t}.$$

Compare this to the predicted asymptotic behavior based on a graph analysis.

**Ex 7.10.** Let $X_{t+1} = MX_t$ where $M = \begin{bmatrix} 0 & 1 & .1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & .9 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix}$. Draw the graph and DAG for this system. Is this a Markov chain? Predict the asymptotic behavior by a graph analysis. Use a computer to follow the evolution of $X_t$ starting from various nonnegative initial conditions. Do the computed results for $X_t$ agree with your predictions?

**Ex 7.11.** A graph is in **Leslie normal form** iff it has $n$ vertices $v_1, v_2, \ldots v_n$ and $\forall i \quad v_i \to v_{i+1(\bmod n)}$ and all other edges have the form $v_j \to v_1$, that is, all other edges go to $v_1$. (Multiple edges are not allowed, but a self-loop $v_1 \to v_1$ is allowed.) Two strongly connected graphs are **g-equivalent** iff the greatest common divisor of the set of cycle lengths is the same for each graph. Two strongly connected graphs are **c-equivalent** iff the set of cycle lengths for the two graphs is the same.

(a) Is every strongly connected graph $c$-equivalent to a graph in Leslie normal form?
(b) Is every strongly connected graph $g$-equivalent to a graph in Leslie normal form?
(c) If every strongly connected graph is equivalent to a graph in Leslie normal form, is the Leslie graph unique? If not, how many Leslie graphs are equivalent to a given strongly connected graph?
(d) Is there a *minimum* Leslie graph equivalent to a given strongly connected graph?
(e) If there is such a minimum Leslie graph, does it have a special form?

**Ex 7.12.** Show that a matrix may be similar to more than one blockwise companion matrix. (Hint: Consider the effect of various vectors on the matrix $\begin{bmatrix}1 & 1\\ 0 & 2\end{bmatrix}$.)

**Ex 7.13.** Show that a $k \times k$ Jordan block is similar to a companion matrix whose companion polynomial is $(x-\lambda)^k$ where $\lambda$ is the eigenvalue of the Jordan block. (Hint: Consider the effect of this companion matrix on the column vector $(\lambda^{k-1}, \lambda^{k-2}, \ldots, \lambda, 1)^T$, and on the vectors formed by taking the derivatives of this vector with respect to $\lambda$ and normalizing so that the constant term in the vector is 1.)

**Ex 7.14.** Show the following: If $\lambda_i$ is an eigenvalue of multiplicity 1 of the matrix $M$, then the solution to

$$X_{t+1} = MX_t$$

can be written as

$$X_t = a\lambda_i^t C + Y_t$$

where $C$ is the column eigenvector for $\lambda_i$, $R$ is the row eigenvector for $\lambda_i$, $a = (R \cdot X_0)/(R \cdot C)$ and $Y_t$ is a vector so that $(R \cdot Y_t) = 0$.

**Ex 7.15.** Let $X_{t+1} = MX_t$ with $M = \begin{bmatrix}3 & -3/2 & -1/2\\ 1 & 0 & 0\\ 3 & -2 & 0\end{bmatrix}$. Find the eigenvalues of the matrix and use them to predict the asymptotic behavior of solutions to the difference equation. Find the solution for the initial value vector $(2,0,0)^T$. Compare this solution to the solution with initial value vector $(1,1,1)^T$. Find the companion form and the Jordan form for the matrix and use them to explain the differences between the two solutions.

**Ex 7.16.** Let $X_{t+1} = HX_t$ where $H = \begin{bmatrix}1 & 1 & 1 & 1\\ 1 & -1 & 1 & -1\\ 1 & 1 & -1 & -1\\ 1 & -1 & -1 & 1\end{bmatrix}$. Show that $H$ has only two eigenvalues but 4 linearly independent eigenvectors. Show that the solution obeys

$$X_t = \begin{cases} 2^t X_0 & \text{if } t \text{ even} \\ 2^{t-1} X_1 & \text{if } t \text{ odd} \end{cases}$$

where $X_1 = HX_0$. Give some initial conditions so that $X_1$ is very different from $X_0$.

**Ex 7.17.** Let $X_{t+1} = MX_t$ with $M = \begin{bmatrix}5 & 3 & 2\\ 3 & 2 & 1\\ 2 & 1 & 1\end{bmatrix}$. Find the eigenvalues and eigenvectors of the matrix $M$ and use them to predict the asymptotic

behavior of solutions to the difference equation. Show that if $X_0 \gg 0$ then $X_t = \Theta(\lambda_0^t)$ where $\lambda_0$ is the unique positive eigenvalue of $M$. More strongly, show that if $X_0 \gg 0$ then $\lim_{t\to\infty}(X_t - \lambda_0^t\, E_1) = 0$ where $ME_1 = \lambda_0 E_1$. Show that for all $X_0$, $\lim_{t\to\infty}(X_t - \lambda_0^t\, E_1) = 0$ where $ME_1 = \lambda_0 E_1$, but for some $X_0$'s one must take $E_1 = 0$.

**Ex 7.18.** Let $X_{t+1} = MX_t$ with $M = \begin{bmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{bmatrix}$. Find conditions under which $X_t = \Theta((2+\sqrt{2})^t)$ and conditions under which $X_t = \Theta((2)^t)$.

**Ex 7.19.** Show that for a strongly connected graph on $n$ vertices exactly one of the following two statements holds:

(a) the graph is primitive and there is a vertex $v$ so that the lengths of all cycles of length $\leq n$, which contain $v$, have gcd $= 1$,

(b) the graph is <u>not</u> primitive and for all vertices $v$ the lengths of all cycles of length $\leq 2n-1$ which contain $v$ have gcd $> 1$.

Use this result to create an algorithm which takes a strongly connected graph as input and decides whether or not the graph is primitive. Show that the time complexity of your algorithm is $O(nE)$ for a graph with $n$ vertices and $E$ edges.

**Ex 7.20.** If $X_{t+1} = MX_t + Y_t$, and $Y_{t+1} = MY_t$, and $M$ has a complete set of eigenvectors $V_1, \ldots, V_k$, show that every solution $X_t$ can be written as

$$X_t \quad = \quad \sum_{i=1}^{k} (\alpha_i\, t \,+\, \beta_i) V_i$$

where the $\alpha_i$'s depend only on the the initial conditions for $Y_t$.

# 8
# Modular Recurrences

In this chapter we consider recurrence relations modulo a fixed positive integer. For any positive integer $m \geq 2$, the output of the operation of reducing an integer modulo $m$ is the remainder after division by $m$ where the remainder lies in the set $\{0, 1, \ldots, m-1\}$. For instance, computing a few terms of the Fibonacci sequence modulo 6 gives

$$1, 1, 2, 3, 5, 2, 1, 3, 4, 1, 5, 0, 5, \ldots, \tag{8.1}$$

where each term is the "mod 6 sum" of the two previous terms. We are interested in answers to the following types of questions:

- Is this sequence periodic or eventually periodic modulo 6?
- What is its period?
- What is the largest period of a sequence which satisfies the Fibonacci recurrence modulo 6?
- How many different sequences satisfy the Fibonacci recurrence modulo 6?

The first question has a fairly quick answer and we give it in the next section. If we have only basic properties of modular arithmetic at our disposal,

answers to the other questions are quite complicated. Instead of using only these basic properties, we give a more sophisticated but accessible point of view which will illuminate the general structure of recurrences modulo $m$ and provide straightforward answers to these questions. We end the chapter with applications of modular recurrences to pseudorandom number generation and to integer factorization.

## 8.1 Periodicity

There's no apparent pattern in the first few terms of the Fibonacci numbers modulo 6 as listed in (8.1). Continuing farther into the sequence,

$$1,1,2,3,5,2,1,3,4,1,5,0,5,5,4,3,1,4,5,3,2,5,1,0,1,1,2,3,\ldots, \tag{8.2}$$

and we see that the initial pair $1,1$ occurs again. Once the initial pair recurs the sequence will repeat because the recurrence is second-order. Also, at least one repetition must occur within the first 37 terms of the sequence because there are only 36 different pairs of elements with entries modulo 6. A rewording this of this is helpful. Let $X$ be the set of all ordered pairs with entries in the set of integers modulo 6, and let $f$ be the Fibonacci function modulo 6 defined on $X$ by

$$f(x,y) = (y, x+y) \pmod 6, \tag{8.3}$$

where this notation means that each coordinate of the point $(y, x+y)$ is reduced modulo 6. Since there are only $6^2$ different pairs of integers modulo 6, any list of 37 consecutive pairs from a recurrence modulo 6 must contain a repeated pair, which means a repetition occurs within the first 37 terms. This argument holds for any second-order recurrence modulo 6 with any pair of initial values.

Putting this in a general context: For a function $f$ defined on a set $X$, let's use $f^{(n)}$ to denote the **$\mathbf{n^{th}}$ iterate of** $f$,

$$f^{(n)} = \overbrace{f \circ f \circ f \circ \cdots \circ f}^{n \text{ times}},$$

and the **orbit** of $x \in X$ under $f$ is the sequence

$$x, f(x), f^{(2)}(x), f^{(3)}(x), f^{(4)}(x) \ldots,$$

the sequence formed by starting with the value $x$ and applying $f$ again and again. If there is some $n > 0$ such that $f^{(n)}(x) = x$, the orbit of $x$ is called a **periodic orbit**, $x$ is called a **periodic point**, and its **period** is the least positive $t$ such that $f^{(t)}(x) = x$. When $t = 1$ holds, $x$ is called a **fixed point of** $f$. From (8.2) we see that $(1,1)$ is a periodic point of the Fibonacci function modulo 6 and its period equals 24.

The orbit of a *periodic point* with period $t$ can be visualized as forming a closed loop which returns to its starting point after $t$ iterations of the function $f$. Sometimes an orbit might get into a loop without returning to its initial value. We will call $x$ an **eventually periodic point** of $f$ if there exists an integer $t$ such that $f^{(n+t)}(x) = f^{(n)}(x)$ holds for all sufficiently large $n$, and the smallest such $t$ is its **period**. (Refer to Figure 8.1.) Notice that under this definition every periodic point is eventually periodic.




FIGURE 8.1. Periodic and Eventually Periodic Orbits.

For example, for the function $f(x) = -x^2 + 2x + 1$ on the set $X = \mathbb{Z}$ the orbit of $x = 0$ is the sequence $0, 1, 2, 1, 2, 1, 2, \cdots$, and $x = 0$ is an eventually periodic point of $f$ which is not periodic, while both $x = 1, 2$ are periodic points of $f$. For each of these three values of $x$ the period is two.

**Theorem 8.1.1.** *If $f$ is a function on a finite set $X$ then every element of $X$ is an eventually periodic point of $f$ and its period is at most the number of elements in $X$. If $f$ is a one-to-one function on any (not necessarily finite) set $X$, then every eventually periodic point of $f$ is periodic. In particular, if $f$ is a one-to-one function on a finite set, then every element of $X$ is a periodic point of $f$.*

*Proof.* If $X$ has $n$ elements, the set $\{x, f(x), \ldots, f^{(n)}(x)\}$ contains $n+1$ elements from $X$ and must contain a duplicate, say $f^{(i)}(x) = f^{(j)}(x)$ for some $0 \le i < j \le n$. Then

$$\begin{aligned} f^{(i+1)}(x) &= f(f^{(i)}(x)) = f(f^{(j)}(x)) = f^{(j+1)}(x)\,, \\ &\vdots \\ f^{(i+k)}(x) &= f^{(j+k)}(x) \text{ for all } k \ge 0\,. \end{aligned}$$

Therefore, for $m \geq i$

$$f^{(m+(j-i))}(x) = f^{(j+(m-i))}(x) = f^{(i+(m-i))}(x) = f^{(m)}(x)\,,$$

and we see $x$ is an eventually periodic point whose period is at most $j-i \leq n$.

Suppose $f$ is one-to-one and the period of $x \in X$ equals $t$. Then there exists a minimal $N_0$ such that $f^{(n+t)}(x) = f^{(n)}(x)$ for all $n \geq N_0$. To prove $x$ is periodic, we prove $N_0 = 0$ holds. If $N$ were non-zero, then

$$f(f^{(N-1)}(x)) = f^{(N)}(x) = f^{(N+t)}(x) = f(f^{(N+t-1)}(x))$$

and the fact that $f$ is one-to-one gives $f^{(N-1)}(x) = f^{(N-1+t)}(x)$, contradicting the minimality of $N$. Therefore, $N = 0$, that is, $x$ is a periodic point of $f$. □

For any modulus $m \geq 2$, the Fibonacci function modulo $m$ is

$$F(x,y) = (y, x+y) \pmod{m}\,,$$

a one-to-one function on the ordered pairs of integers modulo $m$ The previous result gives

**Corollary 8.1.2.** *For any $m \geq 2$, the Fibonacci function modulo $m$ has a periodic orbit for every pair of initial values.*

Let's compute the period of the Fibonacci sequence modulo $m$ for four additional values:

$$\begin{aligned}
m = 2: &\quad 1,1,0,1,1,\ldots \text{ whose period is } 3;\\
m = 3: &\quad 1,1,2,0,2,2,1,0,1,1,\ldots \text{ whose period is } 8;\\
m = 4: &\quad 1,1,2,3,1,0,1,1,\ldots \text{ whose period is } 6;\\
m = 12: &\quad 1,1,2,3,5,8,1,9,10,7,5,0,5,5,10,3,1,4,5,\\
&\quad 9,2,11,1,0,1,1,\ldots \text{ whose period is } 24.
\end{aligned}$$

We have already computed the period modulo 6 (refer to (8.2)) to be 24, which is the product of the periods modulo 2 and modulo 3. Because of this, you might conjecture this holds in general but the period modulo 12 is 24 which does not equal $8 \cdot 6$. After some deliberation you can see that 24 does equal $\operatorname{lcm}(8,6)$, the least common multiple of 8 and 6. The significance of this occurrence of the least common multiple is explained in the next theorem.

**Theorem 8.1.3.** *Let $X = \mathbb{Z}^k$ for some $k$ and let $f$ be a function defined on $X$. For $x \in X$ and positive integers $m_1, m_2$, let $t_i$ be the period of $x$ under $f$ modulo $m_i$ for each of $i = 1, 2$. Then the period of $x$ under $f$ modulo $lcm(m_1, m_2)$ equals $lcm(t_1, t_2)$.*

*Proof.* Let $m = \text{lcm}(m_1, m_2)$ and $L = \text{lcm}(t_1, t_2)$. If $t$ is the period of $x$ under $f$ modulo $m$, there exists $N$ such that for all $n \geq N$,

$$f^{(n+t)}(x_0) = f^{(n)}(x_0) \pmod{m},$$

and since each of $m_1$, $m_2$ divides $m$ then $f^{(n+t)}(x_0) = f^{(n)}(x_0) \pmod{m_i}$ for all $n \geq N$. Therefore, $t$ must be a multiple of each $t_i$, and $L$ divides $t$.

On the other hand, by definition of the period for each of $i = 1, 2$ there exists $N_i$ such that $f^{(n+t_i)}(x_0) = f^{(n)}(x_0) \pmod{m_i}$ for all $n \geq N_i$. Using $N = \max\{N_1, N_2\}$ and the fact that $L$ is a multiple of both $t_1$ and $t_2$, for all $n \geq N$

$$f^{(n+L)}(x_0) = f^{(n)}(x_0) \pmod{m_i}.$$

This means that the difference $f^{(n+L)}(x_0) - f^{(n)}(x_0)$ is divisible by both $m_i$ and so also by their least common multiple $m$, giving

$$f^{(n+L)}(x_0) = f^{(n)}(x_0) \pmod{m},$$

and $L$ has been shown to be a multiple of $t$. Since we've already shown that $L$ divides $t$ (and they're both positive), then $L = t$. □

But how do we find the period for a general modulus $m$? If we happened to know two *relatively prime* integers $m_1, m_2$ (that is, the $\gcd(m_1, m_2) = 1$) for which we know (or can easily find) the periods modulo each of $m_1$ and $m_2$ and such that $m = m_1 m_2$, the theorem could be applied to find the period modulo $m$. That's exactly what we noted for $m = 12$. The following result follows directly from this observation.

**Corollary 8.1.4.** *If $m = p_1^{\alpha_1} \cdots p_s^{\alpha_s}$ is the prime factorization of $m$ and $t_i$ is the period of $x$ under $f$ modulo $m_i = p_i^{\alpha_i}$ then the period of $x$ under $f$ modulo $m$ is $lcm(t_1, \cdots, t_s)$.*

### *8.1.1 Periodicity of linear modular recurrences*

We've been working with functions defined on $\mathbb{Z}^k$ which are then reduced modulo $m$ for various values of $m$ but we could have equally well considered $f$ to be defined on the set $X = \mathbb{Z}_m^k$, where $\mathbb{Z}_m$ means the set $\{0, 1, \ldots, m-1\}$ under the operations of addition and multiplication modulo $m$.

As we saw with the Fibonacci recurrence, a $k^{\text{th}}$-order linear recurrence

$$s_{j+k} \equiv c_1 s_{j+k-1} + c_2 s_{j+k-2} + \ldots + c_k s_j + c_{k+1} \pmod{m} \tag{8.4}$$

defines the function

$$S(x_1, \ldots, x_k) = (x_2, \ldots, x_k, c_{k+1} + c_k x_1 + \ldots + c_1 x_k) \tag{8.5}$$

on $\mathbb{Z}_m^k$, where all operations are performed modulo $m$. (This is similar to the shift operator from Chapter 3.) Noticing that

$$S(x_1, \ldots, x_k) = S(y_1, \ldots, y_k) \iff x_2 = y_2, \ldots, x_k = y_k, c_k x_1 = c_k y_1$$

we have

**Lemma 8.1.5.** *$S$ is a one-to-one function on $\mathbb{Z}_m^k$ if and only if $c_k$ has a multiplicative inverse modulo $m$.*

Orbits of $S$ correspond to choices of initial values $s_0, \ldots, s_{k-1}$ in the recurrence (8.4). Since $S$ is a function defined on a finite set with $m^k$ elements, from this lemma and Theorem 8.1.1 we obtain the following result.

**Theorem 8.1.6.** *Every choice of $(s_0, \ldots, s_{k-1})$ is an eventually periodic point of $S$. Moreover, if $c_k$ has a multiplicative inverse modulo $m$ then every orbit under $S$ is periodic.*

Because every element of $\mathbb{Z}_m$ has an additive inverse, we may refer to an element with a multiplicative inverse as an **invertible element** without confusion as to which operation we mean.

For the Fibonacci sequence, $c_k = 1$ which is invertible for every modulus. This means that (regardless of the initial pair) every modular sequence generated by the Fibonacci recurrence is periodic. On the other hand, the orbit of $(1,5)$ under the recurrence $s_{j+2} = s_{j+1} + 3s_j \pmod{18}$ is

$$1\ ,\ 5\ ,\ 8\ ,\ 5\ ,\ 11\ ,\ 8\ ,\ 5\ ,\ 11, \ldots\ ,$$

an eventually periodic sequence which is not periodic. Note also that 3 is not invertible modulo 18 and $S(1,5) = 8 = S(5,11)$ shows that $S$ is not one-to-one.

The periodicity of modular recurrences therefore depends only on the algebraic property of invertibility of $c_k$. We next describe an efficient procedure for determining whether or not an element is invertible, which therefore is a method for deciding when a specific modular linear recurrence is periodic. The procedure is the famous **Euclidean Algorithm** which can be found in the seventh volume of Euclid's Elements. (A translation of this can be found at `aleph0.clarku.edu/~djoyce/java/elements/bookVII/propVII2.html`.) We've already used this algorithm in Chapter 6 (and for polynomials in Chapter 4) where we showed it computes the gcd for $n$-bit numbers in $O(n)$ operations. Because we're interested in using the algorithm to compute more than the gcd, it's helpful to review it here. Recall that the **greatest common divisor** of two non-zero integers $a, m$ [denoted by $\gcd(a,m)$] is a positive integer which divides both $a$ and $m$ and is defined by the property that $\gcd(a,m)$ is divisible by all divisors of both $a$ and $m$.

**Theorem 8.1.7 (Euclidean Algorithm).** *Let $m$ be any positive integer and $a$ be any integer. Then the sequence of divisions*

$$m = q_1 a + r_1\ ;\ a = q_2 r_1 + r_2\ ;\ r_1 = q_3 r_2 + r_3\ ; \ldots$$

*(where $0 \le r_1 < a$ and $0 \le r_{i+1} < r_i$ for all $i > 1$) ends after finitely many steps and the last non-zero remainder $r_K$ is $gcd(a,m)$.*

*Proof.* Since the sequence of remainders $< r_i >$ is a strictly decreasing sequence of nonnegative integers, it must be a finite sequence and its last

remainder is zero:

$$m = q_1 a + r_1\,;\; a = q_2 r_1 + r_2\,;\ldots;\; r_{K-2} = q_K r_{K-1} + r_K\,;\; r_{K-1} = q_{K+1} r_K\,.$$

From the definition of gcd, $\gcd(a,m) = \gcd(m - ka, a)$ holds for all integers $k$, and we successively obtain

$$\begin{aligned}\gcd(a,m) &= \gcd(m - q_1 a, a) = \gcd(r_1, a) = \gcd(r_1, a - q_2 r_1) = \ldots \\ &= \gcd(r_{K-1}, r_{K-2} - q_2 r_{K-1}) = \gcd(r_{K-1}, r_K) = r_K\,,\end{aligned}$$

and the last non-zero remainder is $\gcd(a,m)$. □

For instance, to compute $\gcd(437, 12)$:

$$437 = 36 \cdot 12 + 5\,,\quad 12 = 2 \cdot 5 + 2\,,\quad 5 = 2 \cdot 2 + 1\,,\quad 2 = 2 \cdot 1 + 0\,,$$

and $\gcd(437, 12) = 1$ is obtained. Whenever $\gcd(a,m) = 1$ holds, the steps of the Euclidean Algorithm can be reversed to obtain $a^{-1} \pmod m$. For instance, in this example successively solving backwards for each remainder gives

$$\begin{aligned}1 =& 5 - 2 \cdot 2 = 5 - 2 \cdot (12 - 2 \cdot 5) = 5 \cdot 5 - 2 \cdot 12 \\ =& 5 \cdot (437 - 36 \cdot 12) - 2 \cdot 12 = 5 \cdot 437 - 182 \cdot 12 \\ =& 5 \cdot 437 - 182 \cdot 12 \equiv -182 \cdot 12 \equiv 255 \cdot 12 \pmod{437}\,,\end{aligned}$$

and we see that 255 is the inverse of 12 modulo 437. This method works in general and is the basis for our proof of the following result.

**Theorem 8.1.8.** *Let $m$ be any positive integer and $a$ be any integer. Then $a$ is invertible modulo $m$ if and only if $\gcd(a,m) = 1$.*

*Proof.* Any divisor of both $a$, $m$ must divide $ax + my$ for any integers $x, y$. If $a$ is invertible there exists $x$ such that $ax \equiv 1 \pmod m$ which means that $ax + my = 1$ for some $y$, and $\gcd(a,m) = 1$ holds. On the other hand, when $\gcd(a,m) = 1$ we can "solve the Euclidean Algorithm backwards" to obtain $x, y$ such that $1 = ax + my$, which gives $ax \equiv 1 \pmod m$. □

Combining these results, gives an answer the first question at the beginning of this chapter for general moduli.

**Theorem 8.1.9.** *If $S$ is the function associated with the linear recurrence (8.4), every orbit of $S$ is periodic if and only if $\gcd(c_k, m) = 1$. In particular, when the modulus $m$ is prime every orbit of every linear recurrence modulo $m$ is periodic provided $c_k \neq 0 \pmod m$.*

## 8.2 Fast Modular Computations

In this section we describe Montgomery multiplication, a quick way to perform all modular operations, which was first suggested by Peter Montgomery [99] in 1985. We include a description of it here because of its helpfulness in computing the terms of a linear recurrence modulo $m$ since in general $k$ multiplications modulo $m$ are required to compute

$$s_n \equiv c_1 s_{n-1} + c_2 s_{n-2} + \ldots + c_k s_{n-k} + c_{k+1} \pmod{m}$$

from $s_{n-1}, s_{n-2}, \ldots, s_{n-k}$. The technique also is fundamental for implementations of fast modular exponentiation as used in many cryptosystems, for example RSA [117]. The technique relies on the fact that there are moduli for which arithmetic is quick, for instance a computer's machine word size could be used. For any such modulus $R$ and any $m$ with $\gcd(m, R) = 1$, Montgomery multiplication translates operations modulo $m$ to the faster operations modulo $R$. Our discussion is restricted to $m < R$ and Montgomery's original paper should be consulted for the important case when $m > R$. Since we are assuming that $\gcd(m, R) = 1$, $R$ is an invertible element of $\mathbb{Z}_m$ which means that $RS \equiv 1 \pmod{m}$ for some $1 \leq S < m$ and there exists an integer $1 \leq m' < R$ such that $RS - mm' = 1$.

Normally when $26 \pmod{25}$ is written we think of 1 but $26 \pmod{25}$ is of course an infinite set of integers, all those integers which are congruent to 26 modulo 25. In what follows we use the notation $t \ mod \ m$ (as contrasted with $t \pmod{m}$) to denote the least nonnegative integer congruent to $t$ modulo $m$. For example, $27 \ mod \ 5$ equals 2. Returning to $R, S$ as defined above, we see that the correspondence $\psi : t \mapsto tS \ mod \ m$ is a bijection on $\mathbb{Z}_m$ with inverse $\phi : t \mapsto tR \ mod \ m$, and this allows us to encode any $t \ mod \ m$ as $\psi(t)$ for the purposes of computation, because $\psi(t)$ can be uniquely decoded using the mapping $\phi(\psi(t)) = t$. The advantage of this encoding is that the multiplication $\psi(ab)$ for any $0 < a, b < m$ can be done quickly using operations modulo $R$, which has been chosen to be a modulus for which operations can be done very quickly. The following algorithm calculates $\psi(ab)$ for any $0 < a, b < m$.

```
Procedure MULT(a, b)
n := (ab mod R)m' mod R
t := (ab + nm)/R
WHEN t < m, return t;
     ELSE return t − m.
```

We justify $\text{MULT}(a, b) = \psi(ab)$ before giving an example of its use in modular calculations. First remember division by $R$ can be done quickly,

which means that reduction modulo $R$ is also quick. To show

$$\text{MULT}(a,b) = \psi(ab) = ab \ mod \ m\,,$$

we show that

$$t \text{ is an integer} \quad \text{and} \quad t \equiv abS \pmod{m} \quad \text{and} \quad 0 \le t < 2m\,.$$

¿From the algorithm, $n = abm' + kR$ for some integer $k$ and so

$$ab + nm = ab(1 + mm' + kRm) = ab(RS + kRm)\,,$$

which is divisible by $R$ and with this we know $t$ is an integer. Also $tR \equiv ab \pmod{m}$ and so

$$abS \equiv tRS \equiv t \pmod{m}$$

and from $0 < a, b < m$ and $0 \le n < R$,

$$0 \le tR = ab + nm < m^2 + mR < 2mR\,.$$

Let's use the algorithm to calculate the second element of the solution to the initial value problem

$$s_0 = 12\,,\ s_1 = 10\,,\ s_n = 13s_{n-1} + 9s_{n-2} \pmod{27}\,.$$

We use $R = 32$, and obtain $S = 11$ and $m' = 13$, to compute $T = \psi(s_2)$ and then find $\phi(T) = s_2$. First we calculate $\psi(13{\cdot}10) = \text{MULT}(13, 10)$ using the notation of the algorithm. Tnen $130 \ mod \ 32 = 130 - 128 = 2$ ,

$$n = 2 * 13 = 26 \ mod \ 32 \text{ and } t = (130 + 26 * 27)/32 = 26 < m$$

gives $\psi(13 \cdot 10) = 26$. Similarly, for $\psi(9 \cdot 12) = \text{MULT}(9, 12)$,

$108 \ mod \ 32 \equiv 108{-}128 \pmod{32} = 12\,,\ n \equiv 12{*}13 \pmod{32} = 156{-}128 = 28 \ mod \ 32$

and

$$t = (108 + 28 * 27)/32 = 27 < m\,,$$

which gives $\psi(9 \cdot 12) = 27 - 27 = 0$. Therefore,

$$\psi(s_2) \equiv \psi(13 \cdot 10) + \psi(9 \cdot 12) = 26\,.$$

How do we find $s_2$ from this? Notice that in general

$$\psi(\psi(t) * R^2) = t * R^2 * S^2 \ mod \ m = t * (RS)^2 \ mod \ m = t \ mod \ m\,,$$

that is, $t = \text{MULT}(\psi(t), R^2 \ mod \ m)$ where $R^2 \ mod \ m$ is a one-time calculation. Applying this result here, $R^2 \equiv (32)^2 = 25 \ mod \ m$ and for $s_2 = \text{MULT}(26, 25)$,

$$\begin{aligned} n &= (26 * 25 \ mod \ 32)13 \ mod \ 32 = (650 - 512 \ mod \ 32)13 \ mod \ 32 \\ &= (138 - 128 \ mod \ 32)13 \ mod \ 32 = 130 \ mod \ 32 \\ &= 2 \ mod \ 32 \end{aligned}$$

which gives $s_2 = (650 + 2 * 27)/32 = 22$.

## 8.3 Periodicity of Linear Recurrences in Finite Fields

In this section we begin the study of finite fields, which are generalizations of the set of integers modulo a prime and provide a more sophisticated context for investigating modular recurrences. Évariste Galois [57] was the first to use some of the properties of finite fields and the first systematic theory was written by Leonard E. Dickson in [45]. Because they're *finite*, recurrences in finite fields satisfy the periodicity results proved in Section 8.1.

Let $\mathbb{F}$ be a finite set with two operations $+$ and $*$. Then $\mathbb{F}$ **is a finite field** (under $+$ and $*$) if all of the following hold:
(i) *The set* $\mathbb{F}$ *is an* **abelian group** *under* $+$, which means that $+$ is an associative and commutative operation on $\mathbb{F}$ and has a special **identity element** denoted by 0 such that $a + 0 = a$ holds for all $a \in \mathbb{F}$ and each element $a$ has an (additive) inverse $b \in \mathbb{F}$ satisfying $a + b = 0$.
(ii) *The set of non-zero elements of* $\mathbb{F}$ *forms an abelian group under* $*$, which as above means that $*$ is an associative and commutative operation and has the special *identity element* denoted by 1 such that $a * 1 = a$ holds for all $a \in \mathbb{F}$ and each non-zero element $a$ has a (multiplicative) inverse $b \in \mathbb{F}$ satisfying $a * b = 1$.
(iii) The operations of $+$ and $*$ are connected by the distributive law, where $a_1 * (a_2 + a_3) = a_1 * a_2 + a_1 * a_3$ for all $a_1, a_2, a_3 \in \mathbb{F}$.

For example, $S = \{0, 1, 2, 3, 4\}$ is a field under the operations of addition and multiplication modulo 5, where the additive and multiplicative identities are respectively 0 and 1; 0 is its own additive inverse and $1, 2, 3, 4$ have additive inverses $4, 3, 2, 1$ and multiplicative inverses $1, 3, 2, 4$. The associative, commutative and distributive properties are inherited from the set of integers. Although $S = \{0, 1, 2, 3\}$ is an abelian group under the operation of addition modulo 4, the fact that $\gcd(2, 4) = 2$ implies 2 does not have a multiplicative inverse under multiplication modulo 4, which means that $S$ is not a finite field under these operations.

**Theorem 8.3.1.** *Let* $\mathbb{F}$ *be the set of integers modulo* $m$*. Then* $\mathbb{F}$ *is a finite field under the operations of addition modulo* $m$ *and multiplication modulo* $m$ *if and only if* $m$ *is prime.*

*Proof.* As in the two examples above, we observe that the associative, commutative and distributive properties are all inherited from the set of integers. Also, 0 is the additive identity and $\mathbb{F}$ is an abelian group under addition modulo $m$. By Theorem 8.1.8, $a$ is invertible modulo $m$ if and only if $\gcd(a, m) = 1$. Ensuring that $\gcd(a, m) = 1$ for all $1 \leq a < m$ is equivalent to requiring that $m$ be prime. □

The number of elements in a finite field can be composite, but must be a prime power. (Refer to Exercise 8.5.)

Let $\mathbb{F}$ be the set of polynomials $\{0, 1, x, x+1\}$. If we define addition on $\mathbb{F}$ to be the usual polynomial addition followed by reducing the coefficients modulo 2, we see that 0 is the additive identity and $\mathbb{F}$ is an abelian group under addition. Multiplication is defined as in Table 8.1.

TABLE 8.1. The multiplication table for the field with four elements

| * | 0 | 1 | x | x+1 |
|---|---|---|---|---|
| 0 | 0 | 0 | 0 | 0 |
| 1 | 0 | 1 | x | x+1 |
| x | 0 | x | x+1 | 1 |
| x+1 | 0 | x+1 | 1 | x |

Then 1 is the multiplicative identity and each of non-zero elements $1, x, x+1$ has a multiplicative inverse (respectively $1, x+1, x$). Therefore, $\mathbb{F}$ is a field under these operations.

It's helpful to examine this example further. In earlier chapters the notation $\mathbb{R}[x]$ (or $\mathbb{C}[x]$) meant the set of polynomials with coefficients from $\mathbb{R}$ (or $\mathbb{C}$). Likewise, here $\mathbb{F}[x]$ is the set of polynomials with coefficients from the set $\mathbb{F}$. For instance, $\mathbb{Z}_2[x]$ is the set of all polynomials whose coefficients are either 0 or 1. We're usually interested not only in the *set* of polynomials, but in the algebraic properties of $\mathbb{F}[x]$. Here the polynomial operations of addition and multiplication are defined similarly to the operations in $\mathbb{R}[x]$ with coefficients being calculated by the operations in $\mathbb{F}$. For example, you can check that when $f(x) = x^2 + x$ and $g(x) = x^3 + x + 1$ are considered as elements of $\mathbb{Z}_2[x]$,

$$f(x) + g(x) = x^3 + x^2 + 1 \quad \text{and} \quad f(x)g(x) = x^5 + x^4 + x^3 + x\,.$$

We are now in a position to describe the above field with four elements in another way. Let $S$ be the subset of all constant and linear polynomials in $\mathbb{Z}_2[x]$. Since $S$ is an abelian group under the polynomial addition of $\mathbb{Z}_2[x]$, it is a suitable addition for $S$. What about multiplication? The set $S$ is not closed under the usual polynomial multiplication reduced modulo 2, because for instance, $(x+1)^2 = x^2 + 1$ is not an element of $S$. But a slight modification of the usual multiplication using the polynomial $f(x) = x^2 + x + 1$ (which is not in $S$) works. Namely, the product of two elements in the above field $\mathbb{F}$ is defined by first obtaining the product $p(x)$ of the two polynomials as elements of $\mathbb{Z}_2[x]$ and then "reducing $p(x)$ modulo $f(x)$", in other words, finding the remainder when $p(x)$ is divided by $f(x)$. Since the polynomial $f(x)$ is quadratic, the remainder will always be in the set $S$. For example, $p(x) = x(x+1) = x^2 + x$ and $p(x) = f(x) + 1$ gives the product $x * (x+1) = 1$ in $S$.

This procedure can be generalized to obtain a finite field with $p^m$ elements for any prime $p$ and positive integer $m$. The construction relies on

the fact that for every $p$ and $m$ there exists an **irreducible polynomial** in $\mathbb{Z}_p[x]$ with $\deg(f) = m$, where here irreducible means that $f(x)$ cannot be factored into polynomials of smaller degree in $\mathbb{Z}_p[x]$. A proof that this construction works and more information about finite fields can be found in [86, p.91ff]. In particular, it can be shown that for each choice of prime $p$ and exponent $m$ there is essentially only one finite field with $q = p^m$ elements. These fields are frequently denoted by $GF(q)$ and are the finite **Galois fields**, named in honor of É. Galois. (Refer to Exercises 8.7-8.11.) In what follows we'll drop the $*$ symbol for multiplication, and instead use juxtaposition of elements to denote multiplication. Our proof of Theorem 8.1.9 generalizes to finite fields:

**Theorem 8.3.2.** *If $\mathbb{F}$ is a finite field then every linear recurrence in $\mathbb{F}$ is periodic.*

## 8.4 The Period of First-Order Linear Recurrences

In what follows we allow $R$ to be either the set of integers modulo $m$ or a finite field and consider the periods of first-order linear recurrences in $R$. We first note that when $a = 1$, $s_n = s_0 + nb$ and the period of $S(x) = x + b$ is therefore the least integer $t$ such that $tb = 0$.

**Lemma 8.4.1.** *Let $a$ be an element of $R$ such that $a-1$ has a multiplicative inverse $c$ in $R$. If $< s_j >$ is any sequence generated by the recurrence $s_{j+1} = as_j + b$ in $R$, then for all integers $m, j$,*

$$s_{m+j} = a^j s_m + bc(a^j - 1). \tag{8.6}$$

*Proof.* Fix $m$. We prove (8.6) by induction on $j \geq 1$. Since $c(a-1) = 1$ holds in $R$,

$$s_{m+1} = as_m + b = as_m + bc(a-1)\,.$$

Assuming (8.6) holds for all $1 \leq j < J$,

$$\begin{aligned} s_{m+J} &= as_{m+J-1} + b = a(a^{J-1}s_m + bc(a^{J-1} - 1)) + bc(a-1) \\ &= a^J s_m + bc(a^J - a + a - 1) = a^J s_m + bc(a^J - 1), \end{aligned}$$

completing the proof. □

**Theorem 8.4.2.** *Let $S$ be the function defined on $R$ by $S(x) = ax + b$, where $b$ is arbitrary and $a-1$ is any element of $R$ which has a multiplicative inverse $c$ in $R$. Then $s_0 = -bc$ is the only fixed point of $S$, and every $s_0 \in R$ is an eventually periodic point of $S$ whose period equals the least integer $t$ such that $(a^t - 1)(S^{(m+1)}(s_0) - S^{(m)}(s_0)) = 0$ holds for all sufficiently large $m$.*

*Proof.* We see that $s_0$ is a fixed point iff $as_0+b = s_0$, which can be uniquely solved for $s_0 = -bc$. From Theorem 8.1.6, every orbit of $S$ is eventually periodic, and if $t$ is any multiple of the period of $s_0$ then $s_{m+t} = s_m$ for sufficiently large $m$. By (8.6) this becomes

$$a^t s_m + bc(a^t - 1) = s_m \; ; \; (a^t - 1)(s_m + bc) = 0.$$

Multiplying this last congruence by $a - 1$ gives

$$0 = (a^t - 1)((a-1)s_m + b) = (a^t - 1)(s_{m+1} - s_m).$$

Because $a - 1$ is invertible in $R$, each of these steps is reversible and we obtain $(a^t - 1)(s_m + bc) = 0$ iff $s_{m+t} = s_m$, and such $t$ is a multiple of the period of $s_0$ under $S$. □

Let's consider orbits of $S(x) = 12x + 4$ on $R = \mathbb{Z}_{21}$. Since $11^{-1} = 2$, $-bc = 13$ is the only fixed point of $S$. For instance, the orbits of 1 and 2 are

$$1, 16, 7, 4, 10, 19, 1, \ldots \quad \text{and} \quad 2, 7, 4, 10, 19, 1, 16, 7, \ldots,$$

and 1 is a periodic point while 2 is *eventually* periodic. The periods are the same, but is that only because 7 is a common term of both orbits? In Exercise 8.14 you're asked to find the periods of other orbits under this map. When $R$ is restricted to be a finite field, the period depends only on the coefficient $a$ in $S(x) = ax + b$.

**Theorem 8.4.3.** *Consider the iteration of $S(x) = ax + b$ in the finite field $\mathbb{F}$. When $a = 1$, the period of any sequence is the characteristic of $\mathbb{F}$. When $a \neq 1$, let $c$ be the multiplicative inverse of $a - 1$. Then $x = -bc$ is the only fixed point of $S$ and for all other $x \neq -bc$ the period equals the period of 1 under the homogeneous linear recurrence $S_0(x) = ax$. This number is called the* **order of** *$a$ in $\mathbb{F}$ and will be denoted by $ord(a)$.*

*Proof.* From our remark before Lemma 8.4.1, the period of $S(x) = x + b$ is the least integer such that $pb = 0$, which by definition is the characteristic of $\mathbb{F}$. (Refer to Exercise 8.4.) For any $S(x) = ax + b$ with $a \neq 1$, the orbit of any $s_0 \neq -bc$ is periodic with period $t \geq 2$, which implies all $s_{m+1} \neq s_m$, and $s_{m+1} - s_m$ is an invertible element of $\mathbb{F}$. Therefore, the condition in the last theorem becomes $a^t = 1$ and the period is $t = ord(a)$. □

Because $ord(a)$ is the period of the sequence $< a^j >$, we have

$$a^i = a^j \iff i \equiv j \pmod{ord(a)}. \tag{8.7}$$

The following algebraic result is a computational aid for computing $ord(a)$. Its statement appeared in a letter written by Fermat in 1640, and was later proved by Leibniz. It's usually called Fermat's *Little* Theorem to differentiate it from Fermat's *Last* Theorem which was finally proved

in 1994 by Andrew Wiles with the assistance of Richard Taylor [147, 139]. Alf van der Poorten [143] has published a fairly accessible account of the proof, which was the culmination of the work of many mathematicians.

**Theorem 8.4.4 (Fermat's Little Theorem).** *If $\mathbb{F}$ is a finite field with $q$ elements, then every non-zero $a \in \mathbb{F}$ satisfies $a^{q-1} = 1$, and so $ord(a)$ divides $q-1$.*

*Proof.* Let $a \in F$ be non-zero, and let $a_1, \ldots, a_{q-1}$ be any ordering of all non-zero elements of $\mathbb{F}$. If $a^{-1}$ is the multiplicative inverse of $a$, we note that for any $i, j$,

$$aa_i = aa_j \iff a^{-1}aa_i = a^{-1}aa_j \iff a_i = a_j\,,$$

and with this we see that the sets $\{aa_1, \ldots, aa_{q-1}\}$ and $\{a_1, \ldots, a_{q-1}\}$ are equal. Because multiplication is commutative, the product of the elements of the first set equals the product of the elements in the second set,

$$a_1 \ldots a_{q-1} = (aa_1)\ldots(aa_{q-1}) = a^{q-1}(a_1 \ldots a_{q-1}).$$

Since the product $a_1 \ldots a_{q-1}$ is a non-zero element of the field $\mathbb{F}$, and multiplying by its inverse yields $a^{q-1} = 1$. □

**Corollary 8.4.5.** *The period of any first-order linear recurrence in a finite field with $q$ elements divides $q-1$.*

We will now use this result to determine the periods of the orbits of $S(x) = 215x+3$ in $Z_m$ for the prime $m = 12323$ without actually calculating any orbit. Since $m$ is a prime, $\mathbb{F} = \mathbb{Z}_m$ is a finite field and all orbits are periodic. From Exercise 8.3(b) we know that 5816 is the multiplicative inverse of $a - 1 = 214$ in $\mathbb{F}$. Therefore $s_0 = -3 * 5816 = 7198$ is the only fixed point of $S$ and the period of any $s_0 \neq 7198$ equals $ord(215)$ in $\mathbb{F}$. It remains to find $t = ord(215)$. Using Fermat's Little Theorem, $t$ divides $m - 1 = 12322 = (2)(61)(101)$, and so there are a number of possibilities for $t$. Calculating $a^k$ for the divisors $k$ of 12322 can be done by **fast exponentiation**, the same process which we have already used to find powers of matrices. For instance, $101 = 2^6 + 2^5 + 2^2 + 1$ gives

$$a^{101} = a^{2^6}a^{2^5}a^{2^2}a = 4797 * 1822 * 3809 * 215 = 1.$$

Since 101 is prime this means $ord(215) = 101$, and with this we know $S$ has one fixed point and 122 orbits of period 101.

We close this section with one more algebraic result. The polynomial equation $x^2 - 1 = 0$ is quadratic, but has 4 solutions in $\mathbb{Z}_8$. When the coefficient set is a *field*, every polynomial equation

$$x^r + a_{r-1}x^{r-1} + \ldots + a_1x + a_0 = 0 \quad \text{with } a_i \in \mathbb{F}$$

has at most $r$ solutions in $\mathbb{F}$. (Refer to Exercise 8.22.) This simple fact can have some surprising consequences. For instance, suppose $\mathbb{F}$ is a field with $q = p^t$ elements. Since the order of $s \in \mathbb{F}$ divides $p-1$ iff $s$ is a root of $x^{p-1} - 1 = 0$, there are at most $p-1$ elements of $\mathbb{F}$ whose order divides $p-1$ and Fermat's Little Theorem gives the following result which will be useful later.

**Theorem 8.4.6.** *Let $\mathbb{F}$ be a finite field with $q = p^t$ elements, and $s$ be a non-zero element of $\mathbb{F}$. Then the order of $s \in \mathbb{F}$ divides $p-1$ if and only if $s \in \mathbb{Z}_p$.*

### *8.4.1 First-order recurrences with maximal period*

In this section we prove that any finite field with $q$ elements has an element whose order is $q-1$, the largest possible period. These elements are called **primitive elements** of $\mathbb{F}$. When $a \in \mathbb{F}$ is a primitive element it is necessary that $a \neq 0, 1$ which means that both $a$ and $a-1$ are invertible elements of $\mathbb{F}$, and for any linear function of the form $S(x) = ax+b$ (for some $b \in \mathbb{F}$) the period of any non-fixed point is $q-1$. When the coefficient $a$ is a primitive element there are only two orbits: the fixed point and everything else!

**Lemma 8.4.7.** *Let $a, b$ be non-zero elements of $\mathbb{F}$ with $ord(a) = r$ and $ord(b) = s$. If $\gcd(r,s) = 1$ then $ord(ab) = rs$.*

*Proof.* Let $n = ord(ab)$. Then $(ab)^n = 1$. Also,

$$1 = ((ab)^n)^r = (ab)^{rn} = (a^r)^n (b)^{rn} = (b)^{rn},$$

and $rn$ is a multiple of $ord(b) = s$. Since $\gcd(r,s) = 1$ this implies that $s$ divides $n$. Interchanging the roles of $a$ and $b$ we similarly obtain that $r$ also divides $n$, and so $n$ is a common multiple of $r$ and $s$. Again using the fact that $\gcd(r,s) = 1$, this means that $rs$ divides $n$. We complete the proof by showing that $n$ divides $rs$. (Since $rs$ and $n$ are both positive, this will give $n = rs$.) For this,

$$(ab)^{rs} = (a^r)^s (a^s)^r = 1,$$

implying $rs$ is a multiple of $ord(ab) = n$. Therefore, $n = rs$, as required. □

**Theorem 8.4.8 (Primitive Element Theorem).** *Every finite field with $q$ elements has at least one element whose order equals $q-1$.*

*Proof.* Let $N$ be the maximum element in the finite set $\{ord(b) : b \in \mathbb{F}\}$, and let $d$ be some element of $\mathbb{F}$ with $ord(d) = N$. Fermat's Little Theorem implies that $N \leq q-1$. If we can prove that the order of every element divides $N$, then each of the $q-1$ non-zero elements of $\mathbb{F}$ would satisfy $x^N - 1 = 0$, and $N \geq q-1$ also would hold.

By way of contradiction, we assume there exists an element $c \in F$ whose order does not divide $N$. Therefore, there exists a prime $p$ such

that $p^\gamma$ divides $ord(c)$ and $p^\gamma$ does not divide $N$. Let $ord(c) = p^\gamma N_1$ where $\gcd(N_1, p) = 1$ and let $\beta < \gamma$ such $N = p^\beta N_2$ with $\gcd(N_2, p) = 1$. For $a = c^{N_1}$ and $b = d^{p^\beta}$ we have $ord(a) = p^\gamma$ and $ord(b) = N_2$. The preceding lemma therefore implies that $ord(ab) = p^\gamma N_2 > p^\beta N_2 = N$, contrary to the maximality of $N$, and we see that the order of every element of $\mathbb{F}$ does divide $N$. □

In Sections 69-75 of his famous *Disquisitiones Arithmeticae* [61], C. F. Gauss considers the question of finding primitive elements for a prime modulus, and outlines the method we've used in the last example. Gauss also says "Euler admits that it is extremely difficult to pick out these numbers (primitive elements) and that their nature is one of the deepest mysteries of numbers." Finding a primitive element is still thought to be a hard problem and is related to the discrete log problem, which is the basis for the assumed security of several cryptosystems.

In 1927 Emil Artin was the first to formulate the conjecture that every non-square positive integer is a primitive element for infinitely many prime moduli. Some specific examples had already been constructed in the years between Gauss' work and Artin's formulation of this conjecture. For instance, it's known that 2 is a primitive element for all primes $p$ of the form $p = 2q + 1$ where $q$ is prime. In [69] Christopher Hooley proved that Artin's conjecture would follow from the Generalized Riemann Hypothesis, a result which is thought to be true but which is also considered to be very difficult to prove.

Although it is difficult to find primitive elements, our proof of the Primitive Element Theorem can be combined with trial and error to construct primitive elements. For example, to find a primitive element in the finite field $\mathbb{F} = \mathbb{Z}_{31}$, let us first try $a = 2$. The powers of 2 modulo 31 are $2, 4, 8, 16, 1$ and we obtain $ord(2) = 5$ in $\mathbb{F}$. Inspecting this sequence, we see that 5 does not occur. The powers of 5 modulo 31 are $5, 25, 1$ which gives $ord(5) = 3$ in $\mathbb{F}$. Since $\gcd(5, 3) = 1$ then $ord(2 * 5) = 15$. Also, $ord(-1) = 2$ implies $ord(21) = ord(-10) = 30$, and 21 is a primitive element in $\mathbb{F}$.

If $a$ is a primitive element in a field $\mathbb{F}$ with $q$ elements, for any divisor $d$ the order of $a^k$ is $d$ for $k$ such that $kd = q - 1$.

**Corollary 8.4.9.** *In any finite field with $q$ elements, for any divisor $d$ of $q - 1$ there exists a first-order recurrence in $\mathbb{F}$ whose period equals $d$.*

## 8.5 Periodic Second-Order Recurrences

In the last section we proved that when the modulus is prime the period of any orbit of a first-order modular recurrence $s_{n+1} = as_n + b$ is the same for all non-fixed points, and the period is the order of the coefficient $a$.

For higher order sequences the situation is less clear and seems to be more complicated.

Our analysis for second-order recurrences involves the matrix form of the recurrence. Let $R$ be either a finite field or $\mathbb{Z}_m$ for some $m \geq 2$ and let $< s_j >$ be the solution of a second-order homogeneous recurrence in $R$,

$$s_{n+2} = c_1 s_{n+1} + c_2 s_n \, , \; c_2 \neq 0 \, .$$

Although linear algebra cannot be used here unless $R$ is a field, matrices are still a convenient way to represent higher order recurrences. By Theorem 8.1.6, the periodicity implies that $c_2$ is invertible in $R$. Recall that the **companion matrix of the recurrence** is

$$A = \begin{bmatrix} c_1 & c_2 \\ 1 & 0 \end{bmatrix} \tag{8.8}$$

and consecutive pairs $S_i = (s_{i+1}, s_i)^T$ of state vectors are connected by the matrix equation $S_n = AS_{n-1}$, implying

$$S_n = A^n S_0 \, .$$

When $c_2$ is invertible in $R$,

$$A^{-1} = \begin{bmatrix} 0 & 1 \\ -c_2^{-1} & -c_1 c_2^{-1} \end{bmatrix} \, ,$$

and with this we see that $A$ is an element of the finite set $S$ of invertible $2 \times 2$ matrices with entries in $R$. In Exercise 8.13 you prove that this implies there exists a minimal positive integer $k$ (called the **order of the matrix**) such that $A^k = I$, from which we get $S_0 = S_k = A^k S_0$, proving $k$ is a multiple of the period. Although the order of the matrix can be larger than the period (see Exercise 8.24) we now obtain conditions on the intial values which ensure that the period *equals* the order of the matrix. Let $T$ be the map $T(S) = AS$ defined on $R^2$, where $T$ is a linear map since it satisfies $T(cS + X) = cT(S) + T(X)$ even when $R$ is not a field. If $< s_n >$ is any recurrence with companion matrix $A$ for which the first two state vectors, $S_0 = (s_1, s_0)^T$ and $S_1 = (s_2, s_1)^T$, form a spanning set for $R^2$ then each linear map $T^{(n)}$ is determined by its effect on the set $\{S_0, S_1\}$. In particular, $A^n = I$ iff both $T^n(S_0) = S_0$ and $T^n(S_1) = S_1$, that is, the order of the matrix equals the period of the solution. We've proved

**Theorem 8.5.1.** *Let $R$ be either a finite field or $\mathbb{Z}_m$ for some $m \geq 2$. If $s_{n+2} = c_1 s_{n+1} + c_2 s_n$ is a second-order recurrence in $R$ with invertible $c_2$, then every solution is periodic and the period always divides the order of the companion matrix of the recurrence. Moreover, if $\{S_0, S_1\}$ is a spanning set for $R^2$ then the period equals the order of the companion matrix.*

Since our proof relied only on the invertibility of the companion matrix, an analogue of this result holds for $k^{\text{th}}$-order recurrences. (Refer to Exercise 8.27.)

For the case when $R$ a field and the eigenvalues of $A$ are distinct it can be checked that $S_0, S_1$ are linearly independent and we have

**Theorem 8.5.2.** *Let $\mathbb{F}$ be a finite field and let $s_{n+2} = c_1 s_{n+1} + c_2 s_n$ be a second-order recurrence in $\mathbb{F}$ with two distinct eigenvalues $\lambda_1 \neq \lambda_2$ in $\mathbb{F}$. Then every non-zero solution is periodic and its period equals the order of the companion matrix unless it has the form $s_n = \gamma \lambda_i^n$ in which case the period is $ord(\lambda_i)$.*

### *8.5.1 The period of the modular Fibonacci sequences*

We now specialize to the Fibonacci recurrence, and first answer the questions about the Fibonacci orbits modulo 6 which were posed in the beginning of this chapter. As already noted, every Fibonacci orbit is periodic. Since the periods modulo 2 and modulo 3 are 3 and 8, respectively, the period modulo 6 is $\text{lcm}(3, 8) = 24$, which is consistent with our calculations in (8.2). If $< f_j >$ is the Fibonacci sequence, as noted in Chapter 2 we know the orbit of any $(a, b) \in \mathbb{Z}_m^2$ under $F(x, y) = (y, x + y)$ is the sequence $< bf_{j+1} + af_j >$ (where $f_{-1} = 1$). Therefore, the period of any orbit under $F$ must divide the period of the orbit of $(0, 1)$, and so 24 is the largest period obtained from the Fibonacci recurrence modulo 6.

Our final question was how many *different* orbits are generated by the Fibonacci recurrence modulo 6, where two orbits are considered the same when they're translates of each other. First note that $(0, 0)$ is the only fixed point. Among the $6^2 - 1 = 35$ non-zero elements of $\mathbb{Z}_6^2$, the usual Fibonacci recurrence (which is the orbit of $(0, 1)$) generates 24 different pairs of consecutive elements, and $35 - 24 = 11$ non-zero pairs remain to be accounted for. Considering the orbit of $(2, 2)$, since every element of the orbit is even it suffices to find the period of $(1, 1)$ modulo 3 which we've already calculated to be 8. Because its period is not 24 this orbit is completely disjoint from the orbit of $(0, 1)$. Similarly, since the period of $(1, 1)$ modulo 2 equals 3, the orbit of $(3, 3)$ modulo 6 is 3. These two periods account for the other 11 elements and give a total of three different non-zero orbits under the Fibonacci recurrence.

What about the period of the Fibonacci sequence for other moduli? Although many of the results proved by D.D. Wall [146] have been generalized, his 1960 paper contains most of what is currently known about the periods of modular Fibonacci sequences.

Recalling that the period modulo $m = p_1^{a_1} \dots p_r^{a_r}$ is the lcm of the periods modulo $p_i^{a_i}$, it is enough to consider moduli which are powers of a prime number. Also, if $t = t(p^{j+1})$ is the period of the Fibonacci sequence modulo

$p^{j+1}$ for any $j \geq 1$ then

$$f_t = 0 \pmod{p^{j+1}} \text{ and } f_{t+1} = 1 \pmod{p^{j+1}},$$

and these congruences also hold modulo all divisors of $p^{j+1}$. Therefore, for all $i \leq j$ it is also true that $(f_t, f_{t+1}) \equiv (0,1) \pmod{p^i}$ and we have

$$t(p^{j+1}) \text{ is divisible by } t(p^j).$$

For any modulus $m$, since the first two state vectors, $S_0 = (1,0)^T, S_1 = (1,1)^T$, form a spanning set for $\mathbb{Z}_m^2$, the period modulo $m$ is exactly the order of the companion matrix, $F = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$. Using the fact that $F^3 = I + 2F$, in Exercise 8.26 you show that the matrix $F^{3 \cdot 2^n} - I$ is non-zero modulo $2^n$ and is the zero matrix modulo $2^{n+1}$. The periods $t(2^n) = 3 \cdot 2^{n-1}$ and $t(5^n) = 4 \cdot 5^n$ can be found in a similar manner.

What about other prime powers? It has been conjectured that $t(p^j) = p^{j-1}t(p)$ holds for all prime powers, and this was verified by Wall for all $p \leq 10,000$. His paper would be very long indeed if he performed an induction similar to what we gave above for all primes less than $10,000$. Instead he proved the surprising fact that if $p$ is a prime for which $t(p^2) \neq t(p)$, then it is true that $t(p^{j+1}) = p^j t(p)$ for all $j \geq 1$. Wall used several combinatorial identities to derive his result but we don't include a proof of the result here. In January 2003 Jonathan Goff, a graduate student in mathematics at Oregon State University, extended Wall's calculations by verifying $t(p^2) \neq t(p)$ for all primes less than one million (and all primes $p \equiv \pm 1 \pmod{10}$ less than twenty million.)

Let's look again at the Fibonacci sequence modulo 5. Rather than computing all terms until the second occurrence of $0, 1$ is obtained, we'll consider what happens algebraically. In $\mathbb{Z}_5[x]$ the characteristic polynomial $ch(x) = x^2 - x - 1$ factors as $ch(x) = (x-3)^2$. Since $\lambda = 3$ is a repeated eigenvalue and $\mathbb{Z}_5$ is a field, there exists a polynomial $p$ with $\deg(p) \leq 1$ such that

$$f_n = p(n)3^n.$$

From $f_0 = 0$ and $f_1 = 1$ we obtain $p(n) = 3^{-1}n$, which gives

$$f_n = n3^{n-1} \text{ for all } n \geq 0.$$

If $t = t(5)$ is the period,

$$t3^{t-1} = f_t = f_0 = 0 \quad \text{and} \quad (t+1) \cdot 3^t = f_{t+1} = f_1 = 1,$$

which can be solved to find that $t$ is the least positive integer which simultaneously satisfies $t \equiv 0 \pmod{5}$ and $3^t \equiv 1 \pmod{5}$. Since $t = 4$ is the least positive integer for which $3^t \equiv 1 \pmod{5}$, the period is $\text{lcm}(4,5) = 20$. (Note that this argument cannot be used for $m = 25$ since $\mathbb{Z}_{25}$ is not a field.)

For this argument to work for other primes $p \neq 2, 5$, we'd like to find the eigenvalues of the recurrence modulo $p$ and in order to do that we must factor the characteristic polynomial $ch(x)$ in $\mathbb{Z}_p[x]$. But $ch(x)$ might be an irreducible quadratic or it might factor into a product of two linear polynomials in $\mathbb{Z}_p[x]$. When is $ch(x)$ irreducible? Since a *quadratic* polynomial is irreducible in $\mathbb{Z}_p[x]$ if and only if it has no roots in $\mathbb{Z}_p$, we'll use the technique of completing the square to determine whether or not $ch(x)$ has roots. The oddness of $p$ means that 2 is invertible, say $2c = 1$ in $\mathbb{Z}_p$, and

$$ch(x) = x^2 - x - 1 = x^2 - 2cx - 4c^2 = (x - c)^2 - 5c^2,$$

giving $ch(x+c) = x^2 - 5c^2$. Since $ch(x)$ is irreducible exactly when $ch(x+c)$ is, from this we obtain

$$ch(x) \text{ is reducible in } \mathbb{Z}_p[x] \iff p \neq 2 \text{ and } x^2 - 5c^2 \text{ has roots in } \mathbb{Z}_p\ ,$$

which is the same as requiring that $5c^2$, and so 5, is a square in $\mathbb{Z}_p$. For what primes does this happen? For this type of question, number theorists invoke the famous **Law of Quadratic Reciprocity**, which was first stated by Legendre and proved by Gauss [61, Section 123]. [1] Using quadratic reciprocity, it's possible to prove $p \equiv \pm 1 \pmod{10}$ are exactly the odd primes for which 5 is a square in $Z_p[x]$. These are therefore the primes $p$ for which the Fibonacci recurrence modulo $p$ has two different eigenvalues in $\mathbb{Z}_p[x]$. What about the other odd primes, $p \equiv \pm 3 \pmod{10}$ ? Because for those primes $ch(x)$ is irreducible in $Z_p[x]$, from Exercise 8.11 we know $\mathbb{K} = \mathbb{Z}_p[x]/(ch(x))$ is a field (with $p^2$ elements) in which $ch(x)$ has a root.

We set $\mathbb{K} = \mathbb{Z}_p$ if $p \equiv \pm 1 \pmod{10}$. Then for any prime $p \neq 2, 5$, there exist roots $\lambda_1, \lambda_2$ of the equation $x^2 - x - 1 = 0$ in $\mathbb{K}$. When $\lambda_1 = \lambda_2$ holds, then $(x - \lambda_1)^2 = x^2 - x - 1$, and $\lambda_1^2 = -1$. Since $ch(\lambda_1) = 0$, then $\lambda_1^2 = \lambda_1 + 1$, from which we get $\lambda_1 = -2$. Therefore, $5 = ch(-2) = 0$, yielding $p = 5$. So, for $p \neq 2, 5$ the Fibonacci recurrence modulo $p$ has distinct eigenvalues, $\lambda_1, \lambda_2$. Then $ch(\lambda_1) = 0$ implies

$$A \begin{pmatrix} \lambda_1 \\ 1 \end{pmatrix} = \lambda_1 \begin{pmatrix} \lambda_1 \\ 1 \end{pmatrix}$$

and $(\lambda_1, 1)^T$ is an eigenvector corresponding to $\lambda_1$ and similarly $(\lambda_2, 1)^T$ is an eigenvector which corresponds to $\lambda_2$. The matrix $B = \begin{bmatrix} \lambda_1 & \lambda_2 \\ 1 & 1 \end{bmatrix}$ is invertible with

$$BAB^{-1} = \begin{bmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{bmatrix} \text{ and } BA^jB^{-1} = \begin{bmatrix} \lambda_1^j & 0 \\ 0 & \lambda_2^j \end{bmatrix}.$$

[1] Consult the website `http://www.rzuser.uni-heidelberg.de/~hb3/rchrono.html` for a timeline for the many proofs of the Law of Quadratic Reciprocity.

This means $\text{lcm}(ord(\lambda_1), ord(\lambda_2))$ is the order of the companion matrix and so equals $t(p)$. We already said in Section 8.4.1 that calculating the order of an element is a hard problem. (In particular, refer to our discussion of Artin's conjecture on primitive elements.) Because of that, you might rightfully question the helpfulness of relating the period to the order of certain elements. The answer is that this the connection allows us to prove other results about the period. The next theorem is an example of this. (Other proofs of this result can be found in [146, Theorems 6, 7].)

**Theorem 8.5.3.** *For prime $p \neq 2, 5$, let $t$ be the period of the Fibonacci sequence modulo $p$.*

*(a) If $p \equiv \pm 1 \pmod{10}$ then $t$ divides $p-1$.*

*(b) If $p \equiv \pm 3 \pmod{10}$ then $t$ divides $2(p+1)$ and also $(2p+2)/t$ is odd.*

*Proof.* From $ch(x) = x^2 - x - 1 = (x - \lambda_1)(x - \lambda_2)$ we have $\lambda_1 = -\lambda_2^{-1}$. Since $ord(\lambda) = ord(\lambda^{-1})$ and $ord(-1) = 2$, then

$$\text{lcm}(ord(\lambda_1), ord(\lambda_2)) = \text{lcm}(ord(\lambda_1), 2)$$

and $t = \text{lcm}(ord(\lambda_1), 2)$.

When $p \equiv \pm 1 \pmod{10}$, $\mathbb{K} = \mathbb{Z}_p$ and the order of a non-zero element $\lambda_1 \in \mathbb{K}$ divides $p-1$, an even number, and so $\text{lcm}(ord(\lambda_1), 2)$ divides $p-1$. In the second case the field $\mathbb{K}$ contains an element $\alpha$ such that $\alpha^2 = 5 \in \mathbb{Z}_p$ and so $\alpha^{2(p-1)} = 1$. This means $x = \alpha^{p-1}$ is a root of the polynomial $x^2 - 1 = 0$, which has the two roots $x = \pm 1$ in any field. Since $\alpha \notin \mathbb{Z}_p$, we know $\alpha^{p-1} \neq 1$ (refer to Theorem 8.4.6) and so $\alpha^{p-1} = -1$. What does this have to do with $ord(\lambda_1)$? In the "completing the square" argument above we had

$$ch(x) = (x - c)^2 - 5c^2, \text{ where } 2c = 1\,,$$

implying $\lambda_1 = c(1 \pm \alpha)$ where we can assume the positive sign occurs. Since $\mathbb{K}$ contains has $p^2$ elements by Exercise 8.6 we know

$$(1 + \alpha)^p = 1 + \alpha^p = 1 + \alpha\alpha^{p-1} = 1 - \alpha\,.$$

This combined with $c^{p-1} = 1$ gives

$$\lambda_1^{p+1} = c^{p+1}(1 - \alpha)(1 + \alpha) = c^2(1 - \alpha^2) = c^2(1 - 5) = -(2c)^2 = -1\,,$$

then $\lambda_1^{2(p+1)} = 1$ and so $t_1 = ord(\lambda_1)$ divides $2(p + 1)$. Further, writing $2(p + 1) = 2^a n$ where $n$ is odd, we know $t_1$ divides $2^a n$ and doesn't divide $2^{a-1} n$ (since $\lambda^{p+1} \neq 1$). This means that $t_1$ is even, and $t = \text{lcm}(t_1, 2) = t_1$ with $(2p + 2)/t$ odd. □

When the periods of the Fibonacci sequence modulo $p$ are computed for primes less than 1000, the upper bound in the theorem is attained quite often. Seventy-eight of these primes are congruent to $\pm 1 \pmod{10}$, and for forty of them the period equals $p - 1$. The situation for the "irreducible"

primes is even stronger. Among primes less than 1000 there are eighty-eight primes congruent to $\pm 3 \pmod{10}$ and for all but fifteen of these the upper bound of $2p+2$ is the period. We list the irreducible primes with smaller periods in Table 8.2 where the second column gives the quotient $(2p+2)/t$.

TABLE 8.2. The exceptional primes congruent to $\pm 3 \pmod{10}$

| $p$ | $(2p+2)/t$ |
|---|---|
| $47, 107, 113, 233, 353, 563, 677, 743, 977$ | 3 |
| $307, 797$ | 7 |
| $263, 557, 953$ | 9 |
| $967$ | 11 |

Let us compute the period $t(m)$ of the Fibonacci recurrence modulo $m = 2^3\,3^2\,17\,47^2$. Corollary 8.1.4 implies $t(m) = \text{lcm}(t(8), t(9), t(17), t(47^2))$ where by Wall's calculations we know that $t(8) = 2^2\,t(2)$, $t(9) = 3\,t(3)$ and $t(47^2) = 47\,t(47)$. ¿From earlier calculations given in Section 8.1, we have $t(2) = 3$ and $t(3) = 8$. Since the remaining two primes are congruent to $-3 \mod 10$, from Table 8.2 we see that $t(47) = 2(48)/3 = 32$ and that 17 is not exceptional which means $t(17) = 2(18) = 36$. Combining these facts gives

$$t(m) = \text{lcm}(12, 24, 36, (32)(47)) = (32)(9)(47) = 13536\,.$$

## 8.6 Applications

### *8.6.1 Application 1: Pseudorandom number generation*

Random numbers are often needed in scientific computation, especially in computations that have some stochastic components such as simulations or probabilistic algorithms. They are also used in cryptology and even computer games. The term "random sequence" means a sequence which passes certain statistical tests. For instance, one simple randomness requirement for sequences from the set $\mathbb{Z}_{10}$ might be the property that each digit should occur about the same number of times so that in suitably long subsequences each digit occurs close to one-tenth of the time. In contrast with sequences generated mechanically or by a physical phenomenon, sequences generated by a mathematical iterative process (a recurrence) can be *proved* to have good statistical properties. Another advantage of such deterministic procedures is that they will dependably generate the same sequence when the initial conditions are unchanged. This allows for the exact reproduction of data for numerical experiments and reduction in storage space. Sequences which are generated deterministically and have good statistical properties are called **pseudorandom numbers (PRN)**. Because a PRN generator cannot satisfy all possible statistical properties (which is another reason for

calling them *pseudo*random), the practitioner should know which statistical properties are required for the application before choosing the generator to use. Chapter 3 of Knuth [77] and Chapter 7 in Niederreiter [103] can be consulted for information on the common statistical tests used to test for pseudorandomness. In this section we concentrate on what could be called *structural* properties, in particular, period length and lattice structure.

A linear modular recurrence is the simplest example of a deterministic modular process which can be used for generating PRN. These are frequently called **linear congruential generators**. From the perspective of PRN generation, the Primitive Element Theorem guarantees that every prime $p$ has at least one linear PRN generator modulo $p$ whose period is $p-1$, the longest possible period. Knuth [77, Chapter 3] has an extensive discussion of the periods for general moduli. For many years after their introduction by D.H.Lehmer [83] in 1949, the linear congruential generator was the preferred method of PRN generation. However, an article by G. Marsaglia [93] raised some serious questions about their use by proving that every linear PRN generator has an inherent lattice structure. As he said in the paper, "...for the past 20 years such regularity might have produced bad, but unrecognized, results in Monte Carlo studies ...." In a later article [94] he developed the **Lattice Test for PRN** which can be stated as follows. For any deterministic sequence $< s_n >$ (with period $N \geq 2$) generated in a finite field $\mathbb{F}$ and each $d \geq 1$, define the $d$-dimensional points $U_1, \ldots, U_{N-1}$ by

$$U_i = (s_i - s_0, s_{i+1} - s_1, \cdots, s_{i-1+d} - s_{d-1}) \in \mathbb{F}^d ,$$

and let $L_d$ be the subspace of $\mathbb{F}^d$ spanned by these points. For instance, $L_1 = \text{Span}\{U_1\} = \text{Span}\{s_1 - s_0\} = \mathbb{F}$ (since $s_1 \neq s_0$) and $L_2 = \text{Span}\{U_1, U_2\} = \text{Span}\{(s_1 - s_0, s_2 - s_1), (s_2 - s_0, s_3 - s_1)\}$, which may have dimension one or two over $\mathbb{F}$. The **lattice dimension** for the sequence $< s_n >$ is defined to be the largest integer $D \geq 1$ for which $L_D = \mathbb{F}^D$. When the lattice dimension is small, the points display a high degree of regularity and so the generation is predictable and not random. A linear generator fares poorly under this test, since for $f(x) = ax + b \bmod p$ the points $U_i$ have the form $U_i = (s_i - s_0)(1, a, a^2, ..., a^{d-1})$, which means its lattice dimension always equals 1.

The remainder of this discussion on PRNs is devoted to an analysis of a commonly used type of *nonlinear* generator whose lattice dimension modulo $p$ has been proved to be at least $(p+1)/2$, much better than linear generators. The generator is called **an inversive generator on the finite field** $\mathbb{F}$ and has the form $f(x) = ax^{-1} + b$ where we set $0^{-1} = 0$ but otherwise the arithmetic is performed in $\mathbb{F}$. Because $f$ is a one-to-one function on $\mathbb{F}$, any sequence generated by the first-order recurrence $x_{n+1} = f(x_n)$ defined by an inversive generator $f$ is periodic and the number of elements in $\mathbb{F}$ is an upper bound on the period. Below we prove that in any finite field there are inversive generators which have this maximal period.

Although the Euclidean Algorithm can be used to compute inverses quickly, finding inverses takes longer than performing the operations for polynomial functions and so inversive generators are slower to compute with. On the other hand, research has shown inversive generators do usually have better statistical properties than polynomial generators. (Refer to [103, Chapter 8].) For instance, extensive computations by Poul Petersen [110] show that for all primes $p \leq 10^5$ the lattice dimension of inversive generators modulo $p$ with maximal period is either $p-6$, $p-4$ or $p-2$. (Table 8.3 gives more details. In [55] it's shown the lattice dimension of a maximal period inversive generator is always odd.)

TABLE 8.3. Lattice dimensions for maximal period inversive generators (mod $p$) for $5 \leq p < 10^5$

| | |
|---|---|
| Number of primes less than $10^5$ | 9590 |
| Number of maximal period inversive generators | $5,579,945,320,208$ |
| Number whose dimension is $p-6$ | 1 |
| Number whose dimension is $p-4$ | 1829 |
| Number whose dimension is $p-2$ | $5,579,945,318,378$ |

As said above, we will show that every finite field has inversive generators which have only one orbit and we'll also characterize the generators which have this maximal period. The polynomial $ch_I(x) = x^2 - bx - a$ associated with the generator $f(x) = ax^{-1} + b$ is useful in this analysis.

For example, the generator $f(x) = 3x^{-1} + 2$ in $\mathbb{F} = \mathbb{Z}_{11}$ has $ch_I(x) = x^2 - 2x - 3$, which factors as $ch_I(x) = (x-3)(x-10)$ in $\mathbb{F}[x]$. Its roots $x = 3, 10$ are the fixed points of $f$, and every other element of $\mathbb{F}$ is in the orbit of $x = 2$, which is $2, 9, 6, 8, 1, 5, 7, 4, 0$. This example illustrates a general property of inversive generators, that its fixed points are precisely the roots of $ch_I(x)$ in the field $\mathbb{F}$. To see this, we first note that the fact that $a$ is non-zero implies that 0 is not a fixed point of $f$ and is not a root of $ch_I(x)$. Therefore,

$$c^2 - bc - a = 0 \quad \Longleftrightarrow \quad c^2 = bc + a \quad \Longleftrightarrow \quad c = b + ac^{-1} = f(c),$$

the last obtained from multiplying by $c^{-1}$.

Because our goal is to find inversive generators with maximal period, we will consider generators which have no fixed points, which means that $ch_I(x)$ is irreducible in $\mathbb{F}[x]$. As we used earlier with the Fibonacci recurrence, $\mathbb{K} = \mathbb{F}[x]/(ch_I(x))$ is then a field, $ch_I(x)$ factors completely as $ch_I(x) = (x-\alpha)(x-\beta)$ in $\mathbb{K}[x]$, and we ensure the roots are distinct by requiring that the discriminant $b^2 + 4a$ is non-zero. Although the recurrence is not linear, we get a relation between the orbit of 0 and the roots of $ch_I(x)$, namely, we claim that for every $j \geq 0$ with $f^{(j-1)}(0) \neq 0$,

$$f^{(j)}(0) = \frac{\alpha^{j+1} - \beta^{j+1}}{\alpha^j - \beta^j}. \tag{8.9}$$

From $ch_I(x) = (x-\alpha)(x-\beta) = x^2 - bx - a$ it is true that $a = -\alpha\beta$ and $b = \alpha + \beta$ and this gives

$$f(0) = b = \alpha + \beta = \frac{\alpha^2 - \beta^2}{\alpha - \beta}.$$

Further, if for $j \geq 1$ we have $f^{(j)}(0) = \dfrac{\alpha^{j+1} - \beta^{j+1}}{\alpha^j - \beta^j} \neq 0$, then

$$f^{(j+1)}(0) = -\alpha\beta\frac{\alpha^j - \beta^j}{\alpha^{j+1} - \beta^{j+1}} + \alpha + \beta = \frac{\alpha^{j+2} - \beta^{j+2}}{\alpha^{j+1} - \beta^{j+1}},$$

and the $(j+1)^{\text{st}}$ element of the orbit of 0 is as given in (8.9).

**Theorem 8.6.1.** *Let $\mathbb{F}$ be a finite field with $q$ elements and let $f$ be an inversive generator on $\mathbb{F}$ such that $ch_I(x)$ is irreducible in $\mathbb{F}[x]$ and has non-zero discriminant. If $t$ is the period of $x = 0$ under $f$, then the order of $\beta^{q-1}$ in $\mathbb{F}[x]/(ch_I(x))$ is $t+1$.*

*Proof.* Since $f$ is a one-to-one function, every orbit of $f$ is periodic and $t$ is the least positive integer such that $f^{(t)}(0) = 0$. As above, the irreducibility of $ch_I(x)$ implies $\mathbb{K} = \mathbb{F}[x]/(ch_I(x))$ is a field which contains elements $\alpha, \beta$ which are not in $\mathbb{F}$ such that (8.9) holds. Therefore,

$$f^{(t)}(0) = 0 \iff \alpha^{t+1} - \beta^{t+1} = 0 \iff (\alpha\beta^{-1})^{t+1} = 1$$

and $t$ is the minimal positive integer with this property. Therefore, $t+1$ is the order of $\alpha\beta^{-1}$ in $\mathbb{K}$. To complete the proof we show that $\alpha = \beta^q$. From Exercise 8.6 we know that $ch_I(\beta^q) = (ch_I(\beta))^q$ which means that $\beta^q$ is also a root of $ch_I(x)$. By Exercise 8.15 the fact that $\alpha \notin \mathbb{F}$ implies $\beta^q \neq \beta$, and $\beta^q$ is forced to be the other root of $ch_I(x)$, and $\alpha\beta^{-1} = \beta^{q-1}$ has order $t+1$. □

As with first-order linear recurrences, we've again related the period of a recurrence to the order of an element in an associated algebraic structure. Because calculating the order of an element can be a quite lengthy process, this result is more useful for theoretical rather than computational purposes. For instance it's used in [55] to show that the lattice dimension of an inversive generator is always odd.

As an example, let's consider the orbit of 0 in $\mathbb{F} = \mathbb{Z}_{11}$ under $f(x) = 7x^{-1} + 10$. The polynomial is $ch_I(x) = x^2 + x + 4$, which can be checked to be irreducible in $\mathbb{F}[x]$ with non-zero discriminant. We won't explicitly determine the field $\mathbb{K} = \mathbb{F}[x]/(ch_I(x))$, but rather recall that it's a finite field with $(11)^2 = 121$ elements in which $ch_I(x)$ factors completely. Since we want to calculate or otherwise determine the order of $\gamma = \beta^{q-1} = \beta^{10}$ where $\beta \in \mathbb{K}$ is a root of $x^2 + x + 4$ and the order of $\beta$ divides $q^2 - 1 = 120$,

which means the order of $\gamma$ divides 12. We first want to find $\gamma = \beta^{10}$ using the technique of **fast exponentiation**. From $\beta^2 = -\beta - 4$ we have

$$\beta^4 = (\beta+4)^2 = 7\beta + 1 \quad ; \quad \beta^8 = (7\beta+1)^2 = -2\beta + 3\,,$$

and

$$\gamma = \beta^8 \cdot \beta^2 = (-2\beta+3)(-\beta-4) = 3\beta + 2\,.$$

Since

$$\gamma^2 = (3\beta+2)^2 = 3\beta + 1 \quad ; \quad \gamma^3 = (3\beta+1)(3\beta+2) = -1,$$

the order of $\gamma = \beta^{q-1}$ equals 6 and the period of 0 is five. This can also be verified by direct calculation, tracing the orbit of 0 as $0, 10, 3, 5, 7$.

**Theorem 8.6.2.** *Every finite field has inversive generators with one orbit. The associated pseudorandom sequence contains every element of* $\mathbb{F}$ *and so has the maximal period among all inversive generators on* $\mathbb{F}$.

*Proof.* Let $\mathbb{F}$ be a finite field with $q$ elements, and let $g$ be any irreducible quadratic polynomial in $\mathbb{F}[x]$ whose discriminant is non-zero. From the Primitive Element Theorem, there exists $\gamma \in \mathbb{F}[x]/(g)$ whose order equals $q^2-1$. Therefore, $\gamma \notin \mathbb{F}$ and the quadratic polynomial $G(x) = (x-\gamma)(x-\gamma^q)$ is irreducible in $\mathbb{F}[x]$. For $G(x) = x^2 - Bx - A$, we consider the associated inversive generator $f(x) = Ax^{-1} + B$ which has $ch_I(x) = G(x)$. Since $ord(\gamma) = q^2 - 1$ then $ord(\gamma^{q-1}) = q + 1$, and the period of $x = 0$ under $f$ does equal $q$. □

### *8.6.2 Application 2: Integer Factorization*

Because the security of RSA [117] relies on the difficulty of factoring integers that are a product of two large primes, factorization algorithms gained widespread interest in the late 1970's when the RSA cryptosystem was introduced.

In this section we discuss one of the early modern methods for factoring integers. The method is called **Pollard's Rho Method** because it was developed by Pollard and produces eventually periodic orbits, which can be drawn to have the shape of the Greek letter $\rho$. (Refer to Figure 8.1.) In the original article [111], Pollard called this algorithm the Monte Carlo Method because it's probabilistic and uses an orbit of a "sufficiently random" function $f(x)$. In practice, a quadratic polynomial, often $f(x) = x^2 + 1$, is used.

What do we mean when we say Pollard Rho is a *probabilistic* method? One feature is that although it's likely that the method will return a factorization, the method might be not be successful for the first choice of initial value $s_0$. Because of this when you use this algorithm you must

already know for certain that $m$ is not prime. At first this might seem circular, since the most obvious way of showing a number is composite is to demonstrate a factorization. Luckily, the contrapositive of Fermat's Little Theorem yields a sufficient condition for compositeness: If $m$ is an integer such that there exists non-zero $a \in \mathbb{Z}_m$ for which $a^{m-1} \neq 1 \pmod{m}$, then $m$ is not prime. This is only a sufficient condition, since for instance $8^{20} = 1 \pmod{m}$ holds for the composite number $m = 21$. A composite integer $m$ for which $a^{m-1} = 1 \pmod{m}$ for $a \neq 1$ is called a **pseudoprime to the base** $a$, because it is a composite number which behaves as if it's a prime for the base $a$. In 1950 Paul Erdös [50] proved that pseudoprimes are relatively rare compared to primes, which says that this is a useful test for verifying compositeness. On the other hand, in 1994 W.R. Alford, Andrew Granville and Carl Pomerance [3] showed the existence of infinitely many composite $m$ (called **Carmichael numbers**) which are pseudoprimes to every possible base. The existence of these numbers lessens the usefulness of the Fermat Test for checking compositeness. But all is not lost, because we can invoke our theory and obtain a slight variant called the **Strong Fermat Test** which doesn't have this problem. The idea behind the test is the following. If $m$ were prime, then by Fermat's Little Theorem, $b = a^{\frac{m-1}{2}}$ would be a solution to $x^2 - 1 = 0$, a quadratic equation which can have only the two solutions $x = \pm 1$ in the field $\mathbb{Z}_m$. Therefore, if we can find a non-zero base $a$ such that $a^{\frac{m-1}{2}}$ is something other than $\pm 1 \pmod{m}$, we are assured that $m$ is not prime. Because the exponent is slightly smaller, this test is a bit easier to perform than the Fermat Test. A more important property is that there are no "strong Carmichael" numbers. In fact, every odd composite $m > 9$ is a strong pseudoprime for at most one-quarter of the bases modulo $m$. (If the widely believed but still unproved the Generalized Riemann Hypothesis were true, we'd be guaranteed that every odd composite $m$ passes a Strong Fermat Test for at least one positive base $a < 2\ log^2(m)$.) In order to guarantee that $m$ is composite it suffices to determine one base $a$ for which $a^{m-1} \neq \pm 1 \pmod{m}$. For example, the Strong Fermat Test with $a = 2$ proves $m = 18923$ is composite since $\frac{m-1}{2} = 9461$ and $2^{9461} = 8144$ in $\mathbb{Z}_m$. (Refer to Chapter 3 in [24] for more information on tests for compositeness.)

Once we've obtained a certificate of compositeness for $m$, how does Pollard's Rho Method find a divisor of $m$? The basic idea is that we want to generate a sequence which repeats modulo some divisor of $m$ well before it begins to repeat modulo $m$ AND we'd like some way of detecting congruent pairs in this sequence. Once we have such pairs, the gcd can be used to find a divisor. For instance, suppose we'd like to factor $m = 712091$ (which has prime factorization $m = 509 * 1399$). If we had some way of identifying the pair $(a, b) = (459, 1986)$ as a likely pair we could calculate $\gcd(b - a, m) = 509$ and so obtain the factor 509 of $m$.

The first idea behind Pollard's Rho Method is that the sequence must be sufficiently random that orbits under $f$ are "very likely" to become periodic modulo a divisor $d$ of $m$ before they become periodic modulo $m$. Writing this in terms of iteration, for $s \in \mathbb{Z}_m$ there exists a divisor $d$ of $m$ such that there are integers $n, k$ (say $n < k$) such that

$$f^{(n)}(s) = f^{(k)}(s) \pmod{d}$$

but

$$f^{(i)}(s) \neq f^{(j)}(s) \pmod{m}) \text{ for all } 0 \leq i, j \leq k\,.$$

If this were the case, then $d$ is a common divisor of both $m$ and $f^{(n)}(s) - f^{(k)}(s)$ and so must necessarily divide their gcd. On the other hand, the second restriction says that $f^{(n)}(s) - f^{(k)}(s)$ is not divisible by $m$. Therefore we've found a nontrivial divisor of $m$.

Recognizing that it's not necessary to find the *smallest* values of $n$ and $k$ for which $f^{(n)}(s) = f^{(k)}(s) \pmod{d}$, popular implementations of the method use the idea of combining a "fast" sequence such as $< f^{(2n)}(s) >$ with the "slow" sequence $< f^{(n)}(s) >$ and then using the sequence $\gcd(f^{(n)}(s) - f^{(2n)}(s) \pmod{m}, m)$. Here's the algorithm.

```
Procedure POLLARD(m, f(x))
          Randomly generate s.
          t := f(s);
          FOR i from 1 to m
                    DO d := gcd(s − t, m);
                       IF d ≠ 1 THEN return d;
                                ELSE s := f(s); t := f^(2)(t);
```

When $d = m$ is obtained, the procedure should be repeated using another choice of seed $s$. Also remember that $m$ must be a certified composite before using Pollard's algorithm.

We already know that $m = 18923$ is composite by the Strong Fermat Test with $a = 2$. Let's now use the Pollard Rho Method with $f(x) = x^2 + 1$ to factor $m$. For instance, the first few elements of the orbit of 2 under $x^2 + 1 \pmod{m}$ are:

$$2, 5, 26, 677, 4178, 8679, 11502, 5312, 3152, 530, 15979, 403, 11026, 11325\,.$$

Applying the Pollard Rho Test gives $d_n = 1$ for all $n \leq 8$ and then the divisor $d_9 = 127$. From this example you can see that the gcd sequence can begin with a rather long string of ones. Because of this, implementations of the method often take giant steps through the gcd sequence, and by calculating products of several $f^{(n)}(s) - f^{(2n)}(s)$ before computing the gcd.

Even for fairly random functions $f$ there can be orbits which won't yield a divisor. Because of this, it would be useful to know when an orbit should be abandoned because it's unlikely to yield a factorization. This translates to wanting a good estimate of when the first duplication is likely to occur for a modulus. In mathematical terms, the probability that $k$ elements chosen from a set with $m$ elements has a repeated element is $1 - P(k, m)$ where

$$P(k,m) = \frac{m \cdot (m-1) \cdot \ldots \cdot (m-(k-1))}{m^k} = \Big(1 - \frac{1}{m}\Big) \ldots \Big(1 - \frac{k-1}{m}\Big)$$

is the probability that the $k$ choices are different. The next theorem finds fairly tight bounds on this probability. It's often referred to as "The Birthday Problem" because it can be used to estimate the probability that at least two people in a crowd of $M$ people have the same birthday. (Refer to Exercise 8.35.)

**Theorem 8.6.3 (The Birthday Problem).** *For any $1 \leq k < m$,*

$$\exp\Big(-\frac{k^2}{m}\Big) < P(k,m) < \exp\Big(-\frac{k(k-1)}{2m}\Big), \tag{8.10}$$

*where $exp(x) = e^x$, the usual exponential function.*

*Proof.* Since $\exp(x)$ is an increasing function, it suffices to prove

$$\frac{k(k-1)}{2m} < -\ln(P(k,m)) < \frac{k^2}{m}. \tag{8.11}$$

We note that for $H(x) = -\ln(1-x)$

$$-\ln(P(k,m)) = H\Big(\frac{1}{m}\Big) + \ldots + H\Big(\frac{k-1}{m}\Big).$$

Since $H(x)$ is an increasing function on $[0,1)$, which is concave upward, H(0)=0, and $H'(0) = 1$ then the graph of $H(x)$ lies above the line $y = x$ and

$$H\Big(\frac{1}{m}\Big) + \ldots + H\Big(\frac{k-1}{m}\Big) > \frac{1}{m} + \ldots + \frac{k-1}{m} = \frac{k(k-1)}{2m},$$

which is the left side of (8.11).

Again using the monotonicity of $H(x)$, for $b = k/m$ we have

$$\int_0^b H(t)\ dt > \frac{1}{m}\Big(H\Big(0\Big) + H\Big(\frac{1}{m}\Big) + \ldots + H\Big(\frac{k-1}{m}\Big)\Big)\ ,$$

and integration by parts gives

$$\int_0^b H(t)\ dt = (1-b)\ln(1-b) + b = -(1-b)H(b) + b < -(1-b)b + b = b^2.$$

Therefore, $H(0) + H(\frac{1}{m}) + \ldots + H(\frac{k-1}{m}) < mb^2 = k^2/m$, which is the other inequality in (8.11). □

For $m \geq 9$, the right side of (8.10) is less than $\frac{1}{2}$ for $k > 1.39\sqrt{m}$; the left side of (8.10) is greater than $\frac{1}{2}$ for $k < .8\sqrt{m}$.

**Corollary 8.6.4.** *Let $1 \leq k < m$ be integers and $m \geq 9$.*
*(a) If $k > 1.39\sqrt{m}$ then $P(k,m) < \frac{1}{2}$.*
*(b) If $k < .8\sqrt{m}$ then $P(k,m) > \frac{1}{2}$.*

From this we obtain the following heuristic

**Corollary 8.6.5.** *Let $m \geq 10$ be a known composite which needs to be factored. Let $f(x)$ be a function which is considered to be sufficiently random modulo $m$ to be used in Pollard's Rho Method. For any initial value $s_0$, the algorithm should be stopped if it doesn't yield a divisor within $1.39\sqrt[4]{m}$ iterations.*

*Proof.* Remember that we don't want a duplication modulo $m$. From the last result a duplication modulo $m$ is unlikely before $.8\sqrt{m}$ iterations. On the other hand, since $m$ is composite it has a divisor $d$ within the interval $(1, \sqrt{m})$. From the first estimate above, we expect a duplication modulo $d$ within $1.39\sqrt{d}$ steps, which gives within $1.39\sqrt[4]{m}$. For $m \geq 10$ we have $.8\sqrt{m} > 1.39\sqrt[4]{m}$. □

## 8.7 Exercises

**Ex 8.1.** For any positive integer $m$, prove there are infinitely many Fibonacci numbers which are divisible by $m$.
HINT: Consider the Fibonacci sequence modulo $m$.

**Ex 8.2.** Find all periodic and eventually periodic points of the recurrence $x_{n+1} = f(x_n)$ where $f(x) = -x^2 + 2x + 1$.

**Ex 8.3.** (a) Prove $\gcd(19, 157) = 1$. Find an integer solution to $19x + 157y = 1$. and show there exist no positive integer solutions to $19x + 157y = 1$.
(b) Find the multiplicative inverse of 214 modulo 12323.
(c) Prove there are integer points $(x, y)$ on the line $179x + 2351y = 1$. Find the integer point on the line which is closest to the origin.

**Ex 8.4.** For the finite field $\mathbb{F}$, define $n \cdot 1$ to be the sum of an integer number $n$ copies of the multiplicative identity 1 in $\mathbb{F}$.
(a) Show that there exists a least positive integer $n_0$ such that $n_0 \cdot 1 = 0$ where 0 is the additive identity in $\mathbb{F}$.
(b) Show that $n_0$ must be prime. This is called the **field characteristic** of $\mathbb{F}$.

**Ex 8.5.** Let $\mathbb{F}$ be a finite field of characteristic $p$. Show that $\mathbb{F}$ can be regarded as a finite dimensional vector space over $\mathbb{Z}_p$ and therefore $\mathbb{F}$ has $p^n$ elements for some $n \geq 1$.

**Ex 8.6.** Let $\mathbb{F}$ be a finite field with $q = p^n$ elements. Prove that for any $f \in \mathbb{F}[x]$, $(f(x))^p = f(x^p)$ and from this $(f(x))^q = f(x^q)$.
HINT: Use the Binomial Theorem.

**Ex 8.7.** Let $\mathbb{F}$ be any finite field and $n$ be a positive integer. Let $V$ be the set of all polynomials in $\mathbb{F}[x]$ of degree less than $n$.

(a) Prove that $V$ is an abelian group under the operation of polynomial addition in $\mathbb{F}[x]$.
(b) Prove that $V$ is a vector space of dimension $n$ over $\mathbb{F}$. Therefore, the number of elements in $V$ equals $q^n$, where $q$ is the number of elements in $\mathbb{F}$.

**Ex 8.8.** Let $g(x) \in \mathbb{F}[x]$ be a fixed polynomial, with $n = \deg(g)$. Let $V$ be the set of all polynomials in $\mathbb{F}[x]$ whose degree is less than $n$, which was shown to be a vector space in the last exercise. Impose still more structure on $V$ by defining an additional operation $\star$ as follows. For any $a(x), b(x) \in V$ we use the Division Algorithm for polynomials (refer to Section 4.3) to obtain $q(x), r(x) \in \mathbb{F}[x]$ such that

$$a(x)b(x) = q(x)g(x) + r(x) \quad \text{where } \deg(r) < n\,,$$

and define the operation $\star$ on $V$ by $a(x) \star b(x) = r(x) \in V$. Show that $\star$ is an associative commutative operation on $V$ which is distributive over the usual addition defined as in Exercise 8.7. Is $V$ always a field under the operations of $+$ and $\star$? In what follows we use the notation $\mathbb{F}[x]/(g(x))$ for $V$.

**Ex 8.9.** Let $\mathbb{F} = \mathbb{Z}_7$ and $g(x) = (x^2+1)(x^3+x+1)$. Show that neither $x^2+1$ nor $x^3+x+1$ has a multiplicative inverse in $F[x]/(g(x))$.

**Ex 8.10.** Let $\mathbb{F}$ be any finite field and let $g(x)$ be an **reducible polynomial in** $\mathbb{F}[x]$; that is, there exist polynomials $a(x), b(x) \in \mathbb{F}[x]$ such that $g(x) = a(x)b(x)$ with $1 \leq \deg(a), \deg(b) < \deg(g)$. Show that neither $a(x)$ nor $b(x)$ has an inverse under $\star$ in $\mathbb{F}[x]/(g(x))$.

**Ex 8.11.** Let $\mathbb{F}$ be any finite field. Let $g(x)$ be a polynomial which is not reducible in $\mathbb{F}[x]$. Prove that $L = \mathbb{F}[x]/(g(x))$ is a finite field under the operations of addition and $\star$.

**Ex 8.12.** Verify that $x^2+1$ is irreducible in $\mathbb{Z}_3[x]$ and use this to construct a field with nine elements.

**Ex 8.13.** In Section 8.3 we defined an *abelian* group. When the requirement of commutativity is removed the structure is called a **group**. Let $R$ be either a finite field or $\mathbb{Z}_m$ for some $m \geq 2$.

(a) Show that the set $S$ of all $2 \times 2$ invertible matrices with entries in $R$ is a finite group under multiplication.
(b) Show that for any $A \in S$ there exists a positive integer $k$ such that $A^k = I$, which is called the **order** of $A$ in the group $S$. (For this, you

might mimic the argument that every invertible element of $\mathbb{Z}_m$ has an order.)

**Ex 8.14.** Find the orbits of $x = 3$ and $x = 4$ under $S(x) = 12x + 4$ on $R = \mathbb{Z}_{21}$. Formulate a conjecture about the periods of all orbits under this map. Check your conjecture by computing the orbits.

**Ex 8.15.** Suppose $\mathbb{F}, \mathbb{K}$ are finite fields with $\mathbb{F} \subseteq \mathbb{K}$, and let $q$ be the number of elements in $\mathbb{F}$. For $\alpha \in \mathbb{K}$, show $\alpha \in \mathbb{F}$ iff $\alpha^q = \alpha$. This is a generalization of Theorem 8.4.6.

For the next three exercises consider a deck $D$ with $n$ cards and denote the set of all permutations of $D$ by $\mathcal{S}_n$.

**Ex 8.16.** Let $S_n$ denote the set of all permutations of $n$ elements, where a **permutation** means a reordering (or bijection) of the integers $1, 2, \ldots, n$. Show that $\mathcal{S}_n$ is a group under composition of functions. What is the identity element in this group?

**Ex 8.17.** Now consider the action of shuffling a deck of $n$ cards. Every shuffle of $n$ cards can be viewed as an element of $\mathcal{S}_n$, and so every shuffle has an order in the group $\mathcal{S}_n$. What is the practical meaning of the order of a shuffle?

**Ex 8.18.** For this problem consider $\mathcal{S}_{12}$, the group of shuffles of a deck with twelve cards. We'll call a shuffle **perfect** if the shuffle begins by splitting the deck into two equal piles and then alternates between the two piles with the card on the top of the original second pile becoming the bottom card. For instance, an example of a perfect shuffle of six cards changes $1, 2, 3, 4, 5, 6$ into $4, 1, 5, 2, 6, 3$.

(a) Prove that after $i$ perfect shuffles the original first card is in the $2^i$ position modulo 13. After how many shuffles is the first card returned to its original place?

(b) What is the order of a perfect shuffle in $\mathcal{S}_{12}$?

**Ex 8.19.** What is the order of a perfect shuffle in $\mathcal{S}_{52}$?

**Ex 8.20.** (This problem is based on the work of Joseph Keller in [72].) Consider a "riffle shuffle" of a deck with $k$ cards. By this we mean cutting the deck once at random and then riffling together the two parts formed by the cut. For this we assume that cutting satisfies a uniform distribution, that is, the probability of any cut is $1/(k-1)$. Let $p_n$ be the probability that the original bottom card is on the bottom of the deck after $n$ riffle shuffles.

(a) Show $p_0 = 1$ and for all $n \geq 0$

$$p_{n+1} = \frac{1}{2}\left(p_n + (1 - p_n)\frac{1}{k-1}\right).$$

(b) Use the theory of linear recurrences to show

$$p_n = \frac{1}{k} + \frac{k-1}{k}\left(\frac{k-2}{2(k-1)}\right)^n.$$

**Ex 8.21.** For this exercise use $\mathbb{F} = \mathbb{Z}_p$ where p equals the prime 5009. Without calculating any orbits, find the period of $x = 0$ under each of $f(x) = 3x + 1$; $f(x) = 4x + 7$.

**Ex 8.22.** Use induction to show that the number of solutions to any polynomial equation with coefficients in a field is bounded by the degree of the equation.

**Ex 8.23.** For this exercise, use $\mathbb{F} = \mathbb{Z}_p$, where p equals the prime 1361. Check that 3 is a primitive element in $\mathbb{F}$. Find a first-order linear recurrence modulo $p$ whose period equals 85.

**Ex 8.24.** Let $A$ be the companion matrix for the Fibonacci recurrence modulo 6.

(a) What is the order of the matrix $A$?

(b) Let $< s_n >$ be the sequence which satisfies the Fibonacci recurrence modulo 6 and has initial state vector $(3, 3)$. Find the period of this sequence by considering the sequence modulo each of the primes 2 and 3.

**Ex 8.25.** (a) Use induction to prove that the Fibonacci sequence $(f_j)$ satisfies

$$f_{2n} = f_n(2f_{n-1} + f_n) \text{ and } f_{2n+1} = f_n^2 + f_{n+1}^2.$$

HINT: Prove the two identities in tandem.

(b) Use part (a) to show that the Fibonacci period modulo $2^{n+1}$ is $3 \cdot 2^n$.

**Ex 8.26.** For this problem let $F$ be the companion matrix for the Fibonacci recurrence. Show that $F^3 = I + 2F$ and use this to find the period of the Fibonacci sequence modulo $2^n$.

**Ex 8.27.** Let $R$ be either $\mathbb{Z}_m$ or a finite field. Let $s_{n+k} = c_1 s_{n+k-1} + \ldots + c_k s_n$ be a linear recurrence in $R$ with $c_k$ an invertible element of $R$.

(a) Show the companion matrix is invertible.

(b) If the first $k$ state vectors form a spanning set for $R^k$ show that the period of any non-zero solution to the recurrence equals the order of the matrix.

**Ex 8.28.** Let $\mathbb{F}$ be a finite field and $< s_n >$ be a non-zero sequence which satisfies a homogeneous second-order recurrence in $\mathbb{F}$ with $s_0 \neq 0$.

(a) Show that the period of $< s_n >$ is less than the order of the companion matrix of the recurrence iff $s_1 s_0^{-1}$ is a root of the characteristic polynomial of the recurrence.

(b) If $\mathbb{F} = \mathbb{Z}_p$ for some prime $p \equiv \pm 3 \pmod{10}$ show that the period of any non-zero solution to the Fibonacci recurrence is the order of the companion matrix.

**Ex 8.29.** For any $m \geq 2$, let $X = \mathbb{Z}_m^2$ and $f(x_1, x_2) = (x_2, x_1 + x_2) \pmod{m}$.

(a) If $t$ is the period of the usual Fibonacci sequence $(f_j)$, prove

$$f_{t-j} = (-1)^j f_j \text{ for all } 0 \leq j \leq t.$$

(b) If the period of $(a, b) \in X$ under $f$ is odd, show that $m = 2$ must hold.

(c) Show that $(0, 0)$ is the only fixed point of $f$.

(d) Show that no orbit of $f$ has period equal to 2.

**Ex 8.30.** (a) What is the largest period for a sequence which satisfies the Fibonacci recurrence modulo 5?

(b) How many *different* sequences satisfy the Fibonacci recurrence modulo 5?

(c) Show that there is only one non-zero Fibonacci orbit modulo 3.

(d) Without calculating any sequences, prove that every Fibonacci period modulo 7 must divide 16.

(e) Find a Fibonacci period modulo 7 which equals 16.

**Ex 8.31.** (a) Factor $x^2 - x - 1$ in $\mathbb{Z}_{11}[x]$.

(b) Without calculating the actual orbit, find the period of the orbit of $(0, 1)$ under the Fibonacci recurrence modulo 11. Check your answer by calculating the orbit.

**Ex 8.32.** Find the period of 0 under $f(x) = 5x^{-1} - 2$ in $\mathbb{Z}_{11}$ (where $0^{-1}$ is defined to be 0).

**Ex 8.33.** This is a problem from the *American Mathematical Monthly*, March 1992, page 278.

(a) Given a positive integer $m$, show that the modular Fibonacci period $t(m)$ satisfies $t(m) \leq 6m$ for all $m$ and that equality holds for infinitely many $m$.

(b) Show that an analogous result holds for the **Lucas sequence** with the upper bound of $6m$ replaced by $4m$.

**Ex 8.34.** (a) Use the Fermat Test to prove that $2047 = 2^{11} - 1$ is a composite number.

(b) Show $m = 561$ is a Carmichael number.

(c) Prove: If $m$ fails the Fermat Test for $a = 2$ then $N = 2^m - 1$ fails the Strong Fermat Test for $a = 2$.

**Ex 8.35.** For this problem, estimate the probability that at least two people in a crowd of $M$ people have the same birthday. Estimate the number

of people needed to ensure this probability is greater than $1/2$. What size crowd ensures the probability to be greater than $2/3$?

**Ex 8.36.** Using an analysis similar to what we've done for the Birthday Problem, show that there exists a constant $c$ such that for sufficiently large $n$, the partial sum $\sum_{i=1}^{n} \frac{1}{i}$ is approximated by $\ln(n) + c$.

**Ex 8.37.** Let $m \geq 2$ be an integer and $f$ be a function defined on $\mathbb{Z}^k$ for some $k \geq 1$ and $s \in Z^k$. Then $< f^{(k)}(s) >$ is periodic modulo $m$ iff there exists a positive integer $n$ such that

$$< f^{(2n)}(s) > \equiv < f^{(n)}(s) > \pmod{m}.$$

**Ex 8.38.** Factor $m = 7031$ using the Pollard Rho Method with the orbit of 3 under $f(x) = x^2 + 1$.

# 9
# Computational Complexity

Analysis of algorithms is intimately related to recurrence relations. In this chapter we present many algorithms which are *recursive* in the sense that they call themselves. We'll see that the analysis of each algorithm quickly leads to a recurrence relation which we can solve using the techniques of the previous chapters. The solution of this recurrence provides information about the amount of resources used by the algorithm.

We'll also see that many easily stated unsolved problems can be found close to the edge of standard material. The analysis and improvement of basic algorithms provides a treasure chest of research problems which are fun (and maybe even profitable) to solve and are accessible to students.

An **algorithm** is a procedure which solves a problem *and* is suitable for implementation as a program on a digital computer. This informal definition makes two important points. First, an algorithm solves a problem. There are computer programs which never terminate. It would be very difficult to say that such a program does anything, let alone solves a problem. Second, each step of the algorithm should be well defined and should be representable, at least in principle, by a program. For example, $s := p/q$ is not well defined if $q$ is allowed to be zero. Also, "Find the smallest $x \in X$ for which the statement $P(x)$ is true" is not necessarily well defined. It depends on the truth value of the statement $P(x)$ and the set $X$. For instance, if $P(x)$ were true for all negative integers, then "Find the smallest $x$ for which the statement $P(x)$ is true" is not well-defined for $X = \mathbb{Z}$ but is well-defined for $X = \mathbb{N}$.

For our purposes and for many purposes, the above somewhat informal definition of algorithm is sufficient. Formal definitions of the term algorithm

were created by Turing, Markoff and others (refer to [118]). It's very satisfying that all these formal definitions of algorithm define the same class of algorithms: When $A$ is an algorithm in one of these senses, then in every one of the other senses there exists an algorithm which is equivalent to $A$.

## 9.1 Analysis of Algorithms

The **analysis of an algorithm** attempts to assay the amount of resources used by the algorithm. For any *solvable* problem there will be a limitless number of algorithms which solve the problem. How do we decide which is the best algorithm? Typical resources are time and space, and in this chapter our analyses concentrate on time analysis.

An abstract world of abstract computers and abstract programs is constructed from the real world of actual computers and actual computer programs. This construction is rarely formal because exact definitions of abstract entities are often not stated. A number of real world limitations disappear in the abstract world. For example, real computers have a fixed finite memory time and there's an upper bound on the size of numbers which can be represented. In the abstract world these limitations don't exist. There, computers are assumed to have finite but unbounded memories with no bound on the size of numbers which can be used. In practice there are examples of algorithms which work very quickly when arbitrarily large numbers are used but implementing them on real computers results in much slower algorithms. These algorithms make perfect sense in the abstract world, but have little or no relevance for the real world.

### *9.1.1 Measuring Running Time*

We want to know how much time it takes an algorithm to perform a particular task. For a real computer program this can be done using a stopwatch to time the execution of a program. This elapsed time is often called the **wall clock time**. Another way of timing a real program is to use your computer's `TIME` command so that the computer will type out the time used when the program is run. This time measure is often called **CPU time** and may differ drastically from wall clock time.

Both wall clock time and CPU time suffer from the real world problem of inexact repeatability. Two different runs of the same program may not take the same amount of time, although it is certainly possible to gather valid statistical data. A more serious problem is that both wall clock time and CPU time are highly dependent on the exact hardware and software implementation used, as well as on the input data. Specifically, if you change operating systems or run the program on a different computer or change

the input data, you will often be unable to reliably predict how long the "same" program will take to run.

We want to call one algorithm faster (it uses less time) than another algorithm if when we run the two algorithms on a computer the faster one will always finish first. To make this a fair test some variables have to be removed. For example, we'd have to code the two algorithms in the same programming language; compile the two programs using the same compiler; run the two programs under the same operating system on the same computer and not interfere with either program while it's running. In practice, even if we could control all these conditions, to our chagrin we might find that algorithm A is faster under conditions C, while algorithm B is faster under conditions D.

To avoid this unhappy situation we calculate **unitless time**. For this we find the running time $T(n)$ as a function of $n$, where $n$ is some measure of the size of the problem. For example, we could use the number of digits as the measure of problem size if the problem is addition of two integers. We could use the number of elements in a list if the problem is to sort a finite list. We could use the number of edges (or the number of vertices) in a graph if the problem is to determine whether or not a finite graph has a certain property.

We consider two algorithms to use the same time if their running times have the same order. For our purposes two running times, $T_1(n)$ and $T_2(n)$, in the variable $n$ **have the same order** when $T_1(n) = \Theta(T_2(n))$ where **Big-Theta** notation was defined in Chapter 1. The statement $T_1(n) = \Theta(T_2(n))$ means there exist positive constants $c_1, c_2$ such that for all sufficiently large $n$

$$c_1 T_1(n) \leq T_2(n) \leq c_2 T_1(n)\,.$$

In particular, we don't distinguish between algorithms whose running times are constant multiples of each other.

It's worthwhile to discuss Big-Theta notation further. Assume that $A$ is an algorithm and that the size of the input data for $A$ is represented by the variable $n$. A typical result might be that $A$ has running time $\Theta(n^2)$. Since we're using unitless time, we have no idea what this means in terms of seconds or nanoseconds. Indeed, we can think of the time unit as an unknown function of many details, among them the machine and the programming language. For example, the fact that the running time is $\Theta(n^2)$ allows us to reasonably predict that, when all these details are kept constant, doubling the size of the input will quadruple the running time of $A$.

There is even more hidden in $\Theta$-notation. Because it's asymptotic, the above prediction might not be valid unless $n$ is large enough. For example, the actual CPU time for $n = 2$ may be the same as for $n = 1$ because the output for these values is computed by a table lookup. Further, for some programs the doubling prediction might be valid for $n \geq 100$, whereas




FIGURE 9.1. The Towers of Hanoi Puzzle.

for others the prediction might be valid only for $n \geq 1024$ or some larger number.

How is the measure of running time used to compare two algorithms? Assume for the moment that algorithm $A$ has running time $\Theta(n^2)$ and that algorithm $B$ has running time $\Theta(n \log n)$. If we program these algorithms, will the program for $B$ be faster than the program for $A$? Yes, but only for large enough $n$. Depending on the constants implicit in the $\Theta$ notation, it may be that $B$ is only faster than $A$ for $n > 10^{23}$, in which case for reasonably sized data the "slower" A might actually be faster.

To summarize, if we find that the order of the running time of algorithm A is strictly less than that of algorithm B, then we can be confident that for any large enough problem A will run faster than B regardless of the input. On the other hand, if the running times of A and B have the same order, then we won't be able to predict which one will be faster for a given input.

### *9.1.2 The Towers of Hanoi: An Example*

In this section we illustrate the above discussion by looking at a concrete example, the running time of an algorithm for solving the **Towers of Hanoi puzzle**.

Ball's *Mathematical Recreations and Essays* [5] contains one of the first mathematical formulations of the Towers of Hanoi puzzle. The puzzle consists of three towers (pegs) usually called A,B,C and $n$ disks of different sizes (numbered 1 through $n$) such that the $i^{\text{th}}$ disk is larger than the $j^{\text{th}}$ disk whenever $i > j$. Initially, the disks are stacked on Tower A in order of size, with the largest disk on the bottom and the smallest on top. The problem is to move the stack of disks from Tower A to Tower C, moving the disks one at a time in such a way that a larger disk is never stacked above a smaller disk. (Refer to Figure 9.1.) An extra constraint is the sequence of moves should be as short as possible. An algorithm is therefore said to *solve the Towers of Hanoi problem* if when the number of disks and the names of the three towers are inputted the algorithm returns a sequence of moves which conforms to the above rules.

Three simple observations form the key to the problem. The first is that moving the largest disk requires all other disks to be out of the way, that

is, the other $n-1$ smaller disks must be located on another tower. In order for the final result to be stacked on Tower C, Tower C must be free for the largest disk, that is, the $n-1$ disks have been moved from Tower A to Tower B, which is the Towers of Hanoi puzzle for $n-1$ disks. After the largest disk is moved to Tower C, the other $n-1$ disks must be moved from Tower B to Tower C, which is again the Towers of Hanoi puzzle for $n-1$ disks. This recursive procedure leads to the following recursive algorithm:

```
Procedure HANOI(A, B, C, n)
If n = 1 then move the top disk from A to C
         else HANOI(A, C, B, n − 1)
              move the top disk from A to C
              HANOI(B, A, C, n − 1).
```

This is called a recursive procedure because it calls itself. Also, unlike the worm Ouroboros it doesn't endlessly swallow its own tail. Each time it calls itself it decrements the size parameter $n$ by 1 which means that eventually the sequence of calls "bottoms out" with a call to `HANOI` with $n = 1$. This call makes one move and then returns to the previous call. The operation of this algorithm can be seen in Figure 9.2, where we give the sequence of calls and the states of the puzzle for $n = 3$.

There's one point in the trace which might seem at least slightly strange: How did the `Move from A to C` in the algorithm give `Move from B to C` in the trace? The answer is that "$A$" in the algorithm refers to the first parameter in the call, and "$C$" refers to the third parameter in the call. So when `Move from A to C` is referred to within a call to `HANOI(C,A,B,n)`, it makes a move from $C$ to $B$ in the execution of the solution of the puzzle.

Now that we have an algorithm for solving the Towers of Hanoi puzzle, let's analyze its running time. Because we're calculating *unitless* time, we don't have to know how long it takes to perform the various operations such as moving a disk or issuing a procedure call. We do have to distinguish between operations which take a constant amount of time (that is, time independent of the value of $n$) and operations whose running time depends on $n$. Aho, Hopcroft, and Ullman in [2] call the assumption that each operation takes the same amount of time the **uniform cost criterion**. Under this condition the running time $T(n)$ used for $n$ disks satisfies the nonhomogeneous first-order recurrence

$$T(n) = 2T(n-1) + c \quad \text{for all } n > 1\,, \tag{9.1}$$

where $c$ is a positive constant. This is true because there are two calls to the procedure with $n-1$ disks and the constant $c$ is the sum of the constant running times for the various operations. From the work in Chapter 3 we




FIGURE 9.2. Trace of the program `HANOI` for $n = 3$.

know (9.1) has the solution

$$T(n) = T(1)2^{n-1} + c(2^{n-1} - 1)\,,$$

and

$$\frac{T(1)}{2}\,2^n \le T(n) \le \frac{T(1)+c}{2}\,2^n$$

which gives $T(n) = \Theta(2^n)$

Instead of assuming the uniform cost criterion, we could have assumed that the running time is proportional to the number of moves. Under that assumption, our argument could proceed as follows: Let $M(n)$ be the number of moves performed when the algorithm is called with $n$ disks. Then

$$M(n) = 2M(n-1) + 1 \text{ for all } n > 1\,,$$

with initial condition $M(1) = 1$ which has the solution $M(n) = 2^n - 1$. Because the running time $T(n)$ is proportional to $M(n)$, then $T(n) = \Theta(M(n)) = \Theta(2^n)$, which is the same as we obtained above.

In our analysis of the algorithm `HANOI` we've made several other assumptions which bear further examination. In any actual implementation on a real digital computer, the number $n$ needs to be stored and manipulated internally. The time required for this is at the very least dependent upon the number of computer words required to represent the number $n$. However, our argument assumed that each operation takes a constant time, independent of $n$. In addition, representing a state of the puzzle in a computer's memory requires increasing memory as $n$ increases and, presumably, increasing time to manipulate this memory. All of the operations in `HANOI` depend on the space needed to store $n$ in memory and this is proportional to the number of bits needed to represent $n$, which is $\lceil \log_2 n \rceil$. We can ignore the ceiling function and consider space to be proportional to $\log n$ where the constant of proportionality depends on the word size of the computer. Aho, Hopcroft, and Ullman [2] call this the **logarithmic cost criterion** and suggest it should be used when the numbers in the algorithm don't have fixed bounds. Using this criterion, the running time for the $n$-tower puzzle is

$$T(n) = 2T(n-1) + c\log(n)\,,$$

with initial condition $T(1) = t_1$. (As above, $c$ and $t_1$ are unknown positive constants.) Again referring to Chapter 3, the solution to this recurrence relation is

$$T(n) = 2^n \left[ \frac{t_1}{2} + c \sum_{i=1}^{n} \frac{\log i}{2^i} \right]\,,$$

where each summand is positive. Therefore, the sum is less than the infinite sum,

$$0 < \sum_{i=1}^{n} \frac{\log i}{2^i} \le \sum_{i=1}^{\infty} \frac{\log i}{2^i}$$

which converges by the Ratio Test. So, $T(n) = \Theta(2^n)$ since this sum is bounded above and below by positive constants and

$$c_1 2^n \leq T(n) \leq c_2 2^n \,,$$

where for instance $c_1, c_2$ can be chosen as $c_1 = \frac{t_1}{2}$ and $c_2 = \frac{t_1}{2} + c\sum_{i=1}^{\infty} \frac{\log i}{2^i}$. The increase in operation time due to the size of $n$ therefore has no effect on the order of our asymptotic estimate of the running time because any changes were absorbed into the "implied constant" in $\Theta(2^n)$.

If we consider building a physical Towers of Hanoi puzzle and the time to move disk $n$ grows as some function $g(n)$, we would have

$$T(n) = 2T(n-1) + g(n)\,.$$

Provided the eminently reasonable assumption that $g(n+1) < 2g(n)$ holds for all sufficiently large $n$, a modification of the above argument would again yield $T(n) = \Theta(2^n)$. From this we see that the conclusion that `HANOI` has $\Theta(2^n)$ running time is **robust**: Specific details of the implementation of the algorithm has no effect on the time. Table 9.1 gives some actual running times and compares them to the predicted running time $c2^n$ where $c$ is computed as $c = T(10)/2^{10}$. Notice that the ratio $T(n)/T(n-1)$ is approximately 2, as predicted by $T(n) = \Theta(2^n)$.

TABLE 9.1. Run times for the `HANOI` Algorithm showing the predicted $\Theta(2^n)$ behavior.

| Run Times for the `HANOI` Algorithm | | | | |
|---|---|---|---|---|
| n | Number of Moves | Actual Time | Predicted Time | T(n)/T(n-1) |
| 5 | 31 | 10 | 74 | |
| 6 | 63 | 30 | 147 | 3 |
| 7 | 127 | 230 | 294 | 7.7 |
| 8 | 255 | 250 | 589 | 1.1 |
| 9 | 511 | 1287 | 1178 | 5.1 |
| 10 | 1023 | 1810 | 2355 | 1.4 |
| 11 | 2047 | 4270 | 4708 | 2.4 |
| 12 | 4095 | 10471 | 9421 | 2.5 |
| 13 | 8191 | 19398 | 18842 | 1.9 |
| 14 | 16383 | 39447 | 37683 | 2.0 |
| 15 | 32767 | 77669 | 75364 | 2.0 |
| 16 | 65535 | 147832 | 150733 | 1.9 |
| 17 | 131071 | 301652 | 301466 | 2.0 |

We close this section with a problem from the Advanced Problem Section of the June-July 1939 issue of the *American Mathematical Monthly* (page

363) in which B.M. Stewart proposed a generalization of the Towers of Hanoi puzzle to $k \geq 3$ towers.

> Given a block in which are fixed $k$ pegs and a set of $n$ washers, no two alike in size, and arranged on one peg so that no washer is above a smaller washer. What is the minimum numer of moves in which the $n$ washers can be placed on another peg, if the washers must be moved one at a time, subject always to the condition that no washer be placed above a smaller washer?

Two years later, two solutions to the problem were published in the Advanced Problems Section of the March 1941 issue of the same journal (pages 216-217). These solutions yield algorithms for solving the Towers of Hanoi puzzle for $k$ towers. Many people believe that these are optimal, but a proof of optimality is still an open question after 60 years. We close this section with a comment from a March 2002 interview with Donald Knuth [76, p. 321]. "In the case of the 4-peg 'Tower of Hanoi', there are many, many ways to achieve what we think is the minimum number of moves, but we have no good way to characterize all solutions. So that's why I personally came to the conclusion that I was never going to solve it, and I stopped working on it in 1972. But I spent a solid week working on it pretty hard."

## 9.2 Computer Arithmetic

To store an integer $N$ in a digital computer we use the binary representation of $N$, and the number of bits for this representation of $N$ is $n = \lceil \log_2(N) \rceil$. This is then broken into blocks which fit into the internal words of the computer's memory and are stored as some sort of list. Every machine has a limit on the size of integers which can be represented in this fashion. In many classical integer data types this was limited to one or two machine words. In contrast, many present day applications require no pre-set bounds on the size of integers.

In any storage scheme there are details which are specific to the machine and the operating system. Among these are the word size and the manner in which the list is stored. In analysis of algorithms these details are avoided by considering the number of bit operations required to perform a calculation. A bit operation is either a **unary operation** (an operation performed on one bit such as reversing a single bit) or a **binary operation** (an operation on two bits such as addition). We assume that the running time is proportional to the number of bit operations. In order for this to make sense, we're assuming that operations on words introduce a constant multiple to the running time, and also that our time estimates are not dominated by the overhead required for keeping track of the list structure.

### *9.2.1 Addition and subtraction*

How long does it take to add two $n$-bit natural numbers $M$ and $N$? That is, how many bit operations are required to add the integers? For this consider the algorithm we all use for adding two binary representations by hand. The maximal number of bit operations for the sum is $n$ additions with carries. If we write the carry bits in a row above $M$ and $N$, this row will always have a zero in the right-most bit and will extend at most one place to the left of $N$. Taking this into account at most $2n$ bit operations will be used to perform the addition.

On the other hand, it takes at least $n$ bit operations to write down the number $M + N$. This is true even in the extreme case when one of the summands is zero and we know a priori that the sum is just equal to the other integer. Because of this, $n$ is a lower bound on the number of bit operations needed to add $M$ and $N$. Since $M - N = M + (-N)$, the cost of subtracting $N$ from $M$ equals the cost of adding $N$ to $M$ plus the cost of negating $N$. Since negation can be accomplished in $n$ bit operations, we conclude that the time required to add or subtract two $n$-bit integers is $\Theta(n)$.

### *9.2.2 Multiplication and division*

The standard algorithm used for hand computation of the product $M * N$ in binary involves multiplying $M$ by each bit of $N$ (resulting either in 0 or $M$ in binary), stacking up the results with appropriate left shifts, and then adding entries in the stack. For example, to compute $7 * 5 = 35$ the standard algorithm is

$$\begin{array}{cccccc} & & & 1 & 1 & 1 \\ & & * & 1 & 0 & 1 \\ \hline & & & 1 & 1 & 1 \\ & & 0 & 0 & 0 & \\ & 1 & 1 & 1 & & \\ \hline 1 & 0 & 0 & 0 & 1 & 1 \end{array} .$$

When $M$ and $N$ are $n$-bit integers, this algorithm requires $n$ additions of integers which have at most $2n$-bits and the complexity of this standard algorithm is $\Theta(n^2)$.

This is not the best algorithm for solving this problem, since there's an easily implemented multiplication algorithm which has complexity $\Theta(n^{\log_2(3)})$ (refer to Exercise 9.17). There's also another algorithm, due to Schönhage and Strassen [126], which has asymptotic complexity $O(n \log n \log(\log n))$ and therefore its running time has order only slightly larger than the order of addition. However, because the constant involved in the asymptotic notation is large the Schönhage-Strassen algorithm isn't better than the more straightforward algorithms until $n$ is quite large.

What about the cost of division? Later in this chapter we use Newton's method to design an algorithm for division which has the same running time as multiplication.

## 9.3 An Introduction to Divide-and-Conquer

Historically the idea of divide-and-conquer is usually attributed to Julius Caesar [151]. In the context of problem solving, divide-and-conquer may reasonably be attributed to René Descartes [41, 42]. This is the same Descartes whose Rule of Signs was used in Chapter 5 to prove every non-negative polynomial has only one positive root. The keystone of Descartes' analytic method is first to break a complicated problem into easier constituent parts and then to solve the individual parts. An understanding of the complicated problem is then obtained from the solution of its parts. In *How to Solve It* [112] George Polya stressed that breaking down a problem into several smaller problems of the same kind is a typical step in solving a mathematical problem, and he further pointed out that this sort of analysis would easily lead to an inductive proof of the correctness of the solution. In the 1930's, philosophers such as Gödel, Kleene, and Ackerman recognized the central role of this recursive technique and their theoretical analyses led Turing to describe an abstract digital computer. After these **Turing machines** were eventually embodied as physical digital computers, the problem of programming these computers led back to the divide-and-conquer technique (for example, refer to [150]).

The term **divide-and-conquer algorithm** is now usually reserved for the algorithm design strategy which solves a problem of size $n$ by solving several problems of size $n/c$ for some constant $c$ so that in its current usage the "divide" means that there is an actual division of the problem size. The `HANOI` algorithm which we discussed in Section 9.1.2 is not a divide-and-conquer algorithm in this sense because it uses subproblems of size $n-1$. (Perhaps a better term for this type of algorithm would be *subtract-and-conquer*?) Many commonly used divide-and-conquer algorithms split the problem into two subproblems, each having half the size of the original problem. Because such an algorithm cannot keep dividing indefinitely, we specify a size limit above which the problem is divided and below which the algorithm calls another algorithm which solves the small size instances. Then those solutions are combined to give a solution to the original problem.

The recursive structure of a divide-and-conquer algorithm leads directly to an inductive proof of correctness and a recurrence for the running time of the algorithm.

### *9.3.1 Example: Polynomial multiplication*

Let's look at an example which makes this discussion more concrete. Our example is **polynomial multiplication** (or **convolution**), where the output is the one polynomial which is the product of two input polynomials. We assume an **arithmetic model of computing** which means both that our computer can perform arithmetic operations and that we want an algorithm which multiplies the two input polynomials using only arithmetic operations on the coefficients. As above, the running time for the algorithm is the number of arithmetic operations used in the algorithm.

Assume each input polynomial has degree $n-1$ (and hence $n$ coefficients which we record in a vector of size $n$). The output polynomial then has degree $2(n-1)$, recorded as a vector with $2n-1$ components. The standard algorithm for multiplying polynomials is similar to the standard algorithm for multiplying integers. If $P(x)$ and $Q(x)$ are the input polynomials, multiplication proceeds by multiplying $P(x)$ by the constant coefficient of $Q(x)$, shifting one space to the left, multiplying $P(x)$ by the linear coefficient of $Q(x)$, and so on. This results in $n$ vectors which are added componentwise to get the final result. Here's an example which multiplies two quadratic polynomials in this fashion:

$$
\begin{array}{cccccccccc}
 & & & & 2x^2 & + & 3x & - & 7 \\
 & & & & 3x^2 & - & 2x & + & 2 \\
\hline
 & & & & 4x^2 & + & 6x & - & 14 \\
 & - & 4x^3 & - & 6x^2 & + & 14x & & \\
6x^4 & + & 9x^3 & - & 21x^2 & & & & \\
\hline
6x^4 & + & 5x^3 & - & 23x^2 & + & 20x & - & 14
\end{array}
.
$$

This method for multiplying two polynomials has running time $\Theta(n^2)$ because each of the $n^2$ table entries is a product of two coefficients and each table entry occurs once as a summand.

```
Procedure
        For i = 0 TO n - 1 DO
            For j = 0 TO n - 1 DO
                c_{i+j} = c_{i+j} + a_i * b_j
```

AN ALGORITHM TO MULTIPLY TWO POLYNOMIALS

Let's construct a divide-and-conquer algorithm for this problem. The basic philosophy of a divide-and-conquer approach to polynomial multiplication is to think of the input polynomial as being constructed from several smaller polynomials. In practice we divide each polynomial "in half". For

simplicity, we first consider polynomials with an even number of coefficients, say $n = 2m$. Then we can write

$$P(x) = a_{2m-1}x^{2m-1} + \ldots + a_1 x + a_0 = x^m P_1(x) + P_0(x)\,,$$

where each of the polynomials $P_1(x) = a_{2m-1}x^{m-1} + \ldots + a_{m+1}x + a_m$ and $P_0(x) = a_{m-1}x^{m-1} + \ldots + a_1 x + a_0$ has $m = n/2$ coefficients. (For example, if $P(x) = 5x^3 + 12x^2 + 7x + 8$ then $P_1(x) = 5x + 12$ and $P_0(x) = 7x + 8$.) Splitting our two input polynomials $P(x)$ and $Q(x)$ of degree $2m$ in this way and suppressing the argument variable $x$ results in

$$PQ = (x^m P_1 + P_0)(x^m Q_1 + Q_0)\,,$$

giving

$$PQ = x^n P_1 Q_1 + x^m [P_1 Q_0 + P_0 Q_1] + P_0 Q_0\,. \tag{9.2}$$

Since multiplying a polynomial by an integer power of $x$ simply shifts the sequence of coefficients, the original problem of finding the product $PQ$ has been reduced to the four subproblems of finding the products $P_1Q_1$, $P_1Q_0$, $P_0Q_1$, and $P_0Q_0$.

Because $m$ (and also the original $n$) might be odd, we need to modify this procedure to get a divide-and-conquer algorithm which works for any pair of polynomials but only a slight modification is needed. For this, let $n-1$ equal the maximal degree of the factors $P(x), Q(x)$ and define $m \geq 1$ to be the least exponent such that $2^m \geq n$, that is, $m = \lceil \log_2(n) \rceil$. Filling in zero entries where necessary, we treat $P(x)$ and $Q(x)$ as polynomials with $2^m$ coefficients. Dividing each of $P(x)$ and $Q(x)$ into their polynomial parts, we obtain four subpolynomials having $2^{m-1}$ coefficients. We continue this subdivision process for $m$ stages, which in general is for as long as we have more than one coefficient. When we reach the stage at which each subpolynomial has one coefficient, we multiply the pairs of constants that remain. Equation (9.2) is then used to retrace the steps and arrive at the product polynomial.

The running time analysis for this divide-and-conquer algorithm is fairly simple. From the first division we obtain four subproblems, each a multiplication of polynomials of degree $2^{m-1}$. The multiplications by $x^{2^m}$ and $x^{2^{m-1}}$ are shifts in the coefficient vectors and are carried out before the vectors are added. Therefore, if $T(n)$ denotes the time needed to multiply two polynomials with $n$ coefficients (where we consider $n = 2^m$), then

$$T(n) = 4T(n/2) + bn\,, \tag{9.3}$$

where $4T(n/2)$ represents the time needed to multiply the four half-size polynomials and the $bn$ term comes from the time needed to shift and add the results. If we were doing an exact operation count we'd write down an explicit value for $b$. However, since we don't know the actual time required

for coefficient addition or multiplication we simply assume that $b$ is some positive constant.

Notice that the recurrence (9.3) is nonlinear since $T(n)$ does not depend on a constant number of preceding terms. Despite this novelty, it's possible to convert (9.3) to a linear recurrence which models the situation. For this, we exploit the fact that $n$ is assumed to be a power of 2. Introducing the new variable $t_k = T(2^k)$, (9.3) becomes

$$t_k = 4t_{k-1} + b2^k \text{ with } t_0 = T(1)\,,$$

which is a nonnegative recurrence with dominant eigenvalue $\lambda_0 = 4$. It has the form

$$t_k = 4t_{k-1} + b\lambda_1^k \text{ where } \lambda_1 < \lambda_0\,,$$

and Theorem 5.3.1 implies that $t_k = \Theta(4^k)$, giving $T(n) = \Theta(n^2)$ when $n$ is a power of two. What about other $n$? Even when $n$ is not a power of two, we've already established that we write the factors as polynomials with $2^m$ coefficients where $m = \lceil log_2(n) \rceil$, and (9.3) becomes

$$4T(n/2) + bn < T(n) = 4T(n/2) + b(cn) < 4T(n/2) + b(2n)\,,$$

since $n = c \cdot 2^m$ for some $1 < c < 2$. This at most quadruples the implied constant in $\Theta(n^2)$ and we've proved that the running time for this divide-and-conquer algorithm is $\Theta(n^2)$.

What's the point of this algorithm when we already have an easy to understand iterative algorithm with running time $\Theta(n^2)$? A further examination of this algorithm allows us to a make only a slight modification which speeds things up.

Returning to the formula in (9.2) we notice that for two of the half-size products, $P_1Q_0$ and $P_0Q_1$, only their sum and not the separate products is needed. This is important, because there's a way to calculate their sum without calculating both products! The trick is to use subtraction and the polynomial identity

$$P_1Q_0 + P_0Q_1 = (P_1 + P_0)(Q_1 + Q_0) - P_1Q_1 - P_0Q_0\,. \tag{9.4}$$

Since each of $P_1$ and $P_0$ has $n/2$ coefficients, so does their sum $P_1 + P_0$. Similarly $Q_1 + Q_0$ has $n/2$ coefficients, and their product can be computed with two half-size polynomial additions and one half-size polynomial multiplication. Since we also need both of $P_1Q_1$,$P_0Q_0$, the product can be computed with three multiplications, four additions, and two subtractions of half-size polynomials. That is, we've replaced one of the multiplications in the earlier algorithm with four additions and two subtractions. This would be a loss if these arithmetic operations all had the same cost. But these operations are *polynomial* operations, and the additions and subtractions cost $\Theta(n)$ and multiplications cost $\Theta(n^2)$. Therefore, the gain is enormous and we obtain the recurrence

$$T(n) = 3T(n/2) + Bn\,, \tag{9.5}$$

where the constant $B$ in this equation is slightly larger than the constant $b$ in the previous recurrence (9.3). It's easy to check that

$$T(n) = an^{\log_2(3)} - 2Bn$$

satisfies this recurrence for some constant $a$ which depends on the initial condition. Therefore, $T(n) = \Theta(n^{\log_2(3)})$, where $\log_2(3) \approx 1.5849$, and this new algorithm is faster than $\Theta(n^2)$ for large enough $n$. In Exercise 9.17 you see that this algorithm can be used to obtain a $\Theta(n^{\log_2(3)})$ multiplication algorithm for integers.

What about the implied constant in this algorithm, as compared to the one in the standard algorithm? Table 9.2 records a comparison of the running times for the two algorithms for various pairs of polynomials with $n$ coefficients. An inspection of the table shows that the asymptotic estimates take over quite quickly. In these examples we were careful to use polynomials with small integer coefficients so that the efficiency of either algorithm is unaffected by the size of the coefficients.

TABLE 9.2. Comparison of the runtimes for two polynomial multiplication algorithms. The iterative algorithm is the classical method with run time $\Theta(n^2)$. The recursive algorithm is the 3 half-size method with runtime $\Theta(n^{\log_2 3})$. The ratios of T(2n)/T(n) are very close to the predicted values of 4 and 3. The "faster" catches up to the "slower" method for large $n$.

| Run Times for $n$ coefficient Polynomial Multiplication. | | | | |
|---|---|---|---|---|
| n | Iterative Time (ms) | | n | Recursive Time (ms) |
| 2 | 0 | | 2 | 0 |
| 4 | 0 | | 4 | 0 |
| 8 | 0 | | 8 | 0 |
| 16 | 0 | | 16 | 0 |
| 32 | 0 | | 32 | 15 |
| 64 | 0 | | 64 | 31 |
| 128 | 15 | | 128 | 78 |
| 256 | 31 | | 256 | 218 |
| 512 | 140 | | 512 | 625 |
| 1024 | 500 | | 1024 | 1875 |
| 2048 | 2000 | | 2048 | 5578 |
| 4096 | 8000 | | 4096 | 16812 |
| 8192 | 32000 | | 8192 | 50438 |
| 16384 | 128094 | | 16384 | 151578 |
| 32768 | 512376 | | 32768 | 454734 |

## 9.4 Simple Divide-and-Conquer Algorithms

Each of the two divide-and-conquer algorithms examined in the previous section led to a recurrence relation for its running time and there was a strong similarity between the two arguments. In this section we generalize these arguments and derive a recurrence relation for the running time of a whole class of divide-and-conquer algorithms. We will give examples of algorithms which illustrate each of the three types of solutions obtained.

In our two previous examples, the problem of polynomial multiplication was split into two subproblems of the same type which were half the original size. Here we generalize this slightly and assume that a problem of size $n$ is split into several (say $a$) subproblems, each of size $n/c$ for some constant $c$. For many divide-and-conquer algorithms $c = 2$ holds, but the same analysis works for general $c$.

In some cases, the splitting process is easy and has negligible computational cost, but we want to allow for the possibility that splitting takes some time. Usually the time-consuming part of a divide-and-conquer algorithm is the combining step when the answers to subproblems are used to compute the answer to the whole problem. The general form of a divide-and-conquer algorithm is

```
Procedure DandC(DATA,n,SOLUTION)
IF n is small
     THEN Solve by some special algorithm
     ELSE SPLIT(DATA,n) into (D_1,n/c)
          DandC(D_1, n/c, S_1)
          D_2 := f_2(DATA, D_1, S_1)
          DandC(D_2, n/c, S_2)
          D_3 := f_3(DATA, D_1, S_1, D_2, S_2)
           ⋮
          D_a := f_a(DATA, D_1, S_1, D_2, S_2, ..., D_{a-1}, S_{a-1})
          DandC(D_a, n/c, S_a)
          SOLUTION:=f_{a+1}(DATA, D_1, S_1, D_2, S_2, ..., D_a, S_a)
          RETURN(SOLUTION)
```

In this general form we've allowed for the results of previous subproblems to be processed and then fed into the next subproblem. The function $f_{a+1}$ performs the final combination of results to produce the solution.

An efficient algorithm should have a running time which is bounded by a function which is *polynomial* in the size of the input or perhaps the size of the output. For this reason we assume that the time for the splitting and combining together is given by a polynomial in $n$. Since we're only inter-

ested in the order of the running time, we can concentrate on the largest term of the polynomial and assume that the running time for the split and combine steps is $bn^m$ for some positive constant $b$ and some constant exponent $m$. For this to be a polynomial, $m$ must be an integer but it costs nothing more to allow $m$ to be any nonnegative real number.

Under these assumptions the running time for a divide-and-conquer algorithm which splits a problem of size $n$ into $a$ subproblems of size $n/c$ is given by the recurrence

$$T(n) = aT(n/c) + bn^m \tag{9.6}$$

(compare this with (9.3) and (9.5)) where the initial condition is specified by the algorithm that handles problems of small size. Since we only run the small size algorithm on data of bounded size, we can bound its running time by some positive constant. This allows us to assume that the initial condition is $T(1)$ and is positive but otherwise unknown.

As in the last section, we essentially use logarithms to the base $c$ by setting $n = c^k$ and $t_k = T(c^k)$. Then (9.6) yields

$$t_k = at_{k-1} + bc^{km} \quad \text{with } t_0 > 0\,.$$

This is a nonnegative difference equation with $\lambda_0 = a$ and Theorem 5.3.1 says that the relative sizes of $a$ and $c^k$ yield three different orders of magnitude for the sequence $< t_k >$,

$$t_k = \begin{cases} \Theta(c^{km}) & \text{for } a < c^m \\ \Theta(kc^{km}) & \text{for } a = c^m \\ \Theta(a^k) & \text{for } a > c^m \end{cases}.$$

In order to translate this back into the running time $T(n)$ we recall $n = c^k$ and $t_k = T(c^k)$ to get

$$T(n) = \begin{cases} \Theta(n^m) & \text{for } a < c^m \\ \Theta(\log_c(n) n^m) & \text{for } a = c^m \\ \Theta(n^{\log_c(a)}) & \text{for } a > c^m \end{cases}.$$

(The last line follows from the observation that $a^k = c^{\log_c(a^k)} = c^{k \log_c(a)} = n^{\log_c(a)}$.) Therefore, there are three different types of behavior that can occur for the running time of a divide-and-conquer recurrence.

**Running Time for Divide-and-Conquer Algorithms**

**Theorem 9.4.1.** *Suppose that a divide-and-conquer algorithm splits a problem of size $n$ into $a$ subproblems, each of size $n/c$, and that the sum of the running times for* `SPLIT` *and* `COMBINE` *is a polynomial $p_m(n)$ of degree $m$. Assuming $T(1)$ is positive, the running time for the divide-and-conquer algorithm satisfies*

$$T(n) = aT(n/c) + p_m(n)\,,$$

*and has order*

$$T(n) = \begin{cases} \Theta(n^m) & \text{if } a < c^m \\ \Theta(n^m \log(n)) & \text{if } a = c^m \\ \Theta(n^{\log_c a}) & \text{if } a > c^m \end{cases}.$$

### *9.4.1 Example 1: A return to polynomial multiplication*

In the two divide-and-conquer algorithms for polynomial multiplication, $c$ was equal to 2 because each subproblem has half the size of the original problem. Since the split and combine operations use only addition and subtraction of polynomials, their combined running time is a linear polynomial and $c^m = 2$. The more straightforward algorithm has $a = 4 > c^m$ while the improved algorithm using (9.4) has $a = 3 > c^m$. The third case of the theorem applies for both. As we saw earlier, the algorithm with $a = 3$ will eventually be faster than that with $a = 4$ since the smaller value of $a = 3$ reduces the *exponent* in the running time from $\Theta(n^2)$ to $\Theta(n^{\log_2(3)})$. If we could somehow further reduce the number of subproblems to two (with $a = 2$) we would move to the second case and then have an asymptotic running time of $\Theta(n \log n)$. No known divide-and-conquer algorithm for polynomial multiplication has $a = 2$, but in Section 9.5 we'll see that a completely different type of algorithm for polynomial multiplication does achieve this complexity of $\Theta(n \log n)$.

### *9.4.2 Example 2: Matrix multiplication*

Another problem in which divide-and-conquer leads to a fast algorithm is matrix multiplication. If two $n \times n$ matrices are broken into four $n/2 \times n/2$ matrices, we then compute the matrix product $C$ using

$$\begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \begin{bmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{bmatrix} = \begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix}.$$

The straightforward algorithm is

$$C_{11} = A_{11}B_{11} + A_{12}B_{21}$$
$$C_{12} = A_{11}B_{12} + A_{12}B_{22}$$
$$C_{21} = A_{21}B_{11} + A_{22}B_{21}$$
$$C_{22} = A_{21}B_{12} + A_{22}B_{22}$$

with eight half-size multiplications. Because the additions can be done in time proportional to $n^2$, the equation for the running time of this divide-and-conquer algorithm is

$$T(n) = 8T(n/2) + bn^2 .$$

This equation has $a = 8$, $c = 2$, $m = 2$, giving $a > c^m$ and $T(n) = \Theta(n^{\log_2 8}) = \Theta(n^3)$ which is the same as the order of magnitude of the running time for the standard "row times column" algorithm for matrix multiplication.

A faster divide-and-conquer algorithm was designed by Strassen [137], and uses seven half-size multiplications. **Strassen's algorithm** computes the eight products

$$M_1 = (A_{12} - A_{22})(B_{21} + B_{22}) \quad ; \quad M_2 = (A_{11} + A_{22})(B_{11} + B_{22})$$

$$M_3 = (A_{11} - A_{21})(B_{11} + B_{12}) \quad ; \quad M_4 = (A_{11} + A_{12})B_{22}$$

$$M_5 = A_{11}(B_{12} - B_{22}) \quad ; \quad M_6 = A_{22}(B_{21} - B_{11})$$

$$M_7 = (A_{21} + A_{22})B_{11}$$

and uses them in

$$C_{11} = M_1 + M_2 - M_4 + M_6 \quad ; \quad C_{12} = M_4 + M_5$$

$$C_{21} = M_6 + M_7 \quad ; \quad C_{22} = M_2 - M_3 + M_5 - M_7 .$$

It is a simple exercise in algebra to prove this algorithm is correct– the real difficulty was discovering the algorithm in the first place. The running time of Strassen's algorithm obeys the equation

$$T(n) = 7T(n/2) + bn^2$$

because addition and subtraction of matrices can be carried out in time proportional to $n^2$, and $T(n) = \Theta(n^{\log_2 7})$ .

### 9.4.3 Example 3: MERGESORT

A concrete example of the second type of the divide-and-conquer behavior is given by the MERGESORT algorithm. For this algorithm the input is a one-dimensional array with $n$ entries and the output is an array with the entries rearranged into increasing order. A divide-and-conquer approach is used, with the array divided into two subarrays that are individually sorted and then combined to give the sorted array. The following algorithm captures this approach.

```
Procedure MERGESORT(A, n)
IF n > 1
  THEN Split A into two arrays of size n/2, A_1 and A_2
       MERGESORT(A_1)
       MERGESORT(A_2)
       MERGE(A_1, A_2)
```

When $n = 1$ is reached there is only one item in the array and so no sorting needs to be done. All of the real work gets done in the MERGE algorithm which combines two sorted arrays $A_1$ and $A_2$ into one sorted array $A$. Without going into the details, we'll say that MERGE can be accomplished by making $n - 1$ comparisons of elements and $n$ moves. This yields the recurrence

$$T(n) = 2T(n/2) + bn\,,$$

for some constant $b$. Since $a = 2$ and $c^m = 2$, we are in the second case and $T(n) = \Theta(n \log n)$.

### 9.4.4 Example 4: Applications of Newton's method

As an example of the first type of behavior we use Newton's method to estimate the values of the reciprocal function $g(x) = 1/x$ and the square root function $g(x) = \sqrt{x}$. These are special cases of estimating functions $g(x)$ at values $x = A$ when $g(x)$ is a one-to-one function in some neighborhood of $A$ where the assumption that $g(x)$ is one-to-one ensures that it has a left inverse $f(x)$ with $f(g(x)) = x$ in a neighborhood of $x = A$. We further assume the left inverse is one-to-one and twice differentiable in this neighborhood. These conditions hold for each of $g(x) = \sqrt{x}$ and $g(x) = 1/x$ on $x > 0$. Since $A = f(g(A))$, any approximation of the solution $f(x) - A = 0$ is close to $g(A)$.

We've already used Newton's method to locate the zeroes of a polynomial. The method uses the iteration

$$N(x) = x - \frac{f(x) - A}{f'(x)}$$

to estimate a root of $f(x) = A$ where $f$ is a twice differentiable function. For this to be computationally feasible, the right-hand side should be in a form that's easy to compute which for us means the right-hand side can be computed in polynomial time $p(n)$ for size $n$ input. Under reasonable hypotheses (refer to [18, 136]), when a good initial approximation is used each iteration of Newton's method doubles the number of correct digits. For instance, when the approximation $x$ agrees with the root on the first five bits then $N(x)$ agrees on about the first ten bits and applying the iteration another time, $N^{(2)}(x)$ agrees on twenty bits, and so forth. Therefore, for input of size $n$ the running time $T(n)$ satisfies the divide-and-conquer recurrence

$$T(n) = T(n/2) + p(n)$$

where the polynomial $p(n)$ records the difficulty of computing $N(x)$ for the particular function under consideration. Newton's method is thus a divide-and-conquer algorithm provided $N(x)$ can be computed from $x$ in polynomial time.

Consider the reciprocal function $g(x) = 1/x$ on $x > 0$, a function which is its own inverse. Since $f'(x) = -1/x^2$, Newton's formula for the iteration is

$$N(x) = x - \frac{1/x - A}{-1/x^2} = x(2 - Ax)\,.$$

In this form, no division is used and only two multiplications and one subtraction of $n$-bit numbers are required which can be accomplished in quadratic time and the running time satisfies

$$T(n) = T(n/2) + p_2(n)\,, \tag{9.7}$$

for some quadratic polynomial $p_2(n)$. Here $a = 1$ and $c^m = 2^2 = 4$, and we're in the first case, with running time $T(n) = \Theta(n^2)$. Since division of real numbers can be performed using the algorithm implicit in $\frac{a}{b} = a * \frac{1}{b}$, this also yields an order $\Theta(n^2)$ algorithm for division.

Finding square roots can also be performed in $\Theta(n^2)$ time using Newton's method. The one-to-one function $g(x) = \sqrt{x}$ has differentiable inverse function $f(x) = x^2$ which is one-to-one on $x > 0$. Since $f'(x) = 2x$, the Newton iteration formula is

$$N(x) = x - \frac{x^2 - A}{2x} = \frac{x}{2} + \frac{A/2}{x}\,.$$

where $A/2$ need only be computed at the outset and not at each iteration. For this calculation it's helpful to represent numbers in binary notation,

because $A/2$ is just a binary shift of $A$ and is computationally trivial. The complicated part is the division by $x$, for which we use the quadratic time algorithm above. Therefore, the running time again satisfies (9.7) for some quadratic polynomial $p_2(x)$ and $T(n) = \Theta(n^2)$.

It's worth noting that this discussion shows that both division and taking square roots can be computed in the same order of time as multiplication. Although we've only demonstrated this for the standard $\Theta(n^2)$ multiplication algorithm, the same argument shows that any multiplication algorithm whose order is at least $\Theta(n)$ will yield division and square root algorithms of the same order. The technique of the next section gives a multiplication whose order is only slightly worse than $\Theta(n)$, and this faster multiplication then can be used to speed up both division and taking square roots.

## 9.5 The Fast Fourier Transform

In our general discussion of divide-and-conquer algorithms we referred to an algorithm for polynomial multiplication which has order $\Theta(n \log n)$, and this section is devoted to a description and explanation of this method. It's based on the technique known as the **Fast Fourier Transform**, usually abbreviated as FFT. The FFT-based polynomial multiplication algorithm is designed for **dense polynomials**, polynomials in which all or almost all of the coefficients are non-zero. Many large-scale polynomial multiplication problems involve **sparse polynomials** and for such problems there are other algorithms which easily outperform the FFT. Because of this, we see that there are practical issues which must be considered before a theoretically good algorithm is used.

The key difference between the FFT method for polynomial multiplication and our earlier methods (pages 254ff) is that here a polynomial is represented by some of its functional values rather than by its coefficients. The basis for the technique is the fact that an $(n-1)^{\text{st}}$ degree polynomial is uniquely determined by its values at $n$ distinct points. To see this, consider the evaluation of a polynomial $f(x) = a_0 + a_1 x + \cdots + a_{n-1}x^{n-1}$ at the $n$ different complex numbers $x = \lambda_1, \lambda_2, \ldots, \lambda_n$. Then

$$(9.8) \qquad (a_0, \ldots, a_{n-1}) \begin{bmatrix} 1 & \ldots & 1 \\ \lambda_1 & \ldots & \lambda_n \\ \vdots & \vdots & \vdots \\ \lambda_1^{n-1} & \ldots & \lambda_n^{n-1} \end{bmatrix} = (f(\lambda_1), f(\lambda_2), \ldots, f(\lambda_n)) \,,$$

where the matrix is **Vandermonde matrix** $V$ associated with $\lambda_1, \lambda_2, \ldots, \lambda_n$. (Refer to Chapter 2.) When the $\lambda_i$ are distinct, we know $V$ is an invertible matrix and accordingly the coefficients of $f(x)$ satisfy

$$(9.9) \qquad (a_0, a_1, \ldots, a_{n-1}) = (f(\lambda_1), f(\lambda_2), \ldots, f(\lambda_n))V^{-1} \,,$$

and so the coefficients are computed from $n$ values of the polynomial. Such a process is called **interpolation** and (9.9) is called an **interpolation formula**. Schematically, we have the bijections

$$\text{Coefficients} \underset{\text{Interpolation}}{\overset{\text{Evaluation}}{\rightleftarrows}} \text{Values}$$

defined by

$$(a_0, a_1, \ldots, a_{n-1}) \xrightarrow{\text{EVALUATION}} (f(\lambda_1), f(\lambda_2), \ldots, f(\lambda_n)) \xrightarrow{\text{INTERPOLATION}} (a_0, a_1, \ldots, a_{n-1})$$

where EVALUATION and INTERPOLATION are the processes which take us from one representation to the other.

For the purpose of polynomial multiplication, encoding a polynomial using $n$ of its functional values is superior to using its coefficients because to obtain the corresponding value for the product polynomial we need only multiply the respective values. But there are some problems. First, the product of two polynomials of degree $n-1$ is a polynomial of degree $2(n-1)$, and therefore the product polynomial is determined by its values at $2n-1$ points. More seriously, you might remember from our discussion of Horner's Method in Chapter 5 that in general it takes $\Theta(n)$ operations to perform one evaluation of a polynomial of degree $n-1$ and hence $\Theta(n^2)$ operations to evaluate it $n$ times. All standard interpolation techniques take $\Theta(n^2)$ operations to interpolate a polynomial of degree $n-1$ through $n$ *generic* points. What saves us here is the freedom to choose the $n$ points in a way that speeds up both evaluation and interpolation. For example, evaluating $f(0)$ takes no work at all and evaluating $f(1)$ takes $n$ additions and no multiplications. Similarly, evaluation of $f(-1)$ also takes no multiplications, since it's the sum of the coefficients which have even indices minus the sum of the other coefficients. We will find algorithms for both EVALUATION and INTERPOLATION which have running time $\Theta(n \log n)$ for the choice of $\lambda_1, \ldots, \lambda_n$ equal to the $n^{\text{th}}$ roots of unity when $n$ is a power of 2.

### *9.5.1 The general form of the Fast Fourier Transform*

Using terminology from Chapter 8, the set $G_n$ of $n^{\text{th}}$ roots of unity is a *group* under multiplication and has $n$ elements. The **principal $\mathbf{n^{th}}$ root of unity**

$$\omega = \cos(\theta) + i \sin(\theta) \text{ where } \theta = \frac{2\pi}{n}$$

has order $n$ in this group, which means every element of $G_n$ can be uniquely written as $\omega^j$ for some $j = 0, 1, \ldots n-1$. Since $1, \omega, \omega^2, \ldots, \omega^{n-1}$ are all

distinct, specializing to $\lambda_j = \omega^{j-1}$ in the **Vandermonde matrix** of (9.8) gives the $n \times n$ matrix

$$V(\omega) = \begin{bmatrix} 1 & 1 & \dots & 1 \\ 1 & \omega^1 & \dots & \omega^{n-1} \\ 1 & \omega^{1\cdot 2} & \dots & \omega^{(n-1)\cdot 2} \\ 1 & \omega^{1\cdot 3} & \dots & \omega^{(n-1)\cdot 3} \\ \vdots & \vdots & \ddots & \vdots \\ 1 & \omega^{1\cdot(n-1)} & \dots & \omega^{(n-1)\cdot(n-1)} \end{bmatrix} . \tag{9.10}$$

Noting that for $0 \le i, j < n$ the dot product

$$(1, \omega^i, \dots, \omega^{(n-1)i}) \cdot (1, \omega^{-j}, \dots, \omega^{-j(n-1)})^T = 1 + \omega^{i-j} + \dots + \omega^{(n-1)(i-j)} ,$$

from (4.36) on page 79 we have

$$(1, \omega^i, \dots, \omega^{(n-1)i}) \cdot (1, \omega^{-j}, \dots, \omega^{-j(n-1)})^T = \begin{cases} n & \text{if } i = j \\ 0 & \text{if } i \neq j \end{cases} .$$

Therefore, the inverse of $V$ is

$$V^{-1}(\omega) = \frac{1}{n} \begin{bmatrix} 1 & 1 & \dots & 1 \\ 1 & \omega^{-1} & \dots & \omega^{-(n-1)} \\ 1 & \omega^{-2} & \dots & \omega^{-2(n-1)} \\ 1 & \omega^{-3} & \dots & \omega^{-3(n-1} \\ \vdots & \vdots & \ddots & \vdots \\ 1 & \omega^{-(n-1)} & \dots & \omega^{-(n-1)^2} \end{bmatrix} = \frac{1}{n} V(\overline{\omega}) ,$$

and

$$V^{-1}(\omega) = \frac{1}{n} V(\overline{\omega}) = \frac{1}{n} \overline{V(\omega)} . \tag{9.11}$$

(Compare this with the inverse of the **generalized Vandermonde matrix** given earlier in Chapter 2.)

Writing $f(x) = a_0 + a_1 x + \dots + a_{n-1} x^{n-1}$, $A = (a_0, \dots, a_{n-1})^T$, and $Y = (f(1), f(\omega), \dots, f(\omega^{n-1}))^T$ equations (9.8) and (9.9) become

$$V(\omega) A = Y \tag{9.12}$$

and

$$A = \frac{1}{n} \cdot \overline{V(\omega)} Y , \tag{9.13}$$

which respectively correspond to EVALUATION and INTERPOLATION. The very special structure of these matrices will allow us to construct quick

divide-and-conquer algorithms for both when $n$ is chosen to be a power of 2.

Historically the term **Discrete Fourier Transform** (or DFT) was used for the general idea described above, with Fast Fourier Transform reserved for the specific implementation with $n = 2^k$. Nowadays this distinction is blurred and all are referred to as the FFT.

### *9.5.2 The FFT when $n = 2^k$*

We set $n = 2^k$ (where $k$ is a natural number), and let $V_k$ denote the Vandermonde matrix $V(\omega)$ constructed using the $n^{\text{th}}$ roots of unity as in (9.10). We assume the powers of the principal $n^{\text{th}}$ root of unity $\omega$ have been found and stored in a table. The FFT also uses the $n/2$ roots of unity, which are already in the table and are found by proceeding through the table in steps of size two.

We now establish an iterative process for getting $V(\omega)$ from $V(\omega^2)$. This will allow us to construct fast divide-and-conquer algorithms for EVALUATION and INTERPOLATION. The matrices for the first, second, and fourth principal roots of unity are respectively

$$V(1) = (1),\ V(-1) = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix},\ \text{and } V(i) = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & i & -1 & -i \\ 1 & -1 & 1 & -1 \\ 1 & -i & -1 & i \end{bmatrix}.$$

To see how to generate $V(\omega)$ from $V(\omega^2)$, note that the first column of $V(-1)$ form a stack of two $V(1)$ and its second column is $\begin{pmatrix} V(1) \\ -V(1) \end{pmatrix}$. Interchanging the second and third columns of $V(i)$ gives

$$\begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & -1 & i & -i \\ 1 & 1 & -1 & -1 \\ 1 & -1 & -i & i \end{bmatrix}, \tag{9.14}$$

whose first two columns form a matrix stack of two copies of $V(-1)$ and its last two columns is the stack $\begin{pmatrix} A \\ -A \end{pmatrix}$ for

$$A = \begin{bmatrix} 1 & 1 \\ i & -i \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & i \end{bmatrix} V(-1).$$

Although it's somewhat of a stretch, we see $V(i)$ can be built in two steps. First construct a block matrix of four $2 \times 2$ blocks, each of which is either $V(i^2)$ or a product of $V(i^2)$ and a diagonal matrix. Then $V(i)$ is obtained from this matrix by performing a suitable permutation of columns. The surprising news is that this procedure can be generalized.

In what follows the indexing of rows and columns of $2^k \times 2^k$ matrices begins with 0. We want to determine the permutation of the columns of $V(\omega)$ which is used. For $k = 2$, we use the binary representations of $0, 1, 2, 3$,

$$00,\ 01,\ 10,\ \text{and}\ 11$$

to define the permutation $\text{Rev}_2$ obtained by *reversing the bits*,

$$00,\ 10,\ 01,\ \text{and}\ 11\,.$$

Applying $\text{Rev}_2$ to the column numbers of $V(i)$ interchanges its second and third columns and keeps the other two fixed, and this *is* the permutation we want. What happens for $k = 3$ is recorded in Table 9.3 where the first row contains the numbers 0 through 7, the second row their binary representations, the third row the bit reversals, and the fourth row contains the decimal equivalents of these reversed representations.

TABLE 9.3. The Bit-Reversal Permutation on $0, \ldots, 7$

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 |
| 000 | 001 | 010 | 011 | 100 | 101 | 110 | 111 |
| 000 | 100 | 010 | 110 | 001 | 101 | 011 | 111 |
| 0 | 4 | 2 | 6 | 1 | 5 | 3 | 7 |

Reading only the top and bottom rows, the permutation $\text{Rev}_3$ of $0, 1, \ldots, 7$ swaps 1 with 4 and 3 with 6.

In general, for any $k$ we define the **bit reversal permutation** $\text{Rev}_k$ of $\{0, 1, \ldots, 2^k - 1\}$ by reversing the bits in this way. The definition of $\text{Rev}_k$ immediately gives the facts that $\text{Rev}_k^2$ is the identity and

$$\text{Rev}_k(j) \text{ is an even integer } \iff 0 \le j < 2^{k-1}\,. \tag{9.15}$$

Application of $\text{Rev}_k$ to the column numbers of the $2^k \times 2^k$ identity matrix, gives a **permutation matrix** which we'll denote by $P_k$. Check that the first three are

(9.16)

$$P_1 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix},\ P_2 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix},\ P_3 = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

Because $\text{Rev}_k$ is the bit-reversal permutation, it has order two in the group of invertible $k \times k$ matrices and $P_k$ is its own inverse. Applying the permutation matrix $P_k$ on the right of $V(\omega)$ gives $F(\omega) = V(\omega)P_k$, the matrix obtained by permuting the columns of $V(\omega)$ by $\text{Rev}_k$.

We refer to $F(\omega)$ as the **Fourier matrix of size** $2^k$. For instance, $F(i)$ is the matrix in (9.14). Since the $(i,j)^{\text{th}}$ entry of $V(\omega)$ is

$$v_{ij} = \omega^{ij} ,$$

then the $(i,j)^{\text{th}}$ entry of $F(\omega)$ is

$$f_{ij} = \omega^{i \operatorname{Rev}_k(j)} .$$

We will use this to show that $F(\omega)$ always has the form we've already noticed for $F(i)$.

**Theorem 9.5.1.** *Let $\omega$ be any primitive $2^k$-th root of unity. Then $F(\omega)$ satisfies the recurrence*

$$F(\omega) = \begin{bmatrix} F(\omega^2) & DF(\omega^2) \\ F(\omega^2) & -DF(\omega^2) \end{bmatrix} ,$$

*where $D$ is the diagonal matrix with diagonal entries $1, \omega, \ldots, \omega^{2^{k-1}-1}$.*

*Proof.* The indices in the upper left quadrant of $F(\omega)$ satisfy $0 \le i, j < 2^{k-1}$, and for such $j$ (9.15) implies $\operatorname{Rev}_k(j)$ is even. Because dividing $\operatorname{Rev}_k(j)$ by 2 removes the least significant bit,

$$f_{ij} = \omega^{i \operatorname{Rev}_k(j)} = (\omega^2)^{i \frac{\operatorname{Rev}_k(j)}{2}} = (\omega^2)^{i \operatorname{Rev}_{k-1}(j)}$$

and this proves the upper left quadrant of $F(\omega)$ is $F(\omega^2)$.

In either lower quadrant, the row index satisfies $2^{k-1} \le i < 2^k$, which we can write as $i = I + 2^{k-1}$ for some $0 \le I < 2^{k-1}$. From this,

$$f_{ij} = \omega^{i \operatorname{Rev}_k(j)} = \omega^{I \operatorname{Rev}_k(j)} \cdot \omega^{2^{k-1} \operatorname{Rev}_k(j)} = f_{Ij} \cdot (-1)^{\operatorname{Rev}_k(j)} .$$

Because $\operatorname{Rev}_k(j)$ is even in both left quadrants this proves that the two left quadrants are the same, and the matrices that form the two right quadrants are negatives of each other.

It remains to show that the upper right quadrant is $DF(\omega^2)$. In that quadrant the indices satisfy $0 \le i < 2^{k-1} \le j$, and $j = J + 2^{k-1}$ for some $0 \le J < 2^{k-1}$. Since $\operatorname{Rev}_k(j) = 2 \operatorname{Rev}_{k-1}(J) + 1$, then

$$\frac{f_{ij}}{\omega^i} = \omega^{i(\operatorname{Rev}_k(j))} \omega^{-i} = \omega^{i(\operatorname{Rev}_k(j)-1)} = (\omega^2)^{i \operatorname{Rev}_{k-1}(J)}$$

which is the $(i,J)^{\text{th}}$ entry of $F(\omega^2)$. This completes the proof. □

We will next show that the recursive structure of this Fourier matrix yields a divide-and-conquer strategy for EVALUATION and INTERPOLATION.

### *9.5.3 Fast evaluation and fast interpolation*

For the polynomial $f(x) = a_0 + a_1 x + \ldots + a_{n-1}x^{n-1}$ recorded as the coefficient vector $A = (a_0, \ldots, a_{n-1})^T$, we recall that (9.12) can be used to obtain the evaluation vector $Y = (f(1), f(\omega), \ldots, f(\omega^{n-1}))$. We have encoded this as

$$Y = F(\omega)P_k A \tag{9.17}$$

and the reverse interpolation process given in (9.13) can be written as

$$A = \frac{1}{n} \cdot \overline{F(\omega)} P_k Y \,. \tag{9.18}$$

Comparing these two equations, we see that *INTERPOLATION is essentially the same process as EVALUATION* ! This means that the analysis of INTERPOLATION is the same as that for EVALUATION.

To analyze EVALUATION we consider the time for performing the permutation $P_k A$, and computing $\mathrm{Rev}_k(j)$ for any $0 \le j < 2^k$ can be done in time proportional to the length of the string $k = \log(n)$ and computing the permutation $\mathrm{Rev}_k$ takes time $O(n \log n)$. Once $\mathrm{Rev}_k$ is known, applying it to $A$ amounts to swapping pairs of indices or pointers and takes time $O(n)$. Therefore, the time for finding $X = P_k A$ is $O(n \log n)$ where $Y = F(\omega)X$.[1]

The decomposition of the Fourier matrix given in Theorem 9.5.1 suggests a divide-and-conquer strategy for computing $Y$. To see this, we write any $X \in \mathbb{C}^{2^k}$ as $X = \begin{pmatrix} X_1 \\ X_2 \end{pmatrix}$ where $X_1$ is the vector consisting of the first $2^{k-1}$ components of $X$ and $X_2$ contains its last $2^{k-1}$ components. Then the special form in the theorem gives

$$F(\omega)X = F(\omega)\begin{pmatrix} X_1 \\ X_2 \end{pmatrix} = \begin{pmatrix} F(\omega^2)X_1 + DF(\omega^2)X_2 \\ F(\omega^2)X_1 - DF(\omega^2)X_2 \end{pmatrix} .$$

Noting that the products $F(\omega^2)X_1$ and $F(\omega^2)X_2$ need only be computed once, this decomposition gives a divide-and-conquer algorithm for computing $F(\omega)X$ whose schematic description is

$$A \xrightarrow{\text{permute}} P_k A =: \begin{pmatrix} X_1 \\ X_2 \end{pmatrix} \xrightarrow{\text{half-size}} \begin{pmatrix} F(\omega^2)X_1 \\ F(\omega^2)X_2 \end{pmatrix} \xrightarrow{\text{combine}} \begin{pmatrix} F(\omega^2)X_1 + DF(\omega^2)X_2 \\ F(\omega^2)X_1 - DF(\omega^2)X_2 \end{pmatrix} = Y \,.$$

[1] Many texts ignore the time necessary to compute the permutation since in practice it takes very little time compared to the rest of the algorithm. Even if we had to apply a permutation in every iteration, the asymptotic running time would not be affected (for this, refer to Exercise 9.19.)

Likewise, INTERPOLATION can be described by the schematic

$$\begin{aligned} Y \xrightarrow{\text{conjugate and permute}} P_k\overline{Y} =: \begin{pmatrix} X_1 \\ X_2 \end{pmatrix} &\xrightarrow{\text{half-size}} \begin{pmatrix} F(\omega^2)X_1 \\ F(\omega^2)X_2 \end{pmatrix} \\ &\xrightarrow{\text{combine}} \begin{pmatrix} F(\omega^2)X_1 + DF(\omega^2)X_2 \\ F(\omega^2)X_1 - DF(\omega^2)X_2 \end{pmatrix} = V(\omega)\overline{Y} \\ &\xrightarrow{\text{conjugate and scale by 1/n}} \frac{1}{n}\overline{V(\omega)}Y = A\,. \end{aligned}$$

Note that in this process the matrix $F(\omega)$ never needs to be calculated because all Fourier matrices in the scheme are simply recursive calls to a procedure. Also, the diagonal entries of $D$ are some of the powers of $\omega$ which are stored in an array and are pulled from the array when necessary.

We've already shown the necessary permuting can be done in $O(n \log n)$ time, and we will now compute the running time of the divide-and-conquer part. Since the combine step involves multiplying by a diagonal matrix and then performing an addition and a subtraction, that step takes $\Theta(n)$ operations. Because we have divided the problem into two half-size subproblems, the divide-and-conquer recurrence has $a = c = 2$, and

$$T(n) = 2T(n/2) + bn\,,$$

which is the second case of the divide-and-conquer formula in Theorem 9.4.1 and so $T(n) = \Theta(n \log n)$. Finally, since appending zero coefficients when $n$ isn't a power of two only increases the implied constant in $\Theta(n \log n)$, multiplying two polynomials of degree less than $n$ can also be accomplished in $\Theta(n \log n)$ operations.

### *9.5.4 The fast polynomial multiplication algorithm*

The entire algorithm for computing the coefficients of the product of two polynomials can be summarized as follows:

**Fast Polynomial Multiplication**

Let $f, g \in \mathbb{C}[x]$ be polynomials with $\deg(f), \deg(g) < n$.

1. Place the coefficients of the polynomial $f$ in the first $n$ components of a vector $A$ of length $2^{k+1}$ where $2^k$ is the smallest power of 2 which is at least $n$. Set the remaining components of $X$ to zero.
2. Place the coefficients of $g$ into a vector $B$ of length $2^{k+1}$.
3. Permute both $A$ and $B$, obtaining $X$ and $Y$ respectively.
4. For the principal $2^{k+1}$-th root of unity $\omega$, use the recursive algorithm to compute both $F(\omega)X$ and $F(\omega)Y$.
5. Let $Z$ be the componentwise product of these two vectors.
6. Permute and conjugate $Z$.
7. Use the recursive algorithm to compute $F(\omega)Z$.
8. Conjugate and divide each component of the result in Step 7 by $2^{k+1}$. This is the vector of coefficients of the product $fg$.

The computation will only be *approximate* because of the round-off in floating point operations and because we only use finite approximations to the powers of $\omega$. The first $2n-1$ components of the output vector are approximations to the coefficients of the product polynomial. Although all higher coefficients in the product $fg$ are zero, some of these components might be non-zero in the output and the size of these extra components gives some indication of the accuracy of the computation.

Schematically the algorithm is




FIGURE 9.3. Schematic Representation of FFT Polynomial Multiplication.

Any divide-and-conquer algorithm must specify what is meant by "small", the size of problem which is completed "by hand". In this algorithm, a vector with two components is the small case, that is, $k = 0$ and the polynomials are constant. We then compute $f(\pm 1)$ and $g(\pm 1)$, and continue into the combine algorithm. Other variations are possible. For instance, $n = 4$

is a popular choice for "small" because $f(\pm i)$ and $g(\pm i)$ can be computed without a multiplication.

We illustrate the FFT algorithm by applying it to the simple example of squaring the polynomial $2 + x$. In Step 1, we form the vector $(2, 1, 0, 0)$, and Step 2 is the same for this example. In Step 3 we use $\text{Rev}_2$ to permute the vectors and obtain $(2, 0, 1, 0)$. In Step 3 the recursive routine is applied to the two (identical) vectors, which we break into two vectors $(2, 0)$ and $(1, 0)$. Since

$$F(-1) = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix},$$

left-multiplication of $(2, 0)$ and $(1, 0)$ by $F(-1)$ gives $(2, 2)$ and $(1, 1)$. We next multiply $(1, 1)$ by the diagonal matrix with 1 and $i$ on the diagonal, yielding $(1, i)$. We now add and subtract this from $(2, 2)$ and place the results in a vector to get

$$(2 + 1, 2 + i, 2 - 1, 2 - i) = (3, 2 + i, 1, 2 - i).$$

These numbers are supposed to be the values of the polynomial $f(x) = 2 + x$ at $x = 1$, $i$, $-1$, and $-i$, which indeed they are.

In Step 5 we multiply the two (in our case identical) value vectors componentwise. This amounts to squaring the vector above and yields

$$Z = (9, 3 + 4i, 1, 3 - 4i),$$

whose coordinates are the values of the polynomial $4 + 4x + 4x^2$ at $x = 1$, $i$, $-1$, and $-i$. In Step 6 we permute using $\text{Rev}_2$ and conjugate to get the vector

$$(9, 1, 3 - 4i, 3 + 4i).$$

In Step 7 we apply the same recursive algorithm, starting at the above vector. It splits into $(9, 1)$ and $(3 - 4i, 3 + 4i)$, and these become $(10, 8)$ and $(6, -8i)$ in the base case of the recursion. We multiply $(6, -8i)$ by the diagonal matrix with 1 and $i$ on its diagonal to get $(6, 8)$, and then $(10, 8)$ and $(6, 8)$ are added, subtracted, and placed in the vector

$$(10 + 6, 8 + 8, 10 - 6, 8 - 8) = (16, 16, 4, 0).$$

In Step 8 we conjugate (which has no effect since our values at this point are real) and divide by $2^2 = 4$ to get the vector

$$(4, 4, 1, 0),$$

the coefficient vector of $f(x)g(x) = (2 + x)^2 = 4 + 4x + x^2$.

Notice that the FFT multiplies *complex* polynomials. Some improvement is possible if all coefficients in both polynomials are real numbers. One minor change is that it is never necessary to conjugate in Step 6. More significantly, since the evaluation is done at $2^{k+1} \approx 2n$ complex points, one can pack a pair of real numbers into one complex number and design a real FFT, which evaluates at only $2^k$ instead of $2^{k+1}$ complex points. (For details on this, refer to Exercise 9.20 and [2].)

## 9.6 Average Case Analysis

Until now we've considered the running time of an algorithm to be a function of the input size. In practice, an algorithm might treat all inputs of the same size in the same way or it could handle some inputs more quickly than others. Because of this, we define the best case, worst case, and average case running times for an algorithm. The maximum running time over all inputs of the same size is called the **worst case running time**. The minimum running time over all inputs of the same size is called the **best case running time**. Averaging the running time over all inputs of the same size is called the **average case running time**. When the probability associated with each of the various inputs of a particular size is unknown it's difficult to calculate the average case time. For definiteness and simplicity it's often assumed that each input of a fixed size is equally likely to occur and then the average case running time can be calculated.

### *9.6.1 The LARGETWO algorithm*

Consider the following algorithm which finds the two largest entries in a one-dimensional array and assigns these values to the variables FIRST and SEC.

```
Procedure  LARGETWO(C)
FIRST:= C[1]
SEC:= C[2]
FOR I = 2 TO n DO
    IF C[I] > FIRST
       THEN SEC:=FIRST ; FIRST:= C[I]
       ELSE IF C[I] > SEC
               THEN SEC:= C[I]
```

Note that this algorithm assumes that the array has at least two components because the results are unpredictable when used with a one-dimentional array.

The ground rules for our running time analysis are that we'll count the number of comparisons of array entries and ignore any comparisons of numbers used to control the FOR loop. In the FOR loop the algorithm makes either one or two comparisons for each $I > 1$ and for an $n$-array the algorithm uses at least $n-1$ comparisons and at most $2(n-1)$ comparisons.

If the data happened to be arranged in order of increasing coordinates,

$$C[1] < C[2] < \ldots < C[n]\,,$$

the algorithm will perform exactly $n-1$ comparisons and this implies that the best case running time is $B(n) = n-1$. The algorithm makes $2(n-1)$ comparisons when $C[1]$ is the largest entry and this is the worst case with running time $W(n) = 2(n-1)$.

What about the average case behavior? Is it close to worst case, close to best case, or midway between the worst and best cases? Let $A(n)$ be the number of comparisons used on average by this algorithm. Since the algorithm begins with the first entry and proceeds in one direction across the array, we may reasonably assume that for the first $n-1$ entries the algorithm will on average use $A(n-1)$ comparisons, the same number it would use if the last entry weren't there. For the last entry, it will use at least one comparison. If $C[n]$ is not the largest element, then $C[n] > FIRST$ is false and a second comparison must be made. If $C[n]$ *is* the largest element, then $C[n] > FIRST$ is true and only that one comparison is made. By our assumption of uniformity, the probability that $C[n]$ is the largest is $1/n$ and from these considerations, for all $n \geq 2$

$$A(n) = A(n-1) + 2\left(1 - \frac{1}{n}\right) + 1 \cdot \frac{1}{n} = A(n-1) + 2 - \frac{1}{n}\,. \tag{9.19}$$

When the array has two entries, uniformity implies that the maximum entry is equally likely to lie in either entry. This implies that two comparisons are required exactly half of the time, giving $A(2) = 3/2$.

Since this is a first-order recurrence with eigenvalue $\lambda = 1$ and forcing function $\psi(i) = 2 - 1/i$, the solution is

$$A(n) = 1 + \sum_{i=1}^{n} (2 - \frac{1}{i}) = 2n + 1 - \sum_{i=1}^{n} \frac{1}{i}\,.$$

Using our result on the partial sums of the harmonic series given in Exercise 8.36, $A(n)$ is therefore asymptotically approximated by $2n+1-\ln(n)+c$, where $c$ is some constant. From this we see that the average case behavior of the algorithm is very close to the worst case, $W(n) = 2(n-1)$, especially as compared with the best case running time, $B(n) = n-1$. Also, notice that the average of the worst and best case running time is $\frac{3}{2}(n-1)$, a severe underestimate of the actual average case behavior.

Here's another analysis of the average case running time for LARGETWO. Let $C[j]$ be the maximal entry. From our assumption of uniformity, the probability that $C[j]$ is the largest entry is $1/n$. As noted above, $2(n-1)$ comparisons are made for $j = 1$. For $j \geq 2$ the algorithm will perform $A(j-1)$ comparisons before the loop reaches $I = j$, then for $I = j$ the loop

makes one comparison and for each of $I = j+1, \ldots, n$ two comparisons are used. This gives

$$nA(n) = \sum_{j=2}^{n} \left[ A(j-1) + 1 + 2(n-j) \right].$$

Expanding the sume gives

$$\begin{aligned} nA(n) &= \sum_{j=2}^{n} A(j-1) + \sum_{j=2}^{n} 1 \; + 2 \sum_{j=1}^{n} (n-j) \\ &= \sum_{j=1}^{n-1} A(j) + n - 1 + 2 \sum_{j=0}^{n-1} j \\ &= \sum_{j=1}^{n-1} A(j) + n - 1 + (n-1)n \\ &= \sum_{j=1}^{n-1} A(j) + n^2 - 1 \,. \end{aligned}$$

Therefore, for all $n \geq 2$

$$nA(n) = \sum_{j=1}^{n-1} A(j) + n^2 - 1 \,. \tag{9.20}$$

This is our first encounter with recurrences of this form, which depend on all previous terms. Similarly to some recurrences in Chapter 4, the solution can be shown to satisfy a first-order recurrence. Since

$$(n-1)A(n-1) = \sum_{j=1}^{n-2} A(j) + (n-1)^2 - 1 \,,$$

then

$$\begin{aligned} nA(n) - (n-1)A(n-1) = A(n-1) + n^2 - (n-1)^2 &= A(n-1) + 2n - 1 \,; \\ nA(n) &= nA(n-1) + 2n - 1 \,, \end{aligned}$$

which is the same recurrence found earlier in (9.19).

### *9.6.2 The QUICKSORT algorithm*

Next we consider QUICKSORT which has a more complicated average case analysis. The algorithm sorts an array $S$ according to increasing order of

entries where we allow repeated entries in $S$. The output of this algorithm is the sorted string of numbers, and the dots in the RETURN statement indicate concatenation of strings.

```
Procedure QUICKSORT(S)
Pick an entry α of S at random.
Divide S into three parts, S1, S2, S3 where
  S1 is the set of elements of S which are less than α.
  S2 is the set of elements of S which equal α.
  S3 is the set of elements of S which are greater than α.
RETURN(QUICKSORT(S1) · S2· QUICKSORT(S3))
```

This is a **random algorithm** because it uses a random choice at some point, in this case for the choice of the entry $\alpha$. The choice could be randomized by using a random number generator, but usually some deterministic rule is used in the hope that the distribution of the input data is not significantly correlated with the choice of rule. We'll assume this correlation is low enough to allow us to treat the choice of $\alpha$ as truly random for the analysis. Three popular rules for choosing $\alpha$ are: always choose the first entry; always choose the last entry; always choose the middle entry. Another popular method is to choose the median of the first, last, and middle entries.

The running time of this algorithm on an array $S$ of size $n$ satisfies

$$T(S) = T(S_1) + bn + T(S_3)\,,$$

where $b$ accounts for the time used to place elements into the three sets and return the answer. The worst case occurs when $S_1$ or $S_3$ contains $n-1$ entries and

$$W(n) = W(n-1) + bn\,,$$

giving

$$W(n) = 1 + b\sum_{i=1}^{n} i = 1 + \frac{b}{2}n(n+1) = \Theta(n^2)\,.$$

The best case for QUICKSORT occurs when $S_1$ and $S_3$ each contain approximately $n/2$ elements. Then

$$B(n) = 2B(n/2) + cn\,,$$

for some constant $c$. This is a divide-and-conquer recurrence with $a = c = 2$ and $m = 1$, and $B(n) = \Theta(n \log n)$.

For this algorithm there's a large gap between the worst case, $W(n) = \Theta(n^2)$, and the best case, $B(n) = \Theta(n \log n)$. Does the algorithm on average

behave closer to worst case or closer to best case? Notice that if the average case running time were the numerical average of worst case and best case, then the average would be $\Theta(n^2)$ and therefore close to the worst case but we'll show the opposite. On average the running time of QUICKSORT is $\Theta(n \log n)$ and hence behaves close to best case. This is intuitively plausible because nearly even splits should on average occur much more frequently than one-sided splits.

As usual, for $n \geq 1$ let $A(n)$ be the average case running time for an array with $n$ entries and assume that $\alpha$ is equally likely to be any of the entries. Letting $A(0) = 0$, for $n \geq 1$

$$A(n) = \frac{1}{n} \sum_{j=1}^{n} [A(j-1) + A(n-j) + b(n+1)]$$

where $b(n+1)$ is the time for split and combine (and is written in this form because it slightly simplifies the computation.) Therefore,

$$nA(n) = \sum_{j=1}^{n} A(j-1) + \sum_{j=1}^{n} A(n-j) + bn(n+1) = 2 \sum_{j=1}^{n-1} A(j) + bn(n+1) .$$

Using the technique of the previous example,

$$nA(n) - (n-1)A(n-1) = 2A(n-1) + 2bn ,$$

giving

$$nA(n) = (n+1)A(n-1) + 2bn ;$$

$$\frac{A(n)}{n+1} = \frac{A(n-1)}{n} + \frac{2b}{n+1} .$$

Setting $Z(n) = A(n)/(n+1)$, this becomes

$$Z(n) = Z(n-1) + \frac{2b}{n+1} ,$$

a first-order recurrence with forcing function $\psi(j) = 2b/(j+1)$, giving

$$Z(n) = c_1 + 2b \sum_{j=1}^{n} \frac{1}{j+1}$$

and

$$A(n) = c_1(n+1) + 2b(n+1) \sum_{j=1}^{n} \frac{1}{j+1} .$$

Using Exercise 8.36 again, we obtain constants $c_2, c_3 > 0$ such that

$$A(n) \to c_2(n+1) + c_3(n+1) \log(n) ,$$

and gives $A(n) = \Theta(n \log(n))$.

This analysis shows that QUICKSORT has an average case behavior which is close to the best case behavior, suggesting that QUICKSORT may be a good method to use to sort an array. However, remember that the analysis assumes that each permutation of the input behavior is equally likely. If that assumption were not true, then it isn't clear that QUICKSORT is a good algorithm to use. For instance, what if the data is in order except for a few items which are out of order? If you always choose the first input for $\alpha$ then the splits will usually give one very large set. It can be proved that the average running time of QUICKSORT for this type of data has order $\Theta(n^2)$.

A lesson to be learned from these two examples is that you know that the average case is *somewhere* between worst case and best case, but it can be anywhere between worst and best. Further, in many practical cases you'll have no idea about the distribution of the input data. The assumption of a uniform probability distribution often makes the average case calculation easy, but it is probably not applicable to a general input distribution.

## 9.7 Exercises

**Ex 9.1.** Use our perspective of moving the largest disk to derive a recurrence relation for the *minimum* number of moves needed to solve the $n$ disk Towers of Hanoi puzzle. Solve your recurrence relation with appropriate initial conditions and conclude that any algorithm which solves the $n$ disk Towers of Hanoi puzzle has running time $\Omega(2^n)$. (Refer to Chapter 1 for the meaning of $\Omega$.)

**Ex 9.2.** Give an algorithm for the Towers of Hanoi puzzle which uses more than the minimal number of moves. Derive and solve a recurrence relation for your algorithm and show that it uses more than the minimum number of moves calculated in the previous exercise. (If you don't see the usefulness of constructing algorithms which take longer than necessary consider the problem of The Cab Driver and the Tourist.)

**Ex 9.3.** In the following algorithm, the towers are labeled 0, 1, and 2 rather than $A$, $B$, and $C$. The variable `COUNT` contains $n$ bits numbered from 1 to $n$ starting at the rightmost bit. The positions in `COUNT` alternate between odd and even (in that order).

```
Procedure TOWERS(n)
          T:=0 (*Tower number computed modulo 3*)
          COUNT:=0
          p:= { +1  if n is even
              { −1  if n is odd
          WHILE TRUE DO
                MOVE DISK 1 FROM T TO T+P
                T := T+P
                COUNT := COUNT+1
                IF COUNT = ALL 1's THEN EXIT
                IF RIGHTMOST 0 IN COUNT IS IN EVEN POSITION
                   THEN MOVE DISK FROM T-P TO T+P
                   ELSE MOVE DISK FROM T+P TO T-P
                COUNT := COUNT+1
          ENDWHILE
```

(a) Without using a recurrence relation show that this algorithm uses the minimal number of moves.
(b) Even if incrementing COUNT and finding the rightmost 0 takes time $cn$ for constant $c$, show that this algorithm has the running time $\Theta(2^n)$.

**Ex 9.4.** (Refer to [17].) The following algorithm for the Towers of Hanoi problem has the towers arranged in a circle. By considering the situation when the largest disk is moved, derive and solve recurrence relations for the number of moves made and for the running time of this algorithm.

```
MOVE SMALLEST DISK ONE TOWER CLOCKWISE
WHILE A DISK OTHER THAN THE SMALLEST DISK CAN BE MOVED
      DO MOVE THAT DISK
         MOVE THE SMALLEST DISK ONE TOWER CLOCKWISE
ENDWHILE
```

(Notice that if the number of disks is even, then the $n$ disks are moved one tower counterclockwise, while if $n$ is odd, the disks are moved one tower clockwise.)

**Ex 9.5.** Which of the three algorithms for the Towers of Hanoi puzzle is fastest? The recursive algorithm and the algorithms from the last two exercises all have running time $\Theta(2^n)$. To determine which is really fastest, we suggest programming them and running them on a real computer. (You

may want to make a small modification in the algorithm from the last exercise, always moving the disks to the same tower regardless of the parity of $n$.) Can you explain why one algorithm might be slower than the others in a real-world programming environment?

**Ex 9.6.** For a given one-dimensional array $A$, let $A[i]$ denote the content of the $i^{\text{th}}$ entry of $A$. Show that the following procedure has running time $\Theta(n)$ by writing down and solving the appropriate recurrence relation. Describe in English what the algorithm accomplishes.

```
Procedure  LARGEST(n)
IF n > 1 THEN LARGEST(n − 1)
              IF A[n − 1] > A[n]
                 THEN TEMP := A[n]
                      A[n] := A[n − 1]
                      A[n − 1] := TEMP
```

**Ex 9.7.** Using the result of the last exercise, give an inductive proof that the following algorithm sorts $(A[1], A[2], \ldots, A[n])$. Derive and solve a recurrence relation for the running time.

```
Procedure SORT(n)
IF n > 1 THEN LARGEST(n)
              SORT(n − 1)
```

**Ex 9.8.** According to an old tale told by Édouard Lucas, the monks in a secret monastery in Hanoi are going through the moves for the Towers of Hanoi puzzle with 64 disks. When they finish, the world will end. If it takes them one minute to move a disk, should you worry about the end of the world? What if they can move one disk per second? After a wily computer salesperson convinces the monks to automate so that they can simulate moving a disk in one nanosecond, should you worry? (Also refer to the title story in *The Nine Billion Names of God: The Best Short Stories of Arthur C. Clarke*, Harcourt, Brace & World, New York, 1967.)

The following secure locking system is used in the next four exercises.

The locking system assumes $n$ locks are connected so that

1. Lock 1 may be changed from locked to unlocked or from unlocked to locked at any time.
2. For any $j > 1$, lock $j$ may be changed from locked to unlocked (or vice versa) only if lock $j-1$ is locked and locks 1 through $j-2$ are unlocked.

**Ex 9.9.** One strategy for designing such a system involves thinking of unlocking all the locks in terms of unlocking a subseries of locks, unlocking the last lock, and then relocking the subseries so that you can repeat this process to unlock all the locks. This strategy naturally leads to a design in which you have two mutually recursive procedures, or one recursive procedure with a switch which indicates whether the procedure will be locking or unlocking.

(a) Use this strategy to design a recursive algorithm whjch gives the sequence of operations needed to unlock a series of $n$ locks which are all initially locked.
(b) Find the pair of recurrence relations which give the time and number of locking/unlocking operations used by your algorithm.
(c) Solve the relations in part (b).

**Ex 9.10.** A second strategy rests on the observation that, after unlocking the last lock, the other locks will be in a specific configuration, and to unlock the next to last lock, it is necessary to re-configure the locks. You can design a simple recursive procedure which takes the locks from the first configuration to the second configuration. This strategy leads to a design with two nested recursive procedures.

(a) Use this strategy to design a recursive algorithm whjch gives the sequence of operations needed to unlock a series of $n$ locks which are all initially locked.
(b) Find the pair of recurrence relations which give the time and number of locking/unlocking operations used by your algorithm.
(c) To solve the relations in part (b), observe that The first depends on the second, but the second does not depend on the first. Thus, you can solve the second equation and plug its solution into the first equation and solve.

**Ex 9.11.** Note the procedures in the previous two exercises use the same amount of time and number of moves because they do the same operations in reverse order. Show that your algorithm uses the minimal number of locking/unlocking operations.

**Ex 9.12.** To get a feel for the puzzle you've investigated in the last problems, go to your local toy store and get SPIN-OUT®. This is a physical embodiment of the puzzle with seven locks. Amaze your friends by solving SPIN-OUT® using the minimum number of moves.

**Ex 9.13.** Give a divide-and-conquer algorithm for finding the largest entry in a vector with $n$ components and show that your algorithm uses $n-1$ comparisons.

**Ex 9.14.** Design a divide-and-conquer algorithm to find the two largest entries in an array. Show your algorithm is correct and calculate the number of comparisons it uses. Give an example which shows that your algorithm uses more comparisons than necessary.

**Ex 9.15.** Suppose you have a large number of coins and a two-pan balance whose pans are as large as needed. The balance will tell you if the coins in one pan weigh the same as the coins in the other pan or which set of coins is heavier. Among your coins is exactly one which has a different weight from the other coins.

(a) Assuming that the number of coins is a power of 3, design a divide-and-conquer algorithm to find the odd coin.
(b) Prove your algorithm is correct.
(c) Find and solve a recurrence for the number of times your algorithm uses the balance.

**Ex 9.16.** Two strings $C_1$ and $C_2$ **commute** (that is, $C_1C_2 = C_2C_1$) iff there is a string $w$ and two natural numbers $k_1$ and $k_2$ so that $C_1 = w^{k_1}$ and $C_2 = w^{k_2}$. Beginning with a constructive proof that such $w$ exists, construct an algorithm which finds $w$. (Note that here $w^k$ denotes the juxtaposition of $k$ copies of $w$, that is, $ww\cdots w$.)

**Ex 9.17.** Show how the divide-and-conquer strategy with the clever identity (9.4) can be used to construct an algorithm for multiplying large integers and which has running time $\Theta(n^{\log_2 3})$, where $n$ is the number of bits in the larger integer.

**Ex 9.18.** Show that the power

$$\begin{bmatrix} 1 & 1 & 2 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}^n$$

can be computed in $O(n)$ bit operations. What special properties of this matrix allows this "fast" algorithm?

**Ex 9.19.** Consider a recurrence of the form $T(n) = 2T(n/2) + g(n)$ with $g(n) \geq 0$. Argue that the asymptotic order of growth of $T(n)$ is monotonic in $g(n)$ in the following sense: If $h(n) = O(g(n))$ and $S(n) = 2S(n/2)+h(n)$,

then $S(n) = O(T(n))$. Then, using the technique of letting $t_k = T(2^k)$, show that $T(n) = O(n \log n)$ if and only if $g(n) = O(n \log n)$.

**Ex 9.20.** Set up and solve a recurrence relation for the number of *real* multiplications the FFT algorithm uses to multiply two real polynomials with $n$ coefficients.

**Ex 9.21.** (a) The number of multiplications used by the FFT depends on the size of the agreed upon "small size" problem. Work out the difference in the number of multiplications among FFTs recursing to one component; two components; four components.

(b) Compare your answers in part (a) with the $n^2$ multiplications used by the standard algorithm for polynomial multiplication.

(c) Calculate the value of $n_0$ for which the FFT is faster than the standard method for polynomials with $n \geq n_0$ coefficients. Do you think the FFT is useful in practice? If you had to multiply two polynomials with $2^{10}$ coefficients, which method would be faster?

(d) Assume that the time for all operations used is dominated by the time for multiplication. What is your prediction for the ratio of the running times of the FFT and the standard algorithm?

**Ex 9.22.** Do an average case analysis for the following procedure:

```
Procedure BIGTWO(C)
FIRST:=C[1]
SEC:=C[2]
FOR I:=2 TO n DO
    IF C[I]≥ SEC
       THEN IF C[I] > FIRST
                THEN SEC:=FIRST ; FIRST:=C[I]
                ELSE SEC:=C[I]
```

Is the average case running time for this algorithm nearer to its worst case or its best case? Which of the two procedures LARGETWO(C) or BIGTWO(C) would you use?

**Ex 9.23.** If the best case $B(n)$ of an algorithm satisfies

$$B\Big(\frac{n}{2} - \gamma\Big) + B\Big(\frac{n}{2} + \gamma\Big) + cn \leq B(n) \leq B\Big(\frac{n}{2} - \alpha\Big) + B\Big(\frac{n}{2} + \alpha\Big) + bn$$

for constants $\alpha, \gamma, b, c$, show that $B(n) = \Theta(n \log n)$.

**Ex 9.24.** Show that if the matrix $C$ is initialized to the zero matrix, then the following algorithm will compute the $n \times n$ matrix product $C = A \times B$.

```
Procedure
        For i = 1 TO n DO
            For j = 1 TO n DO
                For k = 1 TO n DO
                    c_ij = c_ij + a_ik * b_kj
```

Compute the running time of this algorithm. Explain why most computations use this algorithm rather than Strassen's algorithm from Section 9.4.2.

**Ex 9.25.** Show that if multiplication of two $n$ bit numbers can be performed in time $\Theta(n^{1+\epsilon})$ for some $\epsilon \geq 0$ then division of two $n$ bit numbers can also be carried out in time $\Theta(n^{1+\epsilon})$. (Refer to Section 9.4.4.)

**Ex 9.26.** Show that multiplication of two $n$ bit numbers can be performed in the same time order as division of two $n$ bit numbers.

**Ex 9.27.** Compute the product of the Fourier matrices $F(i)$ and $F(-i)$ to show that $F(\omega)^{-1} \neq F(\overline{\omega})$. Find a simple formula for $F(\omega)^{-1}$, the **inverse Fourier matrix**.

# 10
# Some Nonlinear Recurrences

## 10.1 Some Examples

In the previous chapters we have primarily discussed *linear* equations or, said another way, only equations involving *linear operators*. Recall that a function $L$ on the space of sequences is a **linear operator** if it satisfies the two conditions:

1. $L[x+y] = L[x] + L[y]$
2. $L[cx] = cL[x]$,

for all sequences $x =< x_n >$ and $y =< y_n >$ and all constants $c$. (Notice that a square matrix is a linear operator on the space of vectors of appropriate size since a matrix would satisfy these conditions when $x$ and $y$ are any vectors and $c$ is any scalar.) While the $+$ above is our usual addition for sequences, one could consider replacing $+$ by other operations. We will not follow this tack. Instead we want to look at truly nonlinear equations. To keep things simple, we will look at one-dimensional equations of the form:

$$x_{t+1} = f(x_t).$$

For linear equations there is essentially only one linear one-dimensional equation, but to paraphrase Tolstoy: "Equations can be linear in only one way, but equations can be nonlinear in many different ways." So we should not expect to have a general theory for nonlinear equations. Rather, we would hope to have different theories for different classes of nonlinear equations.

Let us consider a very simple nonlinear example:

$$x_{t+1} = \frac{1}{x_t}.$$

If $x_0 = 2$, then $x_1 = 1/2$, $x_2 = 2$, $x_3 = 1/2$, and the sequence will **oscillate** with period 2. In general, the sequence will be $x_0, 1/x_0, x_0, 1/x_0$, that is the sequence will generally have period 2. If $x_0 = 1$, or $x_0 = -1$, then $x_n = x_0$ for all $n \geq 0$ and so the sequence will have period 1, i.e., the sequence will be a constant sequence. The other exceptional case is $x_0 = 0$. Here $x_1 = 1/0$ is undefined. We can define $1/0 = \infty$ and $1/\infty = 0$, and obtain a period 2 oscillation. Even though this equation is nonlinear, we can guess that in some sense it is analogous to the linear equation

$$y_{n+1} = -y_n,$$

which generally has period 2, and has period 1 if $y_0 = 0$. In Section 10.6, we will follow up this analogy to linear equations.

As another simple example, consider

$$x_{n+1} = \sqrt{x_n}.$$

For this equation to make sense, we will assume that $x_0 \geq 0$ and that $\sqrt{x}$ returns the nonnegative square root of $x$. We can calculate some iterates:

$$\begin{aligned} x_1 &= x_0^{1/2} \\ x_2 &= x_1^{1/2} = (x_0^{1/2})^{1/2} = x_0^{1/4} \\ x_3 &= x_2^{1/2} = (x_0^{1/4})^{1/2} = x_0^{1/8} \end{aligned}$$

and see that the solution is

$$x_n = x_0^{1/2^n}.$$

There are four cases

(a) $x_0 = 0$, and then $x_n = 0$ for all $n \geq 0$,
(b) $0 < x_0 < 1$, and then $1 > x_{n+1} > x_n > 0$,
(c) $x_0 > 1$, and then $1 < x_{n+1} < x_n$,
(d) $x_0 = 1$, and then $x_n = 1$ for all $n \geq 0$.

We can summarize these cases by saying that 0 and 1 are **fixed points** of the system since $f(p) = p$ for $p = 0, 1$. The fixed point 0 is **unstable**, while 1 is a **stable fixed point** which attracts all solutions with $x_0 > 0$.

Our point, so far, is that *some* nonlinear equations are not difficult to analyze. But there are nonlinear equations which are more difficult. Consider the example

$$x_{n+1} = a \sin(x_n).$$

One can formally write that

$$x_n = a \sin(a \sin(\dots (x_0) \dots)),$$

but this formula gives no clear information about the behavior of the solutions. Some things are easy to see, for example, if $x_0$ is an integer multiple of $\pi$, then $x_1 = 0$ and $x_n = 0$ for all $n > 0$. On the other hand, it is not clear what sort of solutions will arise for other values of $x_0$. In particular, what will happen if $x_0$ is a small positive number? Will the solution be attracted to 0? Will the solution converge to some positive value? Will the solution become periodic? In general, these questions are difficult to answer.

## 10.2 Nonlinear Systems

The preceding examples suggest that nonlinear difference equations may be difficult to analyze. In the remainder of this chapter we want to describe and give examples of some commonly used methods. Of course, in a short chapter we cannot cover all techniques, so we refer the interested reader to the many books like LaSalle [81], Devaney [43], and Parker and Chua [109] which include more detailed information.

To keep things simple we will concentrate on the one-dimensional equation

$$x_{t+1} = f(x_t)$$

where $f(x)$ is a nonlinear function and $x$ is a real variable which may be restricted to the positive reals or to some interval of the reals. Such equations are sometimes called **discrete dynamical systems**, because they are the discrete analogs of dynamical systems occurring in physics. The function $f(x)$ is sometimes called a **map** to emphasize that this is a discrete system. Notice that these equations have no inputs and so the complexity will come from the nonlinear behavior of $f(x)$ rather than from an external source. Some complexity can come from the chosen initial condition.

The simplest way to deal with such equations is computation. If $f(x)$ is a reasonable function then one should be able to write a computer program which when given an initial value $x_0$ computes as many values of the sequence $x_1, x_2, \dots$ as one desires. As usual there is a caveat, computers do not compute with *real* real numbers, instead they compute with approximations to real numbers. So a sequence generated by a computer may not be the sequence actually defined by the nonlinear recurrence. Of course, for reasonable functions one hopes that the real and the computed sequences are similar. In some cases this can be proved, but we will not tackle this approximation problem.

Since humans tend to understand pictures better than sequences of numbers, these computer calculations are often presented as graphs. (These are not the same kind of graphs we used in discussing nonnegative matrices.)

The most straightforward graph is a plot of $x_t$ as a function of $t$. This is often called a **time plot**. Figure 10.1 shows a time plot of the quadratic difference equation

$$x_{t+1} = x_t[1 + r(1 - x_t)]$$

with $r = 2.99$ and $x_0 = .35$. To make this plot look more like a continuous function, it is not plotted as a sequence of points but rather each point $(t, x_t)$ is connected by a straight line to the next point $(t + 1, x_{t+1})$. This time plot shows that the values of the sequence are positive and don't get too big, but that they do jump around in an irregular manner.




FIGURE 10.1. A time plot of a chaotic trajectory (quadratic model with r=2.99).

Another plotting technique may be more revealing. This technique is called a **web plot**, and it is somewhat similar to the phase plane plot used in differential equations. The idea is to plot $x_{t+1}$ as a function of $x_t$. But from the difference equation this would mean simply plotting $f(x)$ as a function of $x$. While this would give some information, it does not show a solution sequence. To follow a solution it is useful to have both the *curve* $f(x)$ and the *line* $y = x$ plotted on the same axes. To follow a solution:

1. Start with $(x_0, f(x_0))$ which is a point on the *curve*.
2. Then draw a horizontal line from this point to the *line* $y = x$. The point of intersection on the line is $(f(x_0), f(x_0))$, i.e. $(x_1, x_1)$.
3. Then draw a vertical line from this point to the *curve*. The point of intersection on the curve is $(x_1, f(x_1))$, i.e. $(x_1, x_2)$.

4. Now simply repeat the steps going from *curve* to *line* and *line* to *curve* for as many steps as desired.




FIGURE 10.2. A web plot showing chaos (quadratic model with r=2.99).

Figure 10.2 shows the web plot which corresponds to the time plot shown in Figure 10.1. This web plot may be more informative than the time plot. For example, the parabola $x[1 + r(1 - x)]$ is outlined in the web plot suggesting that many or most values of $x$ are visited on the trajectory, but this plot also suggests that some regions are more frequently visited than others. Further the plot suggests that if the trajectory is periodic then it has a very very long period. Also, the fixed point $x = 1$ is visible, and the plot suggests that this fixed point tends to repel nearby trajectories.

A full understanding of a nonlinear system might require computing solutions for essentially all initial conditions. Since this is clearly infeasible, we want some simple ways to summarize the behaviors. For arbitrary $f(x)$'s this will be impossible, that is, there are $f(x)$'s which are complicated enough to simulate the behavior of Turing machines and so there can be no algorithm which determines if an $f(x)$ given an $x_0$ eventually goes into a cycle. [95] But functions used in various applications tend to be simple enough or have enough reasonable properties to allow some sort of analysis. In particular the assumption that $f(x)$ is continuous and is sufficiently differentiable is usually enough to obtain some results.

We usually focus on results that are invariant or asymptotic. By **invariant property** we mean that a property remains the same along a

trajectory. The most common example of this is a closed physical system in which the total energy is always the same. Even as the system's variables change, the total energy remains invariant. **Asymptotic properties** are those which hold in the limit as $t$, time, increases. For example, a system might approach a "steady state" as time increases. The behavior of a nonlinear system is usually analyzed in terms of the system's fixed points and cycles. The next definitions define some of these properties.

**Definition 10.2.1.** The **iterates** of $f$ are defined recursively by $f^{(0)}(x) = x$ and $f^{(i+1)}(x) = f(f^{(i)}(x))$. A set of *distinct* points $p_1, ..., p_K$ forms a **cycle of length** $K$ (sometimes called a $K$-cycle) iff $f^{(K)}(p_i) = p_i$ for all $1 \leq i \leq K$. The points of a $K$-cycle are called **periodic** whose **period** equals $K$. A fixed point $p$ is **attractive** iff for every small enough neighborhood of $p$ and all points $q$ in this neighborhood, $\lim_{n\to\infty} f^{(n)}(q) = p$. Similarly, a cycle $p_1, ..., p_K$ is an **attractive cycle** iff for every small enough neighborhood of the cycle and all points $q$ in this neighborhood $\lim_{n\to\infty} f^{(nK)}(q) \in \{p_1, ..., p_K\}$. A fixed point or cycle is **repelling** iff for every small enough neighborhood of the point or cycle, and all points $q$ in this neighborhood, there is an $n$ so that $f^{(n)}(q)$ is outside the neighborhood.

In the following sections we will look at some simple examples of nonlinear systems and see how fixed points and cycles can be shown to be attractive.

### *10.2.1 Sarkovskii's Theorem*

Nonlinear systems may have many different co-existing cycles, for example, $x_0, f(x_0), f^{(2)}(x_0)$ may all be distinct and $f^{(3)}(x_0) = x_0$, but say $y_0 = f(y_0)$ then there would be a cycle of length 3 and a cycle of length 1. An **interval map** is a function defined on an interval $[a, b]$ which has all its values in $[a, b]$, and in the special case of continuous one-dimensional interval maps, the following theorem tells which cycles will occur. It does not hold for discontinuous maps or for more than one dimension. While the theorem dates from 1964, it was not known in the U.S. until many years later. So, for example Li and Yorke [85] proved a special case in the 1970's and Cull [26] and Rosenkranz [121] also proved special cases. The proof of this theorem is beyond the scope of this book, but it can be found in [43].

**Theorem 10.2.1 (Sarkovskii's Theorem [125]).** *If $f(x)$ is a continuous one-dimensional interval map which has a cycle of length $K$, then $f(x)$ also*

*has cycles of length $L$ for every $L$ less than $K$ in the following ordering:*

$$
\begin{array}{lll}
3 > 5 > 7 > & \ldots\ldots & > \ldots \\
> 2\cdot 3 > 2\cdot 5 > & \ldots\ldots & > \ldots \\
> 2^2\cdot 3 > 2^2 5 > & \ldots\ldots & > \ldots \\
\quad\vdots & \quad\vdots & \\
> 2^n\cdot 3 > 2^n\cdot 5 > & \ldots\ldots & > \ldots \\
\quad\vdots & \quad\vdots & \\
> 2^n > 2^{n-1} > & \ldots\ldots & > 2^2 > 2 > 1.
\end{array}
$$

Notice, in particular that if $f(x)$ has a cycle of length 3, then $f(x)$ has cycles of every length. Also, if $f(x)$ has any cycles other than fixed points, then $f(x)$ has a cycle of length 2.

## 10.3 Chaos

Since the seminal papers of Li and Yorke [85] and May [96, 97], the importance of chaos in science has been evident. A variety of different meanings have been attached to the word *chaos*. For some, it simply means very complicated looking behavior. For instance, Figures 10.1 and 10.2 show behavior that looks complicated. For others, the **butterfly effect** is the signature of chaos. In poetic terms, the butterfly effect means that the flapping of a butterfly's wings in Borneo may cause a tropical storm in the Caribbean which devastates the sugar crop and leads to the downfall of Communism. In the original use of Li and Yorke [85], chaos meant the coexistence of cycles of every length. Other properties which some people like are that chaotic trajectories come near to every point in a space and that there is a measure that is invariant along trajectories. Detailed consideration of these issues are beyond the scope of this book, but we will consider some nonlinear systems and see that they do have some of these properties

### *10.3.1 A Simple Chaotic System*

Several properties of **chaos** are:

(a) cycles of every length
(b) sensitive dependence on initial conditions
(c) bounded but aperiodic orbits
(d) for every open set $A$ and every open set $B$ there is an $x_0 \in A$ and a $K \in \mathbb{N}$ so that $f^{(K)}(x_0) \in B$.

A very simple difference equation which displays these chaotic properties is

$$x_{t+1} = 2x_t \pmod 1 . \tag{10.1}$$

The values for the $x$'s will be limited to the half open interval $[0,1)$. Two real numbers $x$ and $y$ are **equivalent mod 1** exactly when $x - y$ is a (integer) multiple of 1, that is, when $x$ and $y$ differ by an integer. We symbolize this relationship by $x \equiv y \pmod 1$. For example, $3.14 \equiv .14 \pmod 1$. We restrict the $x$'s to $[0,1)$ because $1 \equiv 0 \pmod 1$ and because every real number is equivalent mod 1 to a number in $[0,1)$. Equivalence mod 1 can be visualized as a clockface with circumference 1, so that 0 is at the top position and all the numbers between 0 and 1 are in their standard positions around the dial. Notice that 1 does *not* appear, because $1 \pmod 1 \equiv 0$. That is, if you had a string of length exactly one and you wrapped it around the clockface, both the start and the end of the string would be at 0. If your string had length $3/2$, wrapping it around the clockface would put the end at $1/2$, which agrees with $3/2 \equiv 1/2 \pmod 1$.

Let's consider some initial conditions for this equation. Clearly if $x_0 = 0$ then $x_t = 0$ for all $t \geq 0$ and the equation has a fixed point. Some points are attracted by 0. For example, if $x_0 = 1/2$, then $x_1 = 2(1/2) = 1 = 0 \pmod 1$. In fact for any integer $k$, $1/2^k$ goes through $1/2^k, 1/2^{k-1}, \ldots, 1/2, 0$. This is rather tame behavior. We can also find periodic behavior. Let's try $x_0 = 2/3$. Then $x_1 = 2(2/3) = 4/3 = 1/3 \pmod 1$, and $x_2 = 2/3$. This is a cycle of length 2. More generally, if $x_0 = \dfrac{2^{k-1}}{2^k - 1}$ then

$$\begin{aligned}
x_1 &= \frac{2^k}{2^k - 1} - 1 = \frac{1}{2^k - 1} \\
x_2 &= \frac{2}{2^k - 1} \\
x_3 &= \frac{4}{2^k - 1} \\
&\vdots \\
x_{k-1} &= \frac{2^{k-2}}{2^k - 1} \quad \text{and} \\
x_k &= \frac{2^{k-1}}{2^k - 1} .
\end{aligned}$$

For any $k$ there is an initial condition which gives a cycle of length $k$.

This behavior can be seen more readily in binary notation. For example, $\frac{2}{3} = .10101010\ldots$. Then multiplying by 2 and reducing mod 1 corresponds to shifting the binary point one place to the right and dropping any bit to the left of the binary point. So $2(\frac{2}{3}) \pmod 1$ would be computed by

taking .10101010..., shifting the binary point to get 1.0101010..., and dropping the 1 to the left of the binary point to get .0101010..., which is $\frac{1}{4}+\frac{1}{16}+\frac{1}{64}+\cdots=\frac{1}{4}(1+\frac{1}{4}+\frac{1}{4^2}+\ldots)$ and taking the sum of the geometric series, we get $\frac{1}{4}(\frac{1}{1-\frac{1}{4}})=\frac{1}{3}$. Similarly, starting with .0101010..., shifting the binary point and dropping the leading 0, we get .10101010... which is the number $\frac{2}{3}$ we started with two steps ago. Here, we can *see* the period 2 of the solution to the difference equation follows from the period 2 of the binary expansion of $\frac{2}{3}$.

**Sensitive dependence on initial conditions** is also easy to demonstrate. Since 0 is a fixed point of the equation, if we choose 0 as the initial condition, then the solution will remain at 0 forever thereafter. Now consider an initial condition $\epsilon > 0$ which is very close to 0. Here the solution will start at $\epsilon$, go to $2\epsilon$ to $4\epsilon$, and so forth. If $\epsilon$ is very small then $2\epsilon$ and $4\epsilon$ will also be very small. So for a small number of steps the two solutions – the one with initial condition 0 and the one with intitial condition $\epsilon$ – will both be nearly 0 and we won't see any divergence. But since $\epsilon > 0$, there is at least one 1 in the binary expansion of $\epsilon$. Say the first 1 appears as the $k^{th}$ bit. Then after $k-1$ steps the 1 will have moved to the first bit position and $x_{k-1} \geq 1/2$ for the solution with $\epsilon$ as the initial condition. On the other hand, $x_{k-1}$ will still be 0 for the solution with 0 as the initial condition. So the 2 solutions are now significantly different. The difference is roughly $2^k\epsilon$ and this is much bigger than $\epsilon$.

Another way to describe sensitive dependence is in terms of information. How many bits of the initial condition will we need to know to accurately predict $x_k$? For our example equation (10.1) we need about $k$ bits to predict $x_k$. So if we could only measure the initial condition to 10 correct bits, we would be at a loss to predict the value of the solution after 11 steps. Sometimes this is explained as a loss of bits. If we know the initial condition to 10 correct bits and lose a bit of accuracy at each step, then we will have no bits of accuracy left to predict the $11^{th}$ value.

While our example is extreme for showing sensitive dependence, such dependence can arise in more realistic situations. Consider predicting the weather. One might measure a number of variables like pressure, temperature, and wind direction to 10 bit accuracy (about one part in a thousand). Now assume that we are calculating the weather with a time step of 1/1000 of a day. To compute tomorrow's weather we need to calculate about 1000 steps. If the weather conditions are *smooth* we don't lose many bits per step. When we lose 1/1000 bits per step in smooth conditions, we lose about 1 bit in tomorrow's forecast and we expect that our prediction will be accurate to about 9 bits. On the other hand if in *turbulent* conditions we lose about 1/100 bits per step, then in 1000 steps we will have lost about 10 bits and so we will have no bits of accuracy left when we make tomorrow's prediction. This is not just a story, since some of the original work on chaos was an attempt to describe why weather prediction is so difficult.

Returning to our main theme of exhibiting sensitive dependence on initial conditions, let us note that there was nothing really special about looking at 0 as one of the initial conditions. If we look at any initial conditions say $x_0$ and $x_0 + \epsilon$ then $x_k(x_0)$ will differ by about $1/2$ from $x_k(x_0 + \epsilon)$ when $k$ is about $-\log \epsilon$.

Another characteristic of chaos is bounded but aperiodic orbits. Clearly in our example, all orbits are limited to $[0, 1)$, and so the orbits are always bounded. Now we want to choose some initial condition $x_0$, and show that $x_0, x_1, x_2, \ldots$ is an aperiodic sequence. But the interpretation of the equation as a shift on binary sequences makes this problem easy. All we need to do is find an aperiodic binary sequence. For example $.10100100010000\ldots$ is aperiodic because the number of 0's between consecutive 1's is increasing. Using this number as an initial condition, the solution starts above $1/2$, drops to a lower value, goes above $1/2$, drops to a lower value, builds up for 2 steps, goes above $1/2$, drops to a lower value, builds up for a few steps, goes above $1/2$, and so forth. Why do we write this initial condition as a binary sequence rather than as a ratio? Because no such ratio exists. That is, the number we have written is *not* a rational number because it has a non-periodic expansion.

Next, given two open sets $A$ and $B$ in $[0, 1)$ we want to show that there is an initial condition in $A$ that will lead to a solution that visits $B$. Because $A$ is open, $A$ contains an open interval $(a - 2^{-j}, a + 2^{-j})$. Let $b \in B$. We want to find an element in $A$'s open interval which is eventually mapped to $b$. Let $c = .a_1 a_2 \ldots a_j$ be the number formed from the first $j$ bits in the binary expansion of $a$. Let $K = j + 1$ and $x_0 = c + 2^{-K} b$. We see that $x_0 \in A$ because $a - 2^{-j} < c \leq a$. For this choice of $x_0$, $f^{(K)}(x_0) = b$.

A similar construction yields a trajectory that visits every open set. (In what follows we refer to a open interval with two rational endpoints as a "rational interval.") Since every open set contains a rational interval, a trajectory can be shown to visit every open set by showing it visits every rational interval. The rational numbers are **countable**, that is, they can be put into one to one correspondence with the natural numbers. There are many such correspondences, and any of them gives an ordering of the rationals. So we can speak of the first rational, the second rational and so forth. But since we have just mapped the rationals to the natural numbers, we can map the upper and lower endpoints for a rational interval (and so the interval itself) to a pair of natural numbers, which can be considered to be a rational number. Hence, the set of rational intervals can be countably ordered.

Next, since every rational interval contains a rational $r$ with a terminating binary expansion, there is a natural number $j$ so that $(r - 2^{-j}, r + 2^{-j})$ is inside the interval. So, if our trajectory visits each of these special intervals, it will visit all rational intervals, and hence all open sets. We now pick an initial condition $x_0$ at which to start our trajectory which visits every open set. For the sequence $< r_i >$ of terminating rationals, ordered

according to the ordering of rational intervals, define the initial condition $x_0$ as

$$x_0 = .r_1 \quad (j_1+1)0\text{'s} \quad r_2 \quad (j_2+1)0\text{'s} \quad r_3 \quad (j_3+1)0\text{'s} \quad \ldots .$$

So, $x_0 \in (r_1, r_1 + 2^{-j_1})$ and after an appropriate number of iterations $x_n \in (r_i, r_i + 2^{-j_i})$, which means the trajectory starting at this $x_0$ will eventually visit every open set.

## 10.4 Local Stability

### *10.4.1 Local Stability of a Fixed Point*

If $x_0$ is a fixed point of $f(x)$ we would like to know if nearby values of $x$ are attracted toward $x_0$. The following definition and subsequent theorem make this idea more precise.

**Definition 10.4.1.** A fixed point $x_0$ of $f(x)$ is **locally stable** if for every $y$ in every small enough neighborhood of $x_0$, $f(y)$ is still in the neighborhood and $\lim_{K\to\infty} f^{(K)}(y) = x_0$.

Testing whether a point is locally stable may seem difficult, but if $f(x)$ is differentiable then there is a reasonably easy test which is given in the next theorem.

**Theorem 10.4.1.** *Let $x_0$ be a fixed point of $f(x)$ and assume that $f(x)$ is differentiable at $x_0$. Then $x_0$ is locally stable if $|f'(x_0)| < 1$, and if $x_0$ is locally stable then $|f'(x_0)| \leq 1$.*

*Proof.* If $f(x)$ is differentiable then there is a constant $\alpha$ so that for all $y$ close enough to $x_0$,

$$\frac{|f(y) - f(x_0)|}{|y - x_0|} \leq |f'(x_0)| + \alpha|y - x_0|.$$

If $|f'(x_0)| < 1$, then there is a $\delta > 0$ so that for all $y$ very close to $x_0$, the right hand side of this inequality is less than $1 - \delta$. Since $x_0$ is a fixed point of $f(x)$, this gives

$$|f(y) - x_0| \leq (1 - \delta)|y - x_0|$$

and so $f(y)$ is in the same neighborhood as $y$. Further, letting $d_K = |f^{(K)}(y) - x_0|$ gives the equation $d_K \leq (1-\delta)d_{K-1}$. Then $d_K \leq (1-\delta)^K d_0$ and $\lim_{K\to\infty} d_K = 0$. Therefore, $x_0$ is locally stable. Conversely, if $x_0$ is locally stable then

$$|f(y) - x_0| = |f(y) - f(x_0)| \leq |y - x_0|$$

and since $f(x)$ is differentiable

$$|f'(x_0)| = \lim_{y \to x_0} \frac{|f(y) - f(x_0)|}{|y - x_0|} \leq 1.$$

□

The basic idea here is that close enough to a fixed point, we can approximate a nonlinear difference equation by a linear difference equation. That is, we take the nonlinear equation $x_{t+1} = f(x_t)$ near the fixed point $E$, then define a new variable $d_K = |x_K - E|$, and consider the linear equation $d_K = (1 - \delta)d_{K-1}$. We have translated the fixed point $E$ for $x$ to the fixed point 0 for $d$. If $f(x)$ is differentiable with $|f'(x)| < 1$, then the linear equation gives an upper bound on the the behavior of $|x_K - E|$ in a small enough neighborhood of $E$.

Linear approximations can also be made around non-fixed points, but we will not get as much useful information. When we use a Taylor expansion for $f(x)$ around $E$, then close to $E$, the first term of the expansion, i.e. $f(E)$, will dominate. If $E$ is a fixed point then $f(x) - E$ will have an expansion in which the first term is 0, and so the second term $f'(E)\ (x - E)$ will dominate.

Notice that this theorem has a small hole, it does not say what happens when $|f'(x_0)| = 1$. Essentially this is because we are making a linear approximation to $f(x)$ and this linear approximation dominates the nonlinear behavior locally if $|f'(x_0)| < 1$, but the nonlinear terms are necessary to determine behavior when $|f'(x_0)| = 1$. As a simple example, consider

$$x_{t+1} = f(x_t) = x_t[1 + r(1 - x_t)].$$

This system has a fixed point at $x = 1$. The derivative is $f'(x) = 1 + r - 2rx$, and so $|f'(1)| \leq 1$ if $0 \leq r \leq 2$. By the theorem, $x = 1$ is locally stable when $0 < r < 2$. Let's see what happens in the special cases when $r = 0$ and $r = 2$, that is, when $|f'(1)| = 1$. When $r = 0$, $f(x) = x$, so if the system starts near 1, it stays near 1, but the iterates do not converge to 1. On the other hand when $r = 2$, $f(x) = 1 - (x - 1) - 2(x - 1)^2$ and starting at $1 - \epsilon$ will give $1 + \epsilon - 2\epsilon^2$ which is in the same $\epsilon$ neighborhood of 1. But starting at $1 + \epsilon$ will give $1 - \epsilon - 2\epsilon^2$ which is not in the same $\epsilon$ neighborhood of 1. In spite of this, $f(f(x))$ for $x$ near 1 will always be closer to 1, and $\lim_{K \to \infty} f^{(K)}(x) = 1$ for all $x$ in $(0, 3/2)$. This suggests that our definition of "local stability" may not be ideal, and the interested reader should consult the literature for other definitions of "local stability".

### *10.4.2 Local Stability of a Cycle*

Stability for a cycle is similar to stability for a point. The system

$$x_{t+1} = f(x_t)$$

has a **cycle** $x_1, x_2, \ldots, x_K$ **of length** $K$ if $x_{i+1} = f(x_i)$ for $i \in \{1, \ldots, K-1\}$ and $x_1 = f(x_K)$. It is convenient to consider the $K$-fold iteration of $f$, because each of the points $x_1, x_2, \ldots, x_K$ is a fixed point of $f^{(K)}(x)$, that is, $f^{(K)}(x_i) = x_i$. Since we know when $x_i$ is a locally stable fixed point, we would like to say that $x_1, x_2, \ldots, x_K$ is a **locally stable cycle** when each of the points $x_1, x_2, \ldots, x_K$ is a locally stable fixed point of $f^{(K)}(x)$. To make this simpler, we would like to consider only one point, but we might worry that one of these points could be locally stable while some of the other points would not be locally stable. Luckily, if we assume that $f(x)$ is differentiable then these worries vanish because the stability condition at one point is the same as the stability condition at each of the other points. A fixed point $x_i$ is a locally stable fixed point of $f^{(K)}(x)$ if the absolute value of the derivative of $f^{(K)}(x)$ is less than 1 at $x_i$. To calculate the derivative, we use the Chain Rule

$$D[f^{(K)}(x)] = f'(f^{(K-1)}(x))D[f^{(K-1)}(x)]$$

and so

$$D[f^{(K)}(x)]|_{x=x_1} = f'(x_K)f'(x_{K-1})\ldots f'(x_1).$$

We notice that this is the value of the derivative at *every* point in the cycle, because taking any other point will only result in the terms in the product appearing in different order.

As an example, let us consider

$$x_{t+1} = f(x_t) = x_t[1 + r(1 - x_t)]$$

with $r$ slightly larger than 2. As in a previous example, this system has a fixed point at $x = 1$ but because $r > 2$, this fixed point is repelling. We want to show that there is a pair of points $x_1$ and $x_2$ so that $f(x_1) = x_2$ and $f(x_2) = x_1$ and that the cycle consisting of $x_1$ and $x_2$ is locally stable.

The condition for a cycle of length 2 is

$$f(f(x)) = x = x[1 + r(1 - x)][1 + r(1 - f(x))].$$

We can eliminate the fixed point at $x = 0$ by dividing this equation by $x$. Simplifying the resulting equation by subtracting 1 and dividing by $r$ gives

$$2(1 - x) - rx(1 - x) + r(1 - x)(1 - f(x)) = 0.$$

Now, the fixed point at $x = 1$ can be eliminated by dividing by $1 - x$, and after expanding the remaining $f$, this gives the quadratic equation

$$2 + r - r(2 + r)x + r^2x^2 = 0,$$

which will have two distinct real roots since $r > 2$. (This can be seen by calculating the discriminant of this polynomial.) We could calculate these roots, $x_1$ and $x_2$, but this polynomial already tells us that

$$x_1x_2 = \frac{2 + r}{r^2} \quad \text{and} \quad x_1 + x_2 = \frac{2 + r}{r}. \tag{10.2}$$

Remember that the sufficient condition for local stability of the cycle is

$$|f'(x_1)f'(x_2)| < 1.$$

It is easy to compute

$$f'(x) = 1 + r - 2rx$$

and

$$f'(x_1)f'(x_2) = (1+r)^2 - 2r(1+r)(x_1+x_2) + 4r^2x_1x_2.$$

Using the formulas from (10.2)for $x_1x_2$ and $x_1 + x_2$ gives

$$f'(x_1)f'(x_2) = (1+r)^2 - 2(1+r)(2+r) + 4(2+r) = 5 - r^2$$

and the local stability condition becomes

$$2 < r < \sqrt{6}.$$

Hence, this system has a locally stable 2-cycle when $r$ satisfies this inequality.

### *10.4.3 Local Stability in Two Dimensions*

We can use the techniques of the previous subsections to look at the local stability of a nonlinear system in more than one dimension. The only differences are that we need to generalize the idea of neighborhood and the idea of derivative. A neighborhood in one dimension is an open interval $(x - \epsilon, x + \epsilon)$. For a **neighborhood** in more than one dimension, we can use a ball of radius $\epsilon$. So an $\epsilon$-neighborhood of a point $x$ would be the set of all points $y$ such that $|x - y| < \epsilon$ where we interpret the **absolute value** to mean distance in the appropriate dimension.

When we think of a linear approximation to an $n$-dimensional function the approximation should consist of $n$ linear functions, one for each dimension. Specifically, if

$$F(x,y) = \begin{pmatrix} f_1(x,y) \\ f_2(x,y) \end{pmatrix}$$

is a two-dimensional function then our analog of the derivative is

$$J(x,y) = \begin{bmatrix} \frac{\partial f_1}{\partial x} & \frac{\partial f_1}{\partial y} \\ \frac{\partial f_2}{\partial x} & \frac{\partial f_2}{\partial y} \end{bmatrix}.$$

This is called the **Jacobian matrix** which is also important in vector calculus. Of course, the partial derivatives inside this matrix are functions and have to be evaluated at the point $(x, y)$ at which we are making a linear approximation. For example,

| | |
|---|---|
| if | $f_1(x,y) = xy$ |
| and | $f_2(x,y) = x + y^2,$ |

then the Jacobian matrix is

$$\begin{bmatrix} y & x \\ 1 & 2y \end{bmatrix}.$$

If we want a linear approximation near the point $(0,0)$, then the Jacobian matrix evaluated at $(0,0)$ becomes

$$\begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$$

and the linear approximation at $(0,0)$ is

$$\begin{pmatrix} f_1(x,y) \\ f_2(x,y) \end{pmatrix} \approx \begin{pmatrix} f_1(0,0) \\ f_2(0,0) \end{pmatrix} + \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ x \end{pmatrix}.$$

For the linear approximation at $(1,0)$ we would have

$$\begin{pmatrix} f_1(x,y) \\ f_2(x,y) \end{pmatrix} \approx \begin{pmatrix} f_1(1,0) \\ f_2(1,0) \end{pmatrix} + \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{pmatrix} x-1 \\ y-0 \end{pmatrix} = \begin{pmatrix} y \\ x \end{pmatrix}.$$

As this example shows, we still have a problem because the one-dimensional derivative was a single number, but the Jacobian matrix is an array of numbers. The idea in one dimension was that around a fixed point the linear approximation can be viewed as a linear difference equation and the solutions of this equation will converge to the fixed point if the derivative is less than one in absolute value because the derivative is also an eigenvalue of the difference equation. Similarly the eigenvalues of the Jacobian matrix will be the eigenvalues of a matrix linear difference equation around a fixed point, and the solutions will converge to the fixed point if all the eigenvalues are less than one in absolute value. This result generalizes to all dimensions as the following theorem states. (For a proof, refer to [81].)

**Theorem 10.4.2.** *If $F$ is an $n$-dimensional function with fixed point $X$, and $J$ is the Jacobian matrix of $F$ evaluated at $X$, then $X$ is a locally stable fixed point if all eigenvalues of $J$ have absolute value less than 1. If at least one of these absolute values is strictly greater than 1, the fixed point is unstable.*

As an example, consider the system

$$\begin{aligned} x_{t+1} &= \sin(y_t) \\ y_{t+1} &= \cos(x_t) \end{aligned}$$

whose Jacobian matrix is

$$\begin{bmatrix} 0 & \cos(y) \\ -\sin(x) & 0 \end{bmatrix},$$

its characteristic polynomial is $\lambda^2 + \sin x \cos y$ and its eigenvalues are $\pm\sqrt{-\sin x \cos y}$. Depending on the values of $x$ and $y$, these roots may be

real or complex but we know that $|\sin x| \leq 1$ and $|\cos y| \leq 1$, so these roots will have absolute value $\leq 1$ regardless of whether they are real or complex. More strongly, if there is a fixed point at $(x, y)$ then $|x| \leq 1$ and $|y| \leq 1$, but then $|\sin x| < 1$ and $|\cos y| \leq 1$, so the eigenvalues are less than 1 in absolute value. Therefore, all fixed points are locally stable.

The fixed points of this system can be approximated by plotting $\cos x$ along the $x$-axis and plotting $\sin y$ along the $y$-axis. The fixed point will be where these two curves meet. One can see that there will only be one fixed point and it will occur at a point where $0 < x < 1$ and $0 < y < 1$. The above analysis shows that the eigenvalues for this fixed point will be less than 1 in absolute value and so this fixed point is locally stable.

Another simple example of a two-dimensional system is

$$\begin{aligned} x_{t+1} &= x_t + x_t(1 - x_t - y_t)/6 \\ y_{t+1} &= y_t(1 + x_t - y_t). \end{aligned}$$

For fixed points $p = (x, y)$ of this system,

$$\begin{aligned} x &= x + x(1 - x - y)/6 \\ y &= y(1 + x - y) \end{aligned}$$

and simplifying these gives

$$\begin{aligned} 0 &= x(1 - x - y) \\ 0 &= y(x - y), \end{aligned}$$

which has three solutions:

$$\begin{aligned} x = 0 \quad & y = 0; \\ x = 1 \quad & y = 0; \\ x = 1/2 \quad & y = 1/2. \end{aligned}$$

The matrix of partial derivatives for this system is

$$J(x, y) = \begin{bmatrix} 1 + (1 - x - y)/6 - x/6 & -x/6 \\ y & 1 + x - 2y \end{bmatrix}.$$

The Jacobian matrix for the fixed point $x = 0$, $y = 0$ is

$$\begin{bmatrix} 7/6 & 0 \\ 0 & 1 \end{bmatrix},$$

which has an eigenvalue $7/6$, which is larger than 1, and so this fixed point is unstable. The Jacobian matrix for the fixed point $x = 1$, $y = 0$ is

$$\begin{bmatrix} 5/6 & -1/6 \\ 0 & 2 \end{bmatrix},$$

and again there is an eigenvalue 2 larger than 1, and so this fixed point is unstable. Finally, the fixed point $x = 1/2$, $y = 1/2$ has the Jacobian matrix

$$\begin{bmatrix} 11/12 & -1/12 \\ 1/2 & 1/2 \end{bmatrix},$$

whose characteristic polynomial is $(11/12 - \lambda)(1/2 - \lambda) + 1/12 \cdot 1/2$, which simplifies to $\lambda^2 - \frac{17}{12}\lambda + \frac{1}{2}$. The eigenvalues are $2/3$ and $3/4$, and so this fixed point is locally stable. For this system, one could make the reasonable guess that if the system were started at any point that is not a fixed point, then the trajectory will eventually approach the fixed point $x = 1/2$, $y = 1/2$. While this guess is reasonable, one would still need to rule out other possible behaviors. For example, there could be cycles or aperiodic orbits which simply don't show up in a fixed point analysis.

## 10.5 Global Stability

A locally stable point attracts trajectories that are within a small neighborhood of the point, but one is usually interested in larger neighborhoods. At the extreme, such a neighborhood could be the whole space. This suggests the definition:

**Definition 10.5.1.** If $p$ is a fixed point of $x_{t+1} = f(x_t)$ and $\lim_{K\to\infty} f^{(K)}(b) = p$, for each initial condition $b$ in a set $B$, then $p$ is **globally stable** for $f(x)$ and $B$.

Usually the set $B$ is not mentioned since by context one sees that $B$ is the reals, or the positive reals, or the interval $[0, 1]$, or some other reasonable set. One would also like to say that a cycle or other invariant set is globally stable but this is difficult since fixed points and the special trajectories that lead to those fixed points might not converge to the cycle. People usually solve this problem by saying that the cycle is globally stable without mentioning the *relatively* few points that do not lead to the cycle.

Unlike local stability, there is no nice characterization of global stability even if the function is differentiable. The classical technique to show global stability is to find a "Liapunov" or "energy" function, show that this function is nonnegative and equal to 0 only at the fixed point, and that the "energy" decreases along each trajectory. (See LaSalle [81] for details.) Unfortunately finding a Liapunov function and showing that it has the required properties is a formidable task. In Section 10.6 we will show that some simple functions can be used to prove global stability. Of course, in more than one dimension global stability will be harder to deal with.

We next show how the local stability technique can be slightly extended to prove global stability. For this we use

**Theorem 10.5.1.** *If $f(x)$ is a nonnegative continuous function defined on the real numbers with $x < f(x) < 2 - x$ on $(0,1)$ and $x > f(x) > 2 - x$ on $(1,2)$ then $x = 1$ is globally stable for $(0, \infty)$.*

We will not prove this result, but simply consider the example

$$x_{t+1} = x_t e^{2(1-x_t)} = f(x_t)\,.$$




FIGURE 10.3. Showing global stability for a simple model.

Figure 10.3 shows $f(x) = xe^{2(1-x)}$ and the line $y = 2 - x$. To see what will happen for the iteration of $f$, follow the "web" procedure. Pick a starting $x_0 < 1$ which is also above the peak of $f(x)$. Now go up to the curve, over to the line $y = x$, down to the curve, and then back over to the line $y = x$. At this point you will have reached an $x$-coordinate which is between $x_0$ and 1, and this will be true regardless of which such $x_0$ you chose. So, the iterates will spiral around and toward the point $x = 1$, and this will be the limit of $< x_k >$. Notice that if you began with an $x_0$ lying outside the prescribed interval, then you will eventually find an iterate inside the interval and from there the iterates will follow the spiral convergence which we just described.

How does the line $y = 2 - x$ enter into this discussion? The fact that $f(x)$ lies between the lines $y = 2 - x$ and $y = x$ allows one to show that $|x_k - 1|$ is a decreasing function of $k$. This implies that either $< x_k >$ converges to 1 in a finite number of steps or that there are two convergent subsequences of $< x_k >$–one with increasing elements less than 1 and the

other decreasing with elements greater than 1. Finally, the continuity of $f$ can be used to show that the limit of these subsequences can be neither less than or greater than 1. From this one concludes that the limit of the sequence must equal 1. The unimodal shape of our example is *not* essential to this convergence, but it allows for easier visualization.

In this example we see that convergence to 1 can be shown by verifying that 0 and 1 are the only fixed points of the continuous function $f^{(2)}(x)$(refer to Exercise 10.1). This means that if $< f^{(2k)}(x) >$ converges, it must converge to one of the fixed points. The fact that $f(x)$ is bounded by the lines $y = 2 - x$ and $y = x$ can be used to show that 0 is repelling, 1 is attractive, and $< f^{(2k)}(x) >$ is convergent. This shows that $< x_k >$ converges to 1 for $x_0 \in (0, 2)$. Then the fact that $f(x) > 0$ on $(0, \infty)$ is used to argue that an iterate must eventually visit the interval $(0, 2)$ and thus $< x_k >$ converges to 1 for any initial positive initial value.

## 10.6 Linear Fractional Recurrences

The simplest nonlinear recurrences are the linear fractional recurrences, which are simply the ratio of two linear recurrences. These recurrences are worth studying because

(a) they are simple;
(b) they will give examples of how nonlinear recurrences can work;
(c) techniques from linear recurrences can be used;
(d) linear fractionals can be used to study other nonlinear recurrences.

We will write a **linear fractional recurrence** in the form

$$x_{t+1} = \frac{ax_t + b}{cx_t + d}$$

where we assume that $a, b, c, d$ are real constants and that $x_t$ is a real variable and later we will consider the special case when the constants and variables are restricted to be rational numbers. Linear fractional recurrences are a special subclass of the general recurrence

$$x_{t+1} = f(x_t)$$

which is easy to analyze when $f$ is a linear function but is impossible to analyze when $f$ is a general function. By restricting $f$ to linear fractional form, we will have nonlinear recurrences which we can analyze. The usual questions about nonlinear systems include the existence of fixed points, the existence of cycles of various lengths, the asymptotic behavior of the system (i.e. are the fixed points or cycles attractive in some sense), local and global stability of fixed points and cycles, chaos or chaotic-like behavior, average behavior for a distribution of initial conditions. Pleasantly enough,

all of these questions can be solved in a relatively easy fashion for linear fractional systems.

We would like to analyze linear fractional systems in terms of fixed points, cycles, and other nonlinear behavior. But first we should note that a linear fractional system degenerates into a linear one when $c = 0$ since

$$x_{t+1} = \frac{\hat{a}x_t + \hat{b}}{0x_t + \hat{d}} = ax_t + b. \tag{10.3}$$

As we've already seen, the solution to such a linear recurrence is

$$x_t = a^t x_0 + b\sum_{i=0}^{t-1} a^i = \begin{cases} x_0 + bt & \text{if } a = 1 \\ a^t x_0 + b\frac{a^t-1}{a-1} & \text{if } a \neq 1 \end{cases}. \tag{10.4}$$

It is usual to analyze such a system in terms of the growth rate $a$. Generally, when $|a| > 1$ solutions will be increasing in absolute value, while when $|a| < 1$ solutions will be decreasing in absolute value. In fact by setting $y = x + \frac{b}{a-1}$ the linear system (10.3) can be rewritten as $y_{t+1} = ay_t$ and the analysis in terms of growth rate seems perfectly appropriate. Of course, this transformation does not make sense when $a = 1$, since in the original (10.3) all solutions diverge while in the transformed system every point is a fixed point.

If we analyze (10.3) as a nonlinear system we get a different picture. First, there is a fixed point at $-b/(a-1)$, and this point is attractive when $|a| < 1$ for all real $x_0$. To treat the increasing behavior when $|a| > 1$, we extend the real numbers by including $\infty$, the point at plus and minus infinity, and the linear system will always have $\infty$ as a fixed point. This fixed point will be attractive when $|a| > 1$. Considering $\infty$ when $a = 1$ also explains the discrepancy in the transformed system because then each point is transformed to $\infty$ and is thus a fixed point. In summary, a linear system can be seen to have two fixed points, one at $-b/(a-1)$ and the other at $\infty$. One of these will be attractive depending on whether $|a| < 1$ or $|a| > 1$. In the singular case when $a = 1$, the two fixed points degenerate into a single fixed point at $\infty$, which is an attractive fixed point. When one of the two fixed points is attractive, the system exponentially converges toward the fixed point, but when there is only one fixed point the convergence is only linear.

Periodic behavior occurs when $a = -1$ since all non-fixed points have period 2, and the 2-cycles are not attractive. They are sometimes called **neutrally stable** because if the system starts in a point near a cycle, the system always stays near the cycle, but the system does not approach any nearer to the cycle. In the very special case when $a = 1$ and $b = 0$, every point becomes a neutrally stable fixed point. Finally, in the extremely singular case $a = 0$, all trajectories go to $b$ in one step, and $b$ is called a **superstable point**.

Behavior identical to the linear system (10.3) can also be found in the nonlinear system

$$z_{t+1} = \frac{z_t}{bz_t + a}.$$

The only difference is that $x$ has been replaced by $1/x$, and so the fixed points have been shifted from $-b/(a-1)$ and $\infty$ to $(a-1)/-b$ and 0. This observation suggests that linear fractional systems may have many features in common with linear ones, but that their analysis will depend on functions of several parameters and that these functions should be invariant to such obvious transformations as taking reciprocals of the system's variable.

### *10.6.1 Asymptotic Behavior*

Let us now assume that $c \neq 0$ which makes the system

$$x_{t+1} = \frac{ax_t + b}{cx_t + d}$$

nonlinear. A key parameter is the **determinant**, $Det = ad - bc$, and we first consider the systems with $Det = 0$. Multiplying top and bottom of $f(x)$ by $cd$ gives

$$f(x) = \frac{adcx + bcd}{cd(cx + d)} = \frac{bc(cx + d)}{cd(cx + d)} = \frac{b}{d}.$$

So, after one iteration all trajectories go to $b/d$ which is a superstable point. (The superstable point is $a/c$ if $d = 0$.) Therefore, we may assume $Det$ is nonzero.

Next notice that a linear fractional system can be represented in a linear fashion by

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{pmatrix} x_0 \\ 1 \end{pmatrix}$$

where $x_0$ is the initial point since the ratio of the components of the produced vector gives the value of $x_1$. So $x_n$ can be computed by performing the linear iteration

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix}^n \begin{pmatrix} x_0 \\ 1 \end{pmatrix}$$

and then taking a ratio of components. The eigenvalues of the matrix will be called the **eigenvalues of the linear fractional system**, and they are the roots of $\lambda^2 - (a+d)\lambda + ad - bc = 0$.

A fixed point $p$ will occur when $f(p) = p$, or equivalently when $0 = cp^2 + (d-a)p - b$. This quadratic equation will have two distinct real roots when the **discriminant** $Disc = (a-d)^2 + 4bc$ is positive. This is also the discriminant of the characteristic equation. Assuming that $Disc$ is positive, the eigenvalues are distinct and this means that powers of a matrix can be

computed by diagonalizing the matrix, and the $n^{th}$ linear iteration can be written as

$$\frac{b}{\lambda_2-\lambda_1}\begin{bmatrix} 1 & 1 \\ \frac{\lambda_1-a}{b} & \frac{\lambda_2-a}{b} \end{bmatrix}\begin{bmatrix} \lambda_1^n & 0 \\ 0 & \lambda_2^n \end{bmatrix}\begin{bmatrix} \frac{\lambda_2-a}{b} & -1 \\ -\frac{\lambda_1-a}{b} & 1 \end{bmatrix}\begin{pmatrix} x_0 \\ 1 \end{pmatrix}.$$

If we assume that $|\lambda_1| > |\lambda_2|$ then taking a ratio and a limit gives

$$\lim_{n\to\infty} x_n = \frac{b}{\lambda_1 - a},$$

unless $x_0 = b/(\lambda_2 - a)$. It is easy to check that these two points are in fact the fixed points, and so the linear fractional system will have one unstable fixed point and one globally attractive fixed point when $Disc > 0$. (The given formulas are indeterminate when $b = 0$, but then they can be written in the equivalent forms $(\lambda_1 - d)/c$ and $(\lambda_2 - d)/c$.)

A more interesting case occurs when $Disc$ is zero. When $Det < 0$, then $Disc = (a+d)^2 - 4Det$ implies that $a + d = 0$. A simple calculation shows that $f(f(x)) = x$ which means that every point except for the two fixed points will have period 2. Of course, neither of the fixed points is attractive. For $Disc = 0$ and $Det > 0$, there is only one fixed point. Geometrically this corresponds to the line $y = x$ being tangent to the curve $y = f(x)$. The point of tangency is the fixed point. It is easy to see that every point above the fixed point iterates to a point still above but nearer to the fixed point. While a point below the fixed point will iterate away from the fixed point, eventually one of the iterates will jump across the discontinuity and then jump to a point above the fixed point.

As an example, consider $f(x) = x/(x+1)$. Here, $x_n = x_0/(nx_0 + 1)$ and clearly every trajectory will converge to the fixed point at 0. The jumping between branches is hidden by this formula, but can be displayed by following, for example, the trajectory starting at $-3/4$ which gives $-3/4, -3, 3/4$ and then converges down to 0. Notice that convergence is different in the one and two fixed point cases. For two fixed points, $x_n = F.P. + O(\gamma^n)$ where $F.P.$ is the stable fixed point and $|\gamma| < 1$, while in the one fixed point case $x_n = F.P. + O(1/n)$. Figure 10.4 shows the geometry of the two fixed point case and Figure 10.5 shows the geometry of the one fixed point case.

So far, linear fractional systems have not behaved very differently from linear systems. The remaining case, in which $Disc$ is negative (and so $4Det = (a+d)^2 - Disc$ is positive) will display more nonlinear behavior. We will discuss these cases in the following sections.

### *10.6.2 Rational Coefficients and Periodicity*

Like other nonlinear systems, linear fractional systems can display periodic behavior. But unlike other systems, linear fractionals do not allow the coexistence of cycles of different lengths.




FIGURE 10.4. A plot showing a linear fractional system with two fixed points. All iterates will converge to the upper fixed point.

**Theorem 10.6.1.** *If for a linear fractional map $f$ there exists a point $x$ so that $f^{(K)}(x) = x$ then either $K = 1$ (and $x$ is a fixed point) or for all $y$, $f^{(K)}(y) = y$ (all points are periodic).*

*Proof.* As above, the linear fractional map $f(x) = \dfrac{ax+b}{cx+d}$ can be considered as a $2 \times 2$ matrix $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ acting on a vector $\begin{pmatrix} x \\ 1 \end{pmatrix}$. Since $A$ is a $2 \times 2$ matrix, its characteristic polynomial is quadratic and for every natural number $K$, there exist scalars $\alpha_K$ and $\beta_K$ such that $A^K = \alpha_K A + \beta_K I$, and

$$A^K \begin{pmatrix} x \\ 1 \end{pmatrix} = \alpha_K A \begin{pmatrix} x \\ 1 \end{pmatrix} + \beta_K \begin{pmatrix} x \\ 1 \end{pmatrix}.$$

So if $f^{(K)}(x) = x$,

$$\frac{\alpha_K(ax+b) + \beta_K x}{\alpha_K(cx+d) + \beta_K} = x.$$

There are two possibilities for this equation:

(a) $\alpha_K = 0$ and so $A^K = \beta_K I$ and for every $y$, $A^K \begin{pmatrix} y \\ 1 \end{pmatrix} = \beta_K \begin{pmatrix} y \\ 1 \end{pmatrix}$ which is equivalent to $\beta_K y/\beta_K = y$, so all points have period $K$, or

(b) $\alpha_K \neq 0$ and then $ax + b = (cx + d)x$ which is the equation for fixed points.

□




FIGURE 10.5. A plot showing a linear fractional system with one fixed point. Convergence to the fixed point will be from above.

This theorem allows for the co-existence of periodic points and fixed points. When the linear fractional system has real eigenvalues, two real fixed points will occur and periodic behavior will occur when the two eigenvalues have the same magnitude. In this case, all non-fixed points will have period 2.

When the eigenvalues are complex, there will be no real fixed points, and the magnitudes of the eigenvalues are forced to be equal. Here, let $e^{i\theta} = \lambda_1/\lambda_2$. If $\theta$ is a rational multiple of $\pi$, it is a root of unity and there will be a least positive integer $K$ so that $A^K = I$. In this case, every point will be periodic and have period $K$. If $\theta$ is not a rational multiple of $\pi$, the system will not be periodic and all points will fail to be periodic. We will consider this situation in more detail in the next section.

In general, all values of $\theta$ are possible, since a direct calculation of $\lambda_1/\lambda_2$ shows that any desired complex number of norm 1 can be produced by appropriate choice of the parameters $a, b, c, d$ for the linear fractional map. Of course, these parameters can be chosen as real numbers, but in realistic uses of linear fractional systems one would like to assume that the parameters have finite representations. In particular, one might like to assume that these parameters are rational, and then of course the same linear fractional map can be represented using integer parameters. We would like to know which periods are possible for linear fractional systems with rational parameters. The following theorem gives the answer.

**Theorem 10.6.2.** *A linear fractional system with rational (or integer) parameters can only have periods 1, 2, 3, 4, and 6, and there are linear fractional systems with these periods.*

*Proof.* Let $\gamma = \lambda_1/\lambda_2$, the ratio of the eigenvalues, and $\gamma$ is a root of a polynomial $\gamma^2 - e\gamma + 1$ where $e$ is a rational function of $a, b, c, d$. If the linear fractional system has period $K$, then $\gamma$ is a primitive $K^{th}$ root of unity, i.e., $\gamma^K = 1$ but $\gamma^J \neq 1$ for all $0 < J < K$. For each natural number $K$, it can be shown [122, Section 1.2] that the **cyclotomic polynomial**

$$\Phi_K(x) = \Pi(x - \zeta) \text{ where } \zeta \text{ ranges over all primitive } K^{\text{th}} \text{ roots of unity}$$

has integer coefficients and its degree is $\phi(K)$, where $\phi$ is Euler's Phi Function used in Chapter 8. Further, this polynomial is minimal in the sense that it does not have any smaller degree rational polynomial factors.

Recalling that $\phi(K)$ counts the number of positive integers $i \leq K$ with $\gcd(K, i) = 1$, it can be shown that every integer $K > 6$ has $\phi(K) \geq 3$ because 1 and $K - 1$ and a third number are relatively prime to $K$. We find this third number in each of three cases. One, if $K$ is odd and $K > 5$ then 4 is also relatively prime to $K$. Two, if $K > 6$ and $K = 2m$ with $m$ odd, then $m - 2$ is relatively prime to $K$ because

$$\gcd(2m, m - 2) = \gcd(m - 2, 4) = 1$$

since $m - 2$ is odd. (We need $K > 6$ because for $K = 6$, $m - 2 = 1$.) Three, if $K > 6$ and $K$ is a multiple of 4, then $K/2 - 1$ is relatively prime to $K$ because

$$\gcd(K, K/2 - 1) = \gcd(K/2 - 1, 2) = 1$$

since $K/2 - 1$ is odd.

Returning to the problem at hand, $\gamma$ is a root of $\Phi_K(x)$ which also satisfies a rational *quadratic* polynomial. The minimality of $\Phi_K$ implies $\phi(K)$ must equal 1 or 2. Hence, the only possible values for $K$ are $1, 2, 3, 4, 6$ since we have ruled out all $K \geq 7$ and it's easy to check that $\phi(5) = 4$.

We now give an example of rational linear fractional systems for each of these fiver periods.

Period 1: $f(x) = x$. This is a degenerate linear fractional in which all points have period one.

Period 2: $f(x) = (x-2)/(2x-1)$. Here there are complex eigenvalues and no fixed points. An example period is $2 \longleftrightarrow 0$.
For $f(x) = (5x-2)/(4x-5)$, there are real eigenvalues and two fixed points at $(5 \pm \sqrt{17})/4$. All other points have period 2, for example $1 \longleftrightarrow -3$.

Period 3: $f(x) = (x-1)/x$.
Example period is $3 \longrightarrow 2/3 \longrightarrow -1/2$

Period 4: $f(x) = (x-1)/(x+1)$.
Example period is $2 \longrightarrow 1/3 \longrightarrow -1/2 \longrightarrow -3$

Period 6: $f(x) = (2x-1)/(x+1)$.
Example period is $3 \longrightarrow 5/4 \longrightarrow 2/3 \longrightarrow 1/5 \longrightarrow -1/2 \longrightarrow -4$

□

A pleasant outcome of this analysis is that it is easy to test for periodicity in *rational* linear fractional systems. One simply tries an initial condition and checks to see if it gives periodic behavior of length at most 6. The only minor problem is that one could chance on a fixed point and then one would need to test other points for periodicity.

### *10.6.3 Chaotic-like Behavior*

Three characteristics of **chaos** are:
(a) cycles of every period;
(b) sensitive dependence on initial conditions;
(c) for every open set $A$ and every open set $B$ there is an $x_0 \in A$ and a $K \in N$ so that $f^{(K)}(x_0) \in B$.

For linear fractional systems, (a) is simply false, but (b) and (c) hold when the ratio of the eigenvalues is not a root of unity. Sensitive dependence means that regardless of how closely two trajectories start, eventually these trajectories will be far apart. For linear fractional systems, the pole at $p = -d/c$ will force trajectories to diverge from one another. For example, if one chose to consider trajectories starting at $p-\epsilon$ and $p+\epsilon$, then $|f(p+\epsilon) - f(p-\epsilon)| = 2|Det/c^2\epsilon|$ and this quantity can be made as large as one likes by taking $\epsilon$ small. For some linear fractional systems, this will not be a problem because trajectories will be attracted toward stable fixed points and any initial divergence will disappear in the long term. For *periodic* linear fractional systems, an initial difference will in essence be maintained and not increase. But as we will see, there are complex eigenvalues for which the dependence on initial conditions does not die out. Every trajectory will eventually come close to the pole and then be thrown far away. So two trajectories that start close together will eventually be far apart and there will be no tendency for them to again become close. To see what happens in this case, we will first consider (c) and show that it holds.

**Definition 10.6.1.** A sequence $< S_n >$ is a **source** for $A$ if for every $\alpha \in A$ and for every $\epsilon > 0$ there is a $K$ so that $|\alpha - S_K| < \epsilon$.

**Lemma 10.6.3.** *If $< S_n >$ is a source for $A$ and $g(z)$ is an onto function from $A$ to $B$ so that every pre-range element has a neighborhood of continuity then $g(< S_n >)$ is a source for $B$.*

*Proof.* Let $w$ be the desired point in $B$, then since $g$ is onto there is a $v \in A$ so that $g(v) = w$. Since there is a neighborhood of continuity around $v$, if $|x - v| < \epsilon$ then $|g(x) - g(v)| = |g(x) - w| < \delta$. But since $< S_n >$ is a source for $A$, there is a $K$ so that $|S_K - v| < \epsilon$ and so $|g(S_K) - w| < \delta$ for any desired $w$ and $\delta$, and so $g(< S_n >)$ is a source for $B$. □

**Theorem 10.6.4.** *If $< x_n >= x_0, x_1, ...$ is a trajectory of a linear fractional system in which $\lambda_1/\lambda_2 = e^{i\theta}$ and $\theta$ is an irrational multiple of $\pi$, then $< x_n >$ is a source for $(-\infty, \infty)$.*

*Proof.* The $n^{th}$ term of $< x_n >$ can be written as

$$x_n = \frac{-ax_0 - b + x_0[\cos\theta - \sin\theta \cot n\theta]}{a - cx_0 - [\cos\theta + \sin\theta \cot n\theta]}.$$

Since $\theta$ is an irrational multiple of $\pi$, $< n\theta >$ mod $2\pi$ is a source for $[0, 2\pi]$. [104, Chapter 3] But the transformation from $n\theta$ to $x_n$ satisfies the hypotheses of the lemma, and so $< x_n >$ is a source for $(-\infty, \infty)$. □

Hence, these linear fractional systems obey (c) and they also obey (b) because any two nearby trajectories will eventually hit a small neighborhood of the pole and then be thrown far apart.

Figure 10.6 shows the web diagram for the linear fractional (x-5)/(x+1). One can see that even with the 100 iterates used in the diagram calculation, almost all points are filled in, verifying that this map obeys (c). Figure 10.7 shows sensitive dependence on initial conditions in that the two displayed trajectories are often very close but occasionally they are far apart.

### *10.6.4 Invariant Distributions*

When a system has wandering trajectories that do not converge to cycles or fixed points, one asks if there are perhaps other properties that are conserved. The most obvious question to ask is how long a trajectory stays in a region or how often a trajectory visits a region. It may be difficult to answer these questions and perhaps unenlightening because we want to know how the *system* behaves rather than how a specific trajectory behaves. So, we consider putting a probability distribution on the space and asking how the distribution is changed by the system. In particular, we would like to know if there is a very special distribution that remains the same after the system acts on it. If we could prepare enough copies of a system and start these copies in accord with this invariant distribution, then we would later (even a long time later) still see the same distribution of states. Although the copies that started in particular states would no longer be in those states, other copies would be in those states in proportion to the invariant density. We could also hope for some sort of convergence. It might be possible that the invariant distribution was attractive if not for all initial




FIGURE 10.6. A web plot of the model $\frac{x-5}{x+1}$, showing that most points are visited.

distributions, then at least for a class of initial distributions. By attractive here we mean that under some reasonable definition of distance between distributions, the distance between the system's distribution at time $t$ and the invariant distribution would go to 0, as $t$ increases. We will show linear fractional systems do have invariant distributions but these distributions are not attractive.

Although we spoke of probability *distributions*, it may be easier to work with probability *densities*. A **density function** $g(x)$ defined on $(-\infty, \infty)$ will have the property that for every interval $(\alpha, \beta)$ the probability assigned to that interval is $\int_\alpha^\beta g(x)dx$. For normalization, we want $\int_{-\infty}^{\infty} g(x)dx = 1$. Now let us see how the mass on the interval $(\alpha, \beta)$ is transformed by the mapping $f$. First, the interval $(\alpha, \beta)$ is transformed to $(f(\alpha), f(\beta))$ and then the mass that was on $(\alpha, \beta)$ is spread out on this new interval. But if $g(x)$ is an invariant density, then the density on the new interval must also be $g(x)$. If we let $\hat{g}(x)$ be the distribution on the new interval, we have $\int_\alpha^\beta g(x)dx = \int_{f(\alpha)}^{f(\beta)} \hat{g}(x)dx$ and using $x = f(y)$ and $dx = f'(y)dy$ this becomes $\int_\alpha^\beta g(x)dx = \int_\alpha^\beta \hat{g}(x)f'(y)dy$. Since $\alpha$ and $\beta$ are arbitrary, a limiting argument gives $g(x) = \hat{g}(f(x))f'(x)$ at least for those $x$'s at which these functions are continuous. Said another way, $\hat{g}(x) = g(f^{-1}(x))/f'(f^{-1}(x))$. If $g_I(x)$ is an invariant density then $g_I(x) = g_I(f(x))f'(x)$. Since we want $g_I(x)$ to have a finite integral over $(-\infty, \infty)$, $g_I(x)$ should be close to 0 when $x$ is large in absolute value, and so it may be easier to look at $h(x) = 1/g_I(x)$. Then $h(x) = h(f(x))/f'(x)$. Using a linear fractional map




FIGURE 10.7. Two trajectories for $f(x) = \frac{x-5}{x+1}$. One trajectory starts at $-1.05$ and the other at $-0.95$. These two trajectories are often very close, but occasionally they are far apart.

for $f$, which has been normalized to have $Det = ad - bc = 1$, then

$$h(x) = (cx + d)^2 h\Big(\frac{ax + b}{cx + d}\Big).$$

Assuming that $h(x)$ has a power series, then $h(x)$ must be a polynomial of degree 2. The coefficients of the polynomial can be found by solving a set of linear equations. Up to an unknown scale factor, the unique solution to the set of three linear equations for $h(x)$ is

$$h(x) = cx^2 + (d - a)x - b.$$

**Theorem 10.6.5.** *If $f(x)$ is a linear fractional map with complex eigenvalues, then there is a unique invariant density*

$$g(x) = \frac{\gamma}{cx^2 + (d - a)x - b},$$

*where $\gamma$ is determined from the normalization condition $\int_{-\infty}^{\infty} g(x)dx = 1$.*

The complex eigenvalue condition is equivalent to $g(x)$ having no real poles which in turn implies that $g(x)$ is integrable and that the normalization makes sense.

Conveniently enough, the integral of the density function can also be written in closed form as

$$\int_{-\infty}^{z} g(x)dx = \frac{1}{\pi}\arctan\frac{2cz+d-a}{\sqrt{-(d-a)^2-4bc}} + \frac{1}{2}.$$

If one could choose the states of an ensemble of systems to satisfy the density $g_I(x)$, then as the states evolve by the application of $f$, the same density would be maintained. One might hope that if one chose any density then under $f$ the density might evolve toward $g_I(x)$. The clue that this is not the case is in the theorem's condition that all one needs for an invariant distribution is that the eigenvalues are complex. But it is easy to construct a linear fractional system with complex eigenvalues $i$ and $-i$, and such a system will have all points periodic with period 2. For such a system, *any* initial distribution will repeat itself after two steps and the density will not converge to the invariant density. More specifically, one can show the following theorem.

**Theorem 10.6.6.** *For a linear fractional system with complex eigenvalues the invariant density is not attractive even within the class of densities whose reciprocals are quadratic polynomials. For such densities, the discriminant is a conserved quantity.*

The proof of this theorem consists of showing that the coefficients of the new density can be computed from the coefficients of the old density by applying a matrix to the vector of coefficients. The invariant density corresponds to the 1-eigenvector, the eigenvector associated with the eigenvalue 1. One can then show that the other two eigenvalues of the matrix also have absolute value equal to 1. Hence, even though a density will convert into a new density, the difference from the 1-eigenvector will not decrease and the 1-eigenvector corresponding to the invariant density will not be attractive. In fact, one can also show that the discriminant is preserved. Starting with a density which is the reciprocal of quadratic polynomial and iterating using $f$ will result in a density of the same kind and the discriminant for this density will be identical to the discriminant for the starting density.

What will a typical aperiodic trajectory look like? One way to describe such a trajectory is to use a histogram, that is, to break up the range into small bins and count the number of times the trajectory visits the bin. One can then normalize by the number of iterates, and hope that a limiting histogram exists.

Let $< f^{(n)}(x_0) >$ be the iterates of $f$ starting at $x_0$. If this trajectory is well-behaved, then there is an associated histogram, $H$, so that $H((a,b))$ is the frequency with which the trajectory visits the interval $(a,b)$, i.e.,

$$H((a,b)) = \lim_{N\to\infty}\frac{1}{N}\sum_{i=1}^{N} I_{(a,b)}[f^{(i)}(x_0)]$$




FIGURE 10.8. A histogram for a trajectory of $(x-5)/(x+1)$ showing good agreement with the density $1/(x^2+5)$.

where $I_{(a,b)}[z]$ is an indicator function which gives 1 when its argument is in $(a,b)$ and 0 when its argument is not in $(a,b)$. Assuming that these limits exist and that $H$ is smooth, then

$$H(x) = \lim_{\epsilon \to 0} H((x-\epsilon, x+\epsilon))$$

should exist and behave like a probability density.

Assuming all of this is true, what should $H(x)$ look like? Since we are looking at limiting behavior, it should not matter if the trajectory starts at $x_0$ or at $f(x_0)$. Hence, we can expect the limiting histogram to be a invariant density.

But, by Theorem 10.6.5 there is a unique invariant density, so $H(x)$ should look like $g_I(x)$. Figure 10.8 shows a histogram for 1000 iterates of $(x-5)/(x+1)$. This histogram looks quite smooth and agrees reasonably with the invariant density $1/(x^2+5)$.

These results tell us that in spite of a seemingly irregular trajectory, a fairly simple property is maintained. If one looks at any individual trajectory, then the long-run histogram should look like a fairly simple function. On the other hand, if one took multiple copies of the same linear fractional system and assigned initial conditions for these with the probabilities given by the histogram, and then looked at the distribution of states after one or several time intervals, the probability density will still be the same as the initial density. This is a sort of **ergodic theorem**, which says that the

average over one trajectory is the same as the appropriate average over an ensemble of systems.

Notice that if one picked a density $g(x)$ and picked the initial conditions for an ensemble according to $g(x)$, one would expect a different density say $g_1(x)$ to occur after one time step, and then densities $g_2(x), g_3(x), g_4(x), ...$ for subsequent time steps. There is *no* reason to expect this sequence to converge toward a single density. In fact, Theorem 10.6.6 says this sequence will *not* converge, but if one forms a sequence of averages, these averages will converge. That is, for $G_0(x) = g(x)$ and $G_N(x) = \frac{1}{N}\sum_{i=0}^{N-1} g_i(x)$ then $\lim_{N\to\infty} G_N(x) = g_I(x)$.

TABLE 10.1. The possible asymptotic behavior of a linear fractional system $f(x) = \frac{ax+b}{cx+d}$ (we assume $c \neq 0$ to get a nonlinear system) and set $Det = ad - bc$, $Disc = (a-d)^2 + 4bc$.

| | | | |
|---|---|---|---|
| $Det = 0$ | | $f(x) = constant$ | Superstable fixed point |
| $Det < 0$ | | One stable fixed point | |
| | $d + a \neq 0$ | One unstable fixed point | |
| Two | | Convergence $x_n = F.P. + O(\lambda^n)$, $\lvert\lambda\rvert < 1$ | |
| real | | All points except the fixed points | |
| fixed | $d + a = 0$ | have period 2 | |
| points | | neutral stability | |
| $Det > 0$ | $Disc = 0$ | One globally stable fixed point | |
| | | Convergence $x_n = F.P. + O(\frac{1}{n})$ | |
| | $Disc > 0$ | One stable fixed point | |
| | | One unstable fixed point | |
| | | Convergence $x_n = F.P. + O(\lambda^n)$, $\lvert\lambda\rvert < 1$ | |
| | $Disc < 0$ | Periodic, all points | Rational coeffs - possible periods 1,2,3,4,6 |
| | | have the same period | Irrational coeffs - any period possible |
| | | Chaotic-like | No periodic points |
| | | | All open sets visited |
| | | | Sensitive dependence on initial conditions |
| | | | Invariant density - Non-attractive |

### *10.6.5 Showing Global Stability*

Our initial interest in linear fractional systems came from an application to population models. For some years, it has been known that the usual one-dimensional population models were globally stable exactly when they were locally stable [54, 26, 129]. We were pleased to find that the usual population models were "enveloped" by linear fractional maps of the special form:

$$\phi(x) = \frac{1 - \alpha x}{\alpha - (2\alpha - 1)x} \qquad \text{where } \alpha \in [0, 1) \ .$$

These special linear fractional systems all have period 2, and so by a variation on Sarkovskii's Theorem (refer to Theorem 10.2.1) we were able to show:

**Theorem 10.6.7.** *Let $\phi(x)$ be a monotone decreasing function which is positive on $(0, x_-)$ and so that $\phi(\phi(x)) = x$. Assume that $f(x)$ is a continuous function such that:*

- $\phi(x) > f(x)$ *on* $(0, 1)$
- $\phi(x) < f(x)$ *on* $(1, x_-)$
- $f(x) > x$ *on* $(0, 1)$
- $f(x) < x$ *on* $(1, \infty)$
- $f(x) > 0$ *on* $(1, x_\infty)$

*then for all* $x \in (0, x_\infty)$, $\lim_{k\to\infty} f^{(k)}(x) = 1$.

This theorem enabled us to show local stability implied global stability for the following seven population models:

- $x_{t+1} = x_t e^{r(1-x_t)}$
- $x_{t+1} = x_t[1 + r(1 - x_t)]$
- $x_{t+1} = x_t[1 - r \ln x_t]$
- $x_{t+1} = x_t(\frac{1}{b+cx_t} - d)$
- $x_{t+1} = \frac{(1+ae^b)x}{1+ae^{bx}}$
- $x_{t+1} = \frac{(1+a)^b x}{(1+ax)^b}$ with $a > 0$, $b > 0$
- $x_{t+1} = \frac{rx}{1+(r-1)x^c}$

Details of these results appear in [32].

### *10.6.6 Summary*

Linear fractionals form a simple class of nonlinear systems, yet they display many of the behaviors possible for nonlinear systems. In particular, they exhibit both stable and unstable fixed points, periodic behavior, and even chaotic-like behavior. Table 10.1 gives a summary of the possible behaviors and the corresponding conditions on the parameters.

In contrast to most nonlinear systems, linear fractionals can be analyzed using techniques from elementary mathematics and linear systems. Further, we have shown that linear fractional systems can be used to analyze more complicated systems and suggest that linear fractionals should be a standard part of the toolbox for studying nonlinear systems.

## 10.7 Conclusion

As we said at the beginning of this chapter, nonlinear systems can be nonlinear in many different ways. We have given only some very simple cases, and our examples have usually been one-dimensional. Realistic systems are often high dimensional and the simple analyses possible in one dimension will not apply. On the other hand showing that *chaos* is possible in one dimension both warns us that high dimensional systems may behave in very complicated ways, and suggests that we may be able to explain this complicated behavior by finding a chaotic one-dimensional subsystem. We have also limited our examples to the reals or rationals, but some systems may be easier to understand if one extends the analysis to the complex numbers. For example the linear fractionals may or may not have fixed points in the reals, but such systems always have fixed points in the complex numbers. Other nonlinear systems may be comprised of simple components with finite state sets like neural nets, or have an infinite but locally finite state set like cellular automata, or have state sets like the natural numbers or strings over a finite alphabet like Turing machines and other models of computation. Analyses of such systems can be shown to be impossible. Even with the limitation to finite state sets, the analysis of such systems can be shown to be possible but practically unreasonable.

We close with the warning that realistic systems may be highly nonlinear and highly complicated, but with the hope that, as in the past, future generations of scientists and engineers will be able to find simple enough models to solve societal problems.

## 10.8 Exercises

**Ex 10.1.** Show that if $\lim_{k\to\infty} f^{(k)}(x_0) = p$ and $f$ is a continuous function then $p$ is a fixed point of $f$.

**Ex 10.2.** Let $x_{t+1} = f(x_t)$ where

$$f(x) = \begin{cases} \frac{x}{2} + \frac{1}{4} & \text{if} \quad x < \frac{1}{2} \\ \frac{x}{2} + \frac{1}{2} & \text{if} \quad \frac{1}{2} < x \leq 1 \\ 1 & \text{if} \quad x > 1. \end{cases}$$

Show that for $x_0 \in (-\infty, 1/2)$ $\lim_{K\to\infty} f^{(K)}(x_0) = 1/2$, but for $x_0 > 1/2$, $\lim_{K\to\infty} f^{(K)}(x_0) = 1$. Does this contradict the results of the previous exercise? Can this system enveloped by a linear fractional system? Does this violate theorem 10.6.7?

**Ex 10.3.** Show that $x_{t+1} = \sin x_t$ has one fixed point and that this point is both locally and globally stable. [HINT: A local linear approximation is insufficient. Use a local nonlinear approximation to show local stability.]

**Ex 10.4.** Analyze the local and global stability of $x = 1$ for the system:

$$x_{t+1} = x_t[1 + 2(1 - x_t)].$$

**Ex 10.5.** Analyze the local and global stability of $x = 1$ for the system:

$$x_{t+1} = x_t[1 + 2.00001(1 - x_t)].$$

**Ex 10.6.** Show that

$$x_{t+1} = x_t[1 + r \ln x_t]$$

has a locally stable period 2 cycle when $r$ is slightly bigger than 2.

**Ex 10.7.** Let $g(x) = \sqrt{|x|}$. Show that Newton's method applied to find the root of $g(x)$ fails to converge. Show, in fact, that if $x_0 \neq 0$, Newton's method will oscillate with period 2.

**Ex 10.8.** Let $x_{t+1} = f(x_t)$ where

$$f(x) = \begin{cases} -\sqrt{x} & \text{if} x \geq 0 \\ \sqrt{|x|} & \text{if} x < 0 \end{cases} .$$

Show that this system has a repelling fixed point and an attracting period 2 cycle. Is this period 2 cycle globally stable?

**Ex 10.9.** Let $x_{t+1} = f(x_t)$ where

$$f(x) = \frac{x^2 + 2}{2x - 1}.$$

Show that $x_0 = 1/2$ separates the behavior of this system in the sense that if $x_0 < 1/2$ there is one type of behavior and if $x_0 > 1/2$ there is a different behavior.

**Ex 10.10.** Consider the system

$$\begin{aligned} x_{t+1} &= x_t e^{r(1-x_t-y_t)} \\ y_{t+1} &= y_t e^{r(1-x_t-y_t)}. \end{aligned}$$

Find a quantity which is *invariant* along trajectories. Discuss the convergence properties of this system in light of this invariant. Are there fixed points? Are any of these fixed points locally or globally stable?

**Ex 10.11.** For the difference equation

$$x_{t+1} = 2x_t \pmod 1,$$

plot $x_{t+1}$ as a function of $x_t$. Find the fixed points in your diagram. Does the plot show a discontinuity? Plot $x_{t+2}$ as a function of $x_t$. Can you find the oscillations of period 2?

**Ex 10.12.** For $x_{t+1} = 2x_t \pmod 1$, plot $x_{t+3}$ as a function of $x_t$. How does this show that there are 2 distinct period 3 oscillations? Find these oscillations.

**Ex 10.13.** A sequence is *eventually periodic* if there is an $r$ so that $x_r, x_{r+1}, \ldots$ is periodic of period $p$. Show that $x_{t+1} = 2x_t \pmod 1$ has lots of eventually periodic solutions by giving a procedure which takes $r$ and $p$ as input and outputs an initial condition which gives a solution of eventual period $p$ after a run-in of $r$ steps.

**Ex 10.14.** For $x_{t+1} = 2x_t \pmod 1$ we can find an oscillation of period 5 following our construction for period $K$. Show that this is *not* the only possible construction by finding another solution of period 5 for this equation.

**Ex 10.15.** Show that $x_{t+1} = 2x_t \pmod 1$ has at most $\lfloor (2^K - 1)/K \rfloor$ oscillations of period $K$. Can you find a better formula for the number of distinct oscillations of period $K$?

**Ex 10.16.** Is

$$x_{t+1} = 2x_t \pmod 1$$

a linear equation? Refer to the definition and notice that something is missing. Show that by filling in the missing part of the definition in different ways, you can declare the equation to be linear or you can show that it is nonlinear.

**Ex 10.17.** Consider the one-dimensional system

$$x_{t+1} = f(x_t) = x[1 + r(a - x)]$$

for some fixed positive $a, r$ and $ar > 2$. Show that the system has two fixed points and at least one locally stable 2-cycle when $ar < \sqrt{6}$.

**Ex 10.18.** Consider the two-dimensional system

$$x_{t+1} = f_1(x_t, y_t) = x(5 - x - y)/3$$
$$y_{t+1} = f_2(x_t, y_t) = y(3y - x)/3.$$

Show that $(2, 0)$ is a locally stable fixed point of the system and $(3/4, 5/4)$ is a fixed point which is not locally stable.

**Ex 10.19.** Show that the two-dimensional system

$$x_{t+1} = f_1(x_t, y_t) = x(1 + x + y)/3$$
$$y_{t+1} = f_2(x_t, y_t) = y(1 - x + y)/2.$$

has four fixed points, but only one is locally stable.

**Ex 10.20.** Show that

$$x_{t+1} = \frac{3x_t + 2}{2x_t + 1}$$

has two fixed points, $\lambda_0 > 0$ and $\lambda_1 < 0$, and that $\lambda_0$ is locally stable but $\lambda_1$ is repelling. Further show that the initial value $x_0 = 0$ gives $x_n = f_{3n}/f_{3n-1}$, where $f_i$ is the $i^{\text{th}}$ Fibonacci number. Give a good estimate for how close $x_n$ is to $\lambda_0$. How does the recurrence behave for other initial values?

# References

[1] F. Acton. *Numerical Methods That (Usually) Work.* Harper and Row, New York City, NY, 1970.

[2] A. V. Aho, J. E. Hopcroft, and J. D. Ullman. *The Design and Analysis of Computer Algorithms.* Addison-Wesley, Reading, MA, 1974.

[3] W. R. Alford, A. Granville, and C. Pomerance. There are infinitely many Carmichael numbers. *Annals of Mathematics (2)*, 139:705–722, 1994.

[4] R. Askey and M. Ismail. *Recurrence Relations, Continued Fractions, and Orthogonal Polynomials.* American Mathematical Society, Providence, RI, 1984.

[5] W. W. Rouse Ball and H. M. S. Coxeter. *Mathematical Recreations and Essays.* Dover Publications, New York City, NY, 1987.

[6] L. Beckingham, R. M. Compton, and C. J. Pennycuick. A computer model for simulating the growth of a population, or of two interacting populations. *Journal of Theoretical Biology*, 18:316–329, 1968.

[7] È. G. Belaga. Some problems involved in the calculation of polynomials. *Dokl. Akad. Nauk. SSSR*, 123:775–777, 1958.

[8] A. Berman and R. J. Plemmons. *Nonnegative Matrices in the Mathematical Sciences.* SIAM, Philadelphia, PA, 1994.

[9] H. Bernardelli. Population waves. *J. Burma Res. Soc.*, 31:3–18, 1941.

[10] W. Beyer, R. Roof, and D. Williamson. The lattice structure of multiplicative congruential pseudo-random vectors. *Mathematics of Computation*, 25:345–363, 1971.

[11] B. Bollobás. *Random Graphs*. Academic Press, London, 1985.

[12] B. Boncompagni. *Scritti di Leonardo Pisano mathematico del secolo decimoterzo*. Rome, Italy, 1857-1862.

[13] C. B. Boyer. *A History of Mathematics*. Wiley, New York, 1989.

[14] Brother A. Brousseau. The relation of the zeros to periods in the Fibonacci sequence modulo a prime. *American Mathematical Monthly*, 71:897–899, 1964.

[15] Brother A. Brousseau. *Fibonacci and Related Number Theoretic Tables*. Fibonacci Association, San Jose, CA, 1972.

[16] W. G. Brown. Historical note on a recurrent combinatorial problem. *American Mathematical Monthly*, 72:973–977, 1965.

[17] P. Buneman and L. Levy. The Towers of Hanoi problem. *Information Processing Letters*, 10:243–244, 1980.

[18] R. L. Burden and J. D. Faires. *Numerical Analysis*. Brooks/Cole, Pacific Grove, CA, 2001.

[19] R. M. Capocelli, G. Cerbone, P. Cull, and J. Holloway. Fibonacci facts and formulas. In *Sequences*, pages 123–137. Springer-Verlag, 1990.

[20] R. M. Capocelli and P. Cull. Rounding the solutions of Fibonacci-like difference equations. *Fibonacci Quarterly*, to appear.

[21] E. Catalan. Note sur une équation aux différence finies. *J. Math. Pures Appl.*, 3:508–516, 1838.

[22] A. Cohn. Uber die Anzahl der Wurzeln einer algebraischen Gleichung in einem Kreise. *Mathematische Zeitscrift*, 14:110–148, 1922.

[23] D. Coppersmith and S. Winograd. Matrix multiplication via arithmetic progression. *Journal of Symbolic Computation*, 9:251–280, 1990.

[24] R. Crandall and C. Pomerance. *Prime Numbers: A computational approach*. Springer, New York City, NY, 2001.

[25] P. Cull. The problem of time unit in Leslie's population model. *Bulletin of Mathematical Biology*, 42:719–728, 1980.

[26] P. Cull. Global stability of population models. *Bulletin of Mathematical Biology*, 43:47–58, 1981.

[27] P. Cull. Local and global stability for population models. *Biological Cybernetics*, 54:141–149, 1986.

[28] P. Cull. Local and global stability of discrete one-dimensional population models. In *Biomathematics and Related Computational Problems*, pages 271–278. Kluwer, Dordrecht, 1988.

[29] P. Cull. Stability of discrete one-dimensional population models. *Bulletin of Mathematical Biology*, 50(1):67–75, 1988.

[30] P. Cull. Analysis of algorithms. In *Lectures in Applied Mathematics and Informatics*, pages 1–61. Manchester University Press, 1990.

[31] P. Cull. Linear fractionals - simple models with chaotic-like behavior. In Dubois [47], pages 170–181.

[32] P. Cull and J. Chaffee. Stability in discrete population models. In *Computing Anticipatory Systems: CASYS'99*, pages 263–275. American Institute of Physics, Woodbury, NY, 2000.

[33] P. Cull and J. Chaffee. Stability in simple population models. In *Cybernetics and Systems 2000*, pages 289–294. Austrian Society for Cybernetics Studies, 2000.

[34] P. Cull and J. Holloway. Computing fibonacci numbers quickly. *Information Processing Letters*, 32:143–149, 1989.

[35] P. Cull and E. F. Ecklund Jr. Towers of Hanoi and analysis of algorithms. *American Mathematical Monthly*, 92(6):407–420, June-July 1985.

[36] P. Cull and A. Vogt. Mathematical analysis of the asymptotic behavior of the Leslie population matrix model. *Bulletin of Mathematical Biology*, 35:645–661, 1973.

[37] P. Cull and A. Vogt. The periodic limit for the Leslie model. *Mathematical Biosciences*, 21:39–54, 1974.

[38] P. Cull and A. Vogt. The period of total population. *Bulletin of Mathematical Biology*, 38:317–319, 1976.

[39] J. H. Davenport, Y. Siret, and E. Tournier. *Computer Algebra*. Chapman and Hall, San Diego, CA, 1988.

[40] A. DeMoivre. *Miscellanea Analytica*. Londini, excudebant J. Tonson & J. Watts, London, 1730.

[41] R. Descartes. *Discourse on Method.* Penguin, New York City, NY, 1968.

[42] R. Descartes. *Meditations.* Penguin, New York City, NY, 1968.

[43] R. Devaney. *An Introduction to Chaotic Dynamical Systems.* Benjamin, Redwood City, 1986.

[44] P. Diaconis, M. McGrath, and J. Pitman. Riffle shuffles, cycles, and discents. *Combinatorica*, 15:11–29, 1995.

[45] L. E. Dickson. *Linear Algebraic Groups and an Exposition of Galois Theory.* Dover Publications, New York City, NY, 1958.

[46] D. M. Dubois, editor. *Computing Anticipatory Systems:CASYS 99 - Third International Conference, Conference Proceedings.* American Institue of Physics, Woodbury, N.Y., 2000.

[47] D. M. Dubois, editor. *Computing Anticipatory Systems:CASYS 2002 - Fifth International Conference, Conference Proceedings 627.* American Institue of Physics, Woodbury, N.Y., 2002.

[48] F. Dyson and H. Falk. Period of a discrete cat mapping. *American Mathematical Monthly*, 99:603–614, 1992.

[49] J. Eichenauer-Hermann. Inversive congruential pseudorandom numbers avoid the planes. *Mathematics of Computation*, 56:297–301, 1991.

[50] P. Erdos. On almost primes. *American Mathematical Monthly*, 57:404–407, 1950.

[51] L. Euler. *Novi Commentarii Academiae Scientiarum Imperalis Petropolitanae*, 7:13–14, 1758-1759.

[52] L. Euler. *Introduction to Analysis of the Infinite.* Springer-Verlag, New York City, NY, 1988. Translated by J. D. Blanton.

[53] W. Feller. *An Introduction to Probability Theory and its Applications.* John Wiley, New York City, NY, 1968.

[54] M. E. Fisher, B. S. Goh, and T. L. Vincent. Some stability conditions for discrete-time single species models. *Bulletin of Mathematical Biology*, 41:861–875, 1979.

[55] M. Flahive and H. Niederreiter. On inversive congruential generators for pseudorandon numbers. *Lecture Notes in Pure and Applied Mathematics*, 141:75–80, 1993.

[56] H. Furstenberg. *Recurrence in Ergodic Theory and Combinatorial Number Theory.* Princeton University Press, Princeton, NJ, 1981.

[57] É. Galois. *Oeuvres Mathematiques.* Gauthier-Villars, Paris, France, 1897.

[58] F. R. Gantmacher. *The Theory of Matrices.* Chelsea Publishing Company, New York City, NY, 1959.

[59] M. Garey and D. Johnson. *Computers and Intractability: A Guide to the Theory of NP-Completeness.* W. H. Freeman, San Francisco, 1979.

[60] F. Garvin. *The Maple Book.* Chapman and Hall, Boca Raton, FL, 2002.

[61] C. F. Gauss. *Disquisitiones Arithmeticae.* Springer-Verlag, New York City, NY, 1986.

[62] B. S. Goh. *Management and Analysis of Biological Populations.* Elsevier, New York, 1979.

[63] R. L. Graham, D. E. Knuth, and O. Patashnik. *Concrete Mathematics: A Foundation for Computer Science.* Addison-Wesley, Reading, MA, 1994.

[64] R. T. Gregory. *Methods and Applications of Error-Free Computation.* Springer-Verlag, New York City, NY, 1984.

[65] M. A. Harrison. *Lectures on Linear Sequential Machines.* Academic Press, New York City, NY, 1969.

[66] M. P. Hassel. Density dependence in single species populations. *Journal of Animal Ecology*, 44:283–296, 1974.

[67] K. Hoffman and R. Kunze. *Linear Algebra (second edition).* Prentice-Hall, Englewood Cliffs, NJ, 1971.

[68] J. L. Holloway. *Algorithms for Computing Fibonacci Numbers Quickly.* Oregon State University MS thesis, Corvallis, OR, 1989.

[69] C. Hooley. On Artin's conjecture. *J. reine angew. Math*, 226:209–220, 1967.

[70] Y. N. Huang. A counterexample for P. Cull's theorem. *Kexue Tongbao*, 31:1002–1003, 1986.

[71] D. Kalman. The generalized Vandermonde matrix. *Mathematics Magazine*, 57:15–21, 1984.

[72] J. Keller. How many shuffles to mix a deck? *SIAM Review*, 37:88–89, 1995.

[73] A. Knight. *Basics of MATLAB and Beyond*. Chapman and Hall, Boca Raton, FL, 2000.

[74] D. Knuth. Big Omicron and Big Omega and Big Theta. *SIGACT News*, 8:18–24, April-June 1976.

[75] D. Knuth. *Selected Articles on Analysis of Algorithms*, pages 35–42. Addison-Wesley, Reading, MA, 2000.

[76] D. Knuth. All questions answered. *Notices of the American Mathematical Society*, 49:318–324, 2002.

[77] D. E. Knuth. *The Art of Computer Programming*. Addison-Wesley, New York City, NY, third edition, 1997.

[78] R. J. Kooman. *Convergence Properties of Recurrence Sequences*. Centrum voor Wiskunde en Informatica, Amsterdam, The Netherlands, 1991.

[79] T. Koshy. *Fibonacci and Lucas Numbers*. Wiley-Interscience, New York City, NY, 2001.

[80] G. Lamé. Extrait d'une letter de M. Lamé à M. Liouville sur cette question: un polygone convexe étant donné, de combien de manières peut-on le partager en triangles au moyen de diagonales? *J. Math. Pures Appl.*, 3:505–507, 1838.

[81] J. P. LaSalle. *The Stability of Dynamical Systems*. SIAM, Philadelphia, 1976.

[82] H. Lebesgue. L'oeuvre mathématique de Vandermonde. *L'Enseignement mathématique (2)*, 1:203–223, 1956.

[83] D. H. Lehmer. Mathematical methods in large-scale computing units. In *Proceedings of the Second Symposium on Large-Scale Digital Computing Machinery*, pages 141–146. Harvard University Press, Cambridge, MA, 1951.

[84] P. H. Leslie. On the use of matrices in certain population mathematics. *Biometrika*, 33:183–212, 1945.

[85] T-Y. Li and J. Yorke. Period three implies chaos. *American Mathematical Monthly*, 82:985–992, 1975.

[86] R. Lidl and H. Niederreiter. *Finite Fields*. Cambridge University Press, Cambridge, England, 1997.

[87] D. Lind and B. Marcus. *An Introduction to Symbolic Dynamics and Coding*. Cambridge University Press, New York City, NY, 1995.

[88] A. J. Lotka. *Elements of Mathematical Biology*. Dover Publications, New York, 1956.

[89] U. Manber. *Introduction to Algorithms: A Creative Approach*. Addison-Wesley, Reading, MA, 1989.

[90] M. Marcus and H. Minc. *A Survey of Matrix Matrix Theory and Matrix Inequalities*. Allyn and Bacon, Rockleigh, NJ, 1964.

[91] M. Marden. Much ado about nothing. *American Mathematical Monthly*, 83:788–798, 1976.

[92] M. Marden. *The Geometry of the Zeros of a Polynomial in a Complex Variable*. American Mathematical Society, New York City, NY, 1989.

[93] G. Marsaglia. Random numbers fall mainly in the planes. *Proceedings of the National Academy of Sciences, U.S.A.*, 61:25–28, 1968.

[94] G. Marsaglia. The structure of linear congruential sequences. In *Applications of Number Theory to Numerical Analysis*, pages 249–285. Academic Press, New York City, NY, 1972.

[95] Y. V. Matiyasevich. *Hilbert's Tenth Problem*. The MIT Press, Cambridge, MA, 1993.

[96] R. M. May. Biological populations with nonoverlapping generations: stable points, stable cycles, and chaos. *Science*, 186:645–647, 1974.

[97] R. M. May. Simple mathematical models with very complicated dynamics. *Nature*, 261:459–467, 1976.

[98] K. Mehlhorn. *Data Structures and Algorithms*. Springer-Verlag, New York City, NY, 1984.

[99] P. L. Montgomery. Modular multiplication without trial division. *Mathematics of Computation*, 44:519–522, 1985.

[100] P. A. P. Moran. Some remarks on animal population dynamics. *Biometrics*, 6:250–258, 1950.

[101] T. Muir. *The Theory of Determinants in the Historical Order of Development*, volume 3. Dover Publications, New York City, NY, 1960.

[102] H. Neiderreiter. Quasi-Monte Carlo methods and pseudorandom numbers. *Bulletin of the American Mathematical Society*, 84:957–1041, 1978.

[103] H. Niederreiter. *Random Number Generation and Quasi-Monte Carlo Methods.* SIAM, Philadelphia, PA, 1992.

[104] I. Niven. *Diophantine Approximations.* Interscience Publishers, New York City, NY, 1963.

[105] I. Niven. Formal power series. *American Mathematical Monthly*, 76:871–889, 1969.

[106] A. Nobile, L. M. Ricciardi, and L. Sacerdote. On Gompertz growth model and related difference equations. *Biological Cybernetics*, 42:221–229, 1982.

[107] V. Ya. Pan. On methods of computing polynomial values. *Russian Mathematical Surveys*, 21:105–137, 1966.

[108] F. Parker. Inverses of Vandermonde matrices. *American Mathematical Monthly*, 71:410–411, 1964.

[109] T. S. Parker and L. Chua. *Practical Numerical Algorithms for Chaotic Systems.* Springer-Verlag, New York City, NY, 1989.

[110] P. Petersen. *On computing maximal lattice dimensions of the inversive congruential generator.* Thesis for the Masters of Science degree, Corvallis, OR, 1998.

[111] J. M. Pollard. A Monte Carlo method for factorization. *BIT*, 15:331–334, 1975.

[112] G. Pólya. *How to Solve It.* Princeton University Press, Princeton, NJ, 1945.

[113] G. Pólya. On picture writing. *American Mathematical Monthly*, 63:689–697, 1956.

[114] L. M. Ricciardi, editor. *Biomathematics and Related Computational Problems.* Kluwer, Dordrecht, 1988.

[115] W. E. Ricker. Stock and recruitment. *Journal of the Fisheries Research Board of Canada*, 11:559–623, 1954.

[116] R. H. Risch. The problem of integration in finite terms. *Transactions of the American Mathematical Society*, 139:167–189, 1969.

[117] R. L. Rivest, A. Shamir, and L. M. Adelman. A method for obtaining digital signatures and public-key cryptosystems. *Communications of the ACM*, 21:145–152, 1978.

[118] H. Rogers. *Theory of Recursive Functions and Effective Computability.* McGraw-Hill, New York City, NY, 1967.

[119] K. H. Rosen. *Elementary Number Theory and its Applications.* Addison-Wesley, Reading, MA, 1993.

[120] R. Rosen. *Anticipatory Systems.* Pergamon Press, New York, 1985.

[121] G. Rosenkranz. On global stability of discrete population models. *Mathematical Biosciences*, 64:227–231, 1983.

[122] J. J. Rotman. *Advanced Modern Algebra.* Prentice Hall, Upper Saddle River, NJ, 2002.

[123] D. G. Saari and J. B. Urenko. Newton's method, circle maps, and chaotic motion. *The American Mathematical Monthly*, 91:3–18, 1984.

[124] A. Salomaa. *Automata-theoretic Aspects of Formal Power Series.* Springer-Verlag, New York, NY, 1978.

[125] A. Sarkovskii. Coexistence of cycles of a continuous map of a line to itself. *Ukr. Mat. Z.*, 16:61–71, 1964.

[126] A. Schonhage and V. Strassen. Schnelle Multiplikation grosser Zahlen. *Computing*, 7:281–292, 1971.

[127] E. Seneta. *Non-negative Matrices.* John Wiley & Sons, New York City, NY, 1973.

[128] L. Sigler. *Fibonacci's Liber Abbaci.* Springer Verlag, New York City, NY, 2002.

[129] D. Singer. Stable orbits and bifurcation of maps of the interval. *SIAM Journal on Applied Mathematics*, 35(2):260–267, Sept 1978.

[130] D. Smith and M. Latham. *The Geometry of Rene Descartes.* Open Court Publishing Company, Chicago, IL, 1925.

[131] J. M. Smith. *Mathematical Ideas in Biology.* Cambridge University Press, Cambridge, 1968.

[132] J. M. Smith. *Models in Ecology.* Cambridge University Press, Cambridge, 1974.

[133] R. P. Stanley. Generating functions. In *Studies in Combinatorics*, pages 100–141. Mathematical Association of America, Washington, D.C., 1978.

[134] R. P. Stanley. *Enumerative Combinatorics*, volume 1. Wadsworth & Brooks/Cole, Monterey, CA, 1986.

[135] R. P. Stanley. *Enumerative Combinatorics*, volume 2. Cambridge University Press, New York City, NY, 1999.

[136] J. Stoer and R. Bulirsch. *Introduction to Numerical Analysis.* Springer-Verlag, New York City, NY, 1993.

[137] V. Strassen. Gaussian elimination is not optimal. *Numer. Math*, 13:354–356, 1969.

[138] A. Tarski. *Logic, Semantics, Metamathematics.* Oxford University Press, Oxford, England, 1956.

[139] R. Taylor and A. Wiles. Modular elliptic curves and Fermat's Last Theorem. *Annals of Mathematics (2)*, 141:443–551, 1995.

[140] A. Tucker. *Applied Combinatorics.* John Wiley, New York City, NY, 2002.

[141] J. B. Urenko. Improbability of nonconvergent chaos in Newton's method. *Journal of Mathematical Analysis and Applications*, 117:42–47, 1986.

[142] S. Utida. Population fluctuation, an experimental and theoretical approach. *Cold Spring Harbor Symposium on Quantitative Biology*, 22:139–151, 1957.

[143] A. van der Poorten. *Notes on Fermat's Last Theorem.* John Wiley & Sons, New York City, NY, 1996.

[144] J. Vinson. The relation of the period modulo to the rank of apparition of $m$ in the Fibonacci sequence. *Fibonacci Quarterly*, 1:37–45, 1963.

[145] N. N. Vorobev. *Fibonacci Numbers.* Blaisdell, New York City, NY, 1961.

[146] D. D. Wall. Fibonacci series modulo $m$. *American Mathematical Monthly*, 67:525–532, 1960.

[147] A. Wiles. Ring-theoretic properties of certain Hecke algebras. *Annals of Mathematics (2)*, 141:553–572, 1995.

[148] H. S. Wilf. *generatingfunctionology.* Academic Press, New York City, NY, 1990.

[149] J. Wimp. *Computation with Recurrence Relations.* Pitman, Boston, MA, 1984.

[150] N. Wirth. *Algorithms + Data Structures = Programs.* Prentice-Hall, Englewood Cliffs, NJ, 1976.

[151] A. Wiseman and T. P. Wiseman. *De bello Gallico. English.* D.R. Godine, Boston, MA, 1980.

[152] S. Wolfram. *Mathematica*. Addison-Wesley, Redwood City, CA, 1988.

[153] S. Wolfram. *A New Kind of Science*. Wolfram Media, Champaign, IL, 2002.

[154] O. Wyler. On second-order recurrences. *American Mathematical Monthly*, 72:500–506, 1965.

[155] N. Zierler. Linear recurring sequences and error-correcting codes. In *Error-Correcting Codes*, pages 47–59. Wiley, New York City, NY, 1968.

# Appendix A
## Worked Examples

In the first chapters of this book we considered **$\mathbf{k^{th}}$-order linear recurrences**, that is, equations of the form

$$\text{(L)} \qquad s_n - c_1 s_{n-1} - c_2 s_{n-2} \cdots - c_k s_{n-k} = \psi(n) \text{ for } n \geq k\,,$$

where $c_i \in \mathbb{C}$ with $c_k \neq 0$ and $\psi$ is a complex-valued function. There we found a nice solution for the special case in which the forcing function has the form

$$\psi(n) = \lambda^n p(n)$$

for fixed $\lambda \in \mathbb{C}$ and polynomial $p$ with complex coefficients. The initial value problems we consider in this appendix have this form. Our principal tool is Theorem 3.2.1, which we use to find a particular solution $< v_n >$. Throughout, $\lambda_1, \ldots, \lambda_t$ are the *different* eigenvalues of the recurrence.

## A.1 Every eigenvalue is simple

All examples in this section are second-order linear equations with characteristic polynomial

$$ch(x) = x^2 - x - 6 = (x-3)(x+2)$$

which has the simple roots $\lambda_1 = 3, \lambda_2 = -2$, and the general solution of the *homogeneous* system has the form $s_n = a_1 3^n + a_2(-2)^n$ for some $a_1, a_2 \in \mathbb{C}$. Therefore, for any $\psi$ the equation

$$s_n = s_{n-1} + 6s_{n-2} + \psi(n)$$

has general solution

$$s_n = a_1 3^n + a_2(-2)^n + v_n\,,$$

for some particular solution $v_n$ and the constants $a_1, a_2$ are determined by the initial conditions. Let's first analyze the homogeneous initial value problem.

*Example* A.1.1. For any initial value problem with equation

$$s_n = s_{n-1} + 6s_{n-2}$$

we have $s_n = a_1 3^n + a_2(-2)^n$ and the initial conditions $s_0, s_1$ give

$$s_0 = a_1 + a_2 \quad \text{and} \quad s_1 = 3a_1 - 2a_2. \tag{A.1}$$

Multiplying the first equation by 3 and subtracting that from the second equation, we obtain

$$s_1 - 3s_0 = -5a_2 \;;\; a_2 = \frac{3s_0 - s_1}{5}.$$

Inserting this value for $a_2$ into the first equation of (A.1)

$$a_1 = s_0 - \frac{3s_0 - s_1}{5} = \frac{2s_0 + s_1}{5}\,,$$

and the solution to the initial value problem is

$$s_n = \frac{2s_0 + s_1}{5}3^n + \frac{3s_0 - s_1}{5}(-2)^n.$$

Theorem 3.2.1 can be used to find a particular solution to the *nonhomogeneous* equation

$$s_n = s_{n-1} + 6s_{n-2} + \lambda^n p(n)$$

where $p$ is a non-zero polynomial and $\lambda$ is a scalar. For these equations the following special case of the theorem applies

**Solving (L) when there are no multiple eigenvalues**

*When there are no multiple eigenvalues then* $v_n = \lambda^n n^\delta q(n)$ *is a particular solution to (L) where* $q(x)$ *is a fixed polynomial with* $\deg(q) = \deg(p)$ *and*

$$\delta = \begin{cases} 0 & \text{if } \lambda \notin \{\lambda_1, \ldots, \lambda_t\} \\ 1 & \text{if } \lambda \in \{\lambda_1, \ldots, \lambda_t\} \end{cases}. \tag{A.2}$$

*Example* A.1.2. Consider the second-order equation

$$s_n = s_{n-1} + 6s_{n-2} + 2^n\,,$$

which has $\deg(p) = 0$ and $\lambda = 2 \notin \{3, -2\}$ and so $\deg(q) = 0$ and $\delta = 0$. The result gives the particular solution $v_n = c\,2^n$, and the recurrence can be used to solve for $c$:

$$v_{n-1} + 6v_{n-2} + 2^n - v_n = c2^{n-1} + 6c2^{n-2} + 2^n - c2^n = 2^{n-1}(c + 3c + 2 - 2c)\,.$$

Therefore, $2c + 2 = 0$ which gives the particular solution $v_n = -2^n$, and

$$s_n = a_1 3^n + a_2(-2)^n - 2^n \text{ for some } a_1, a_2 \in \mathbb{C}\,. \tag{A.3}$$

We can write this in terms of the initial values $s_0, s_1$, namely,

$$s_0 = a_1 + a_2 - 1 \text{ and } s_1 = 3a_1 - 2a_2 - 2, \tag{A.4}$$

which can be solved for $a_1, a_2$ as in the previous example. Multiplying the first equation by 2 and adding it to the second, we have

$$2s_0 + s_1 = 5a_1 - 4\ ;\ a_1 = \frac{2s_0 + s_1 + 4}{5}.$$

Substituting this value of $a_1$ into the first equation of (A.4) we obtain

$$a_2 = s_0 - a_1 + 1 = s_0 - \frac{2s_0 + s_1 + 4}{5} + 1 = \frac{3s_0 - s_1 + 1}{5},$$

and (A.3) becomes

$$s_n = \frac{2s_0 + s_1 + 4}{5}3^n + \frac{3s_0 - s_1 + 1}{5}(-2)^n - 2^n.$$

To satisfy ourselves that this has been done correctly, we use the last expression and find

$$s_2 = \frac{2s_0 + s_1 + 4}{5}\,9 + \frac{3s_0 - s_1 + 1}{5}\,4 - 4 = s_1 + 6s_0 + 4\,,$$

which is consistent with the recurrence.

*Example* A.1.3. For the second-order equation

$$s_n = s_{n-1} + 6s_{n-2} + n2^n\,, \tag{A.5}$$

let's first check that there is no constant $c$ for which $w_n = c\,2^n$ is a particular solution, since

$$\begin{aligned} w_{n-1} + 6w_{n-2} + n2^n - w_n \\ &= c2^{n-1} + 6c2^{n-2} + n2^n - c2^n \\ &= 2^{n-1}(c + 3c + 2n - 2c) = 2^n(c + n), \end{aligned}$$

which cannot be zero for all $n$ when $c$ is a constant. Rather, since $\deg(p) = 1$ and $\lambda = 2 \notin \{3, -2\}$ the result says that there exists a particular solution of the form $v_n = (an + b)2^n$ for some constants $a, b$. The recurrence gives

$$\begin{aligned} v_{n-1} + 6v_{n-2} + n2^n - v_n \\ &= (an - a + b)2^{n-1} + 6(an - 2a + b)2^{n-2} + n2^n - (an + b)2^n \\ &= 2^{n-1}((an - a + b) + 3(an - 2a + b) + 2n - 2(an + b)) \\ &= 2^{n-1}((2a + 2)n + (-7a + 2b)), \end{aligned}$$

which must equal zero for all integers $n \geq 0$, that is, $2a + 2 = 0$ and $-7a + b = 0$. This gives $a = -1$, $b = 7a = -7$, and $v_n = -(2n + 7)2^{n-1}$ is a particular solution of (A.5). Therefore, the general solution has the form

$$s_n = a_1 3^n + a_2(-2)^n - (2n + 7)2^{n-1} \text{ for constants } a_1, a_2 \in \mathbb{C}.$$

For initial values $s_0, s_1$ from this we obtain

$$s_0 = a_1 + a_2 - 7/2 \text{ and } s_1 = 3a_1 - 2a_2 - 9\,.$$

Simultaneously solving this system of equations,

$$a_1 = \frac{2s_0 + s_1 + 16}{5} \text{ and } a_2 = \frac{6s_0 - 2s_1 + 3}{10}\,,$$

which gives

$$s_n = \frac{2s_0 + s_1 + 16}{5}3^n - \frac{6s_0 - 2s_1 + 3}{5}(-2)^{n-1} - (2n + 7)2^{n-1}\,.$$

Verifying this calculation for $n = 2$:

$$s_2 = \frac{18s_0 + 9s_1 + 144}{5} + \frac{12s_0 - 4s_1 + 6}{5} - 22 = 6s_0 + s_1 + 8\,,$$

as required.

*Example* A.1.4. For

$$s_n = s_{n-1} + 6s_{n-2} + 3^n,$$

$\deg(p) = 0$ and $\lambda = \lambda_1$, which means $\delta = 1$ and the result gives $v_n = 3^n bn$ for some $b \in \mathbb{C}$. (Note that $w_n = c3^n$ is a solution to the homogeneous equation and so cannot be a particular solution here. Also, $v_n = 3^n(bn + a)$ could be used here, but the constant term can be absorbed into the earlier coefficient of $3^n$.) To find $b$,

$$v_{n-1} + 6v_{n-2} + 3^n - v_n = 3^{n-1}b(n - 1 + 2n - 4 - 3n) + 3^n = 3^{n-1}(-5b + 3)\,,$$

and from this, $b = 3/5$ and $v_n = n3^{n+1}/5$. Therefore,

$$s_n = a_1 3^n + a_2(-2)^n + \frac{n3^{n+1}}{5}\,,$$

and $s_0 = a_1 + a_2$, $s_1 = 3a_1 - 2a_2 + 9/5$ gives

$$a_1 = \frac{2s_0 + s_1 - 9/5}{5}\ ;\ a_2 = \frac{3s_0 - s_1 + 9/5}{5}\,.$$

Hence,

$$s_n = \frac{1}{5}\Big((2s_0 + s_1 - 9/5\ + 3n)3^n + (3s_0 - s_1 + 9/5)(-2)^n\Big).$$

*Example* A.1.5. For

$$s_n = s_{n-1} + 6s_{n-2} + n(-2)^n$$

$\deg(p) = 1$ and $\lambda = \lambda_2$, from which we obtain $\delta = 1$ and $v_n = (-2)^n Q(n)$ where $Q(n) = cn^2 + bn$ for some $b, c \in \mathbb{C}$. Then

$$\begin{aligned} v_{n-1} + 6v_{n-2} + n(-2)^n - v_n \\ =(-2)^{n-1}(Q(n-1) - 3Q(n-2) - 2n + 2Q(n)) \end{aligned}$$

equals zero for all integers $n \geq 2$. In particular, from $n = 2$ and $n = 3$ we obtain

$$\begin{aligned} Q(1) - 3Q(0) + 2Q(2) = 4 \\ Q(2) - 3Q(1) + 2Q(3) = 6\,, \end{aligned}$$

which give $5b + 9c = 4$ and $5b + 19c = 6$, and $c = 1/5$, $b = 11/25$. Therefore, $Q(n) = \frac{5n^2+11n}{25}$ and

$$s_n = a_1 3^n + \Big(a_2 + \frac{5n^2 + 11n}{25}\Big)(-2)^n\,.$$

Substituting $n = 0$ and $n = 1$,

$$s_0 = a_1 + a_2 \quad \text{and} \quad s_1 = 3a_1 - 2a_2 - 32/25$$

and from this

$$a_1 = \frac{2s_0 + s_1 + 32/25}{5}\ ;\ a_2 = \frac{3s_0 - s_1 - 32/25}{5}\,.$$

## A.2 A multiple root

Here we assume $m_2 = \ldots = m_t = 1$, and Theorem 3.2.1 becomes:

**Solving (L) when there is a multiple eigenvalue**

*When $m_1 \geq 2$ and $m_2 = \ldots = m_t = 1$ are the eigenvalues of the recurrence (L), then $v_n = \lambda^n n^\delta q(n)$ is a particular solution where $q$ is a fixed polynomial with* $\deg(q) = \deg(p)$ *and*

$$\delta = \begin{cases} 0 & \text{if } \lambda \notin \{\lambda_1, \ldots, \lambda_t\} \\ m_1 & \text{if } \lambda = \lambda_1 \\ 1 & \text{if } \lambda \in \{\lambda_2, \ldots, \lambda_t\} \end{cases} . \tag{A.6}$$

The first five examples in this section are second-order equations, and each has the characteristic polynomial

$$ch(x) = x^2 - 4x + 4 = (x-2)^2 .$$

This has the double root $\lambda_1 = 2$, which means the general solution is

$$s_n = (a_1 n + a_2)2^n + v_n ,$$

where $v_n$ is a particular solution. As in the last section $v_n$ depends on $\psi$ and $a_1, a_2$ can be calculated from the initial conditions.

*Example* A.2.1. For any initial value problem with

$$s_n = 4s_{n-1} - 4s_{n-2},$$

$s_n = (a_1 n + a_2)2^n$ where the constants $a_1, a_2$ can be computed from the initial conditions $s_0, s_1$ to be $a_2 = s_0$ and $a_1 = (s_1 - 2s_0)/2$. The solution is

$$s_n = ((s_1 - 2s_0)n + 2s_0)2^{n-1} .$$

*Example* A.2.2. Consider the second order equation

$$s_n = 4s_{n-1} - 4s_{n-2} + 3^n,$$

where $\deg(p) = 0$ and $\lambda = 3$ is not an eigenvalue. A particular solution is $v_n = c3^n$ for some constant $c$ and

$$4v_{n-1} - 4v_{n-2} + 3^n - v_n = 3^{n-2}(12c - 4c + 9 - 9c) = 3^{n-2}(9 - c)$$

which equals zero for $c = 9$. This gives $v_n = 3^{n+2}$ and general solution

$$s_n = (a_1 n + a_2)2^n + 3^{n+2}, \text{ for some } a_1, a_2 \in \mathbb{C}.$$

For the initial conditions $s_0, s_1$, computation gives

$$s_n = \Big(\frac{s_1 - 2s_0 - 9}{2} n + (s_0 - 9)\Big)2^n + 3^{n+2}.$$

*Example* A.2.3. The second-order equation

$$s_n = 4s_{n-1} - 4s_{n-2} + 3^n\, n$$

has $\deg(p) = 1$ and $\lambda = 3$ is again not an eigenvalue. Therefore, for constants $a, b$, $v_n = (an + b)3^n$ is a particular solution and

$$\begin{aligned} 4v_{n-1} - 4v_{n-2} + n3^n - v_n \\ &= 3^{n-2}(12(an - a + b) - 4(an - 2a + b) + 9n - 9(an + b)) \\ &= 3^{n-2}((9 - a)n - (4a + b)), \end{aligned}$$

which equals zero for all integers $n \geq 2$ when $a = 9$ and $b = -4a = -36$. Therefore, $v_n = (9n - 36)3^n = (n - 4)3^{n+2}$ is a particular solution, and the general solution has the form

$$s_n = (a_1 n + a_2)2^n + (n - 4)3^{n+2}.$$

For initial values $s_0, s_1$ we obtain

$$s_n = \Big(\frac{s_1 - 2s_0 + 9}{2} n + (s_0 + 36)\Big)2^n + (n - 4)3^{n+2}.$$

*Example* A.2.4. The second-order equation

$$s_n = 4s_{n-1} - 4s_{n-2} + 2^n$$

has $\deg(p) = 0$ and $\lambda = 2$ is the double eigenvalue which gives $\delta = 2$ and $v_n = an^2 2^n$ is a particular solution for some $a \in \mathbb{C}$. Then

$$\begin{aligned} 4v_{n-1} - 4v_{n-2} + 2^n - v_n \\ &= 2^n(2a(n - 1)^2 - a(n - 2)^2 + 1 - an^2) \\ &= 2^n(-2a + 1)\,, \end{aligned}$$

implying $a = \frac{1}{2}$ which means $v_n = n^2 2^{n-1}$ is a particular solution and the general solution therefore has the form

$$s_n = (n^2 + a_1 n + a_2)2^{n-1}.$$

From the initial values $s_0, s_1$ we obtain

$$s_n = (n^2 + (s_1 - 1 - 2s_0)n + 2s_0)2^{n-1}.$$

*Example* A.2.5. The second-order equation

$$s_n = 4s_{n-1} - 4s_{n-2} + 2^n\, n$$

has $\deg(p) = 1$ and $\lambda = 2$ is again the double eigenvalue, giving $\delta = 2$ and particular solution $v_n = 2^n Q(n)$ for $Q(n) = an^3 + bn^2$. Then

$$\begin{aligned} &4v_{n-1} - 4v_{n-2} + n2^n - v_n \\ &\qquad = 2^n[2Q(n-1) - Q(n-2) + n - Q(n)]\,, \end{aligned}$$

which must equal zero for all values of $n \geq 2$. From

$$2Q(n-1) - Q(n-2) + n - Q(n) = 0 \text{ for } n = 2 \text{ and } n = 3\,,$$

we obtain $a = 1/6$ and $b = 3a = 1/2$. Therefore, $v_n = (\frac{1}{3}n^3 + n^2)2^{n-1}$ is a particular solution and the general solution has the form

$$s_n = \Big(\frac{n^3}{3} + n^2 + a_1 n + a_2\Big)2^{n-1}\,,$$

and the initial values $s_0, s_1$ give

$$s_n = \Big(\frac{n^3}{3} + n^2 + (s_1 - 2s_0 - \frac{4}{3})n + 2s_0\Big)2^{n-1}.$$

Since each of the last five examples had only one (double) root, it was impossible to be in the middle alternative of (A.6). The next example illustrates this case.

*Example* A.2.6. The second-order equation

$$s_n = 4s_{n-1} - 5s_{n-2} + 2s_{n-3} + 2^n$$

has $\deg(p) = 0$ and $\lambda = 2$. Since the characteristic polynomial factors as

$$ch(x) = x^3 - 4x^2 + 5x - 2 = (x-1)^2(x-2),$$

$\lambda$ is a simple root of $ch(x)$ and $\delta = 1$, which gives the particular solution $v_n = an2^n$. Then

$$\begin{aligned} &4v_{n-1} - 5v_{n-2} + 2v_{n-3} + 2^n - v_n \\ &\qquad = 2^{n-2}(8a(n-1) - 5a(n-2) + a(n-3) + 4 - 4an) = 2^{n-2}(4-a)\,, \end{aligned}$$

and $v_n = 4n2^n = n2^{n+2}$ is a particular solution with general solution

$$s_n = (a_1 n + a_2) + a_3 2^n + n2^{n+2} = (a_1 n + a_2) + (a_3 + 4n)2^n\,.$$

For initial values $s_0, s_1, s_2$ we obtain

$$s_0 = a_2 + a_3\,;\; s_1 = a_1 + a_2 + 2a_3 + 8\,;\; s_2 = 2a_1 + a_2 + 4a_3 + 32\,,$$

which implies

$$\begin{aligned} s_n = (-2s_0 + 3s_1 - s_2 + 8)n + (2s_1 - s_2 + 16) \\ +(s_0 - 2s_1 + s_2 - 16 + 4n)2^n\,. \end{aligned}$$

# Appendix B
## Complex Numbers

Because the square of a real number cannot be negative, there is no real number whose square is $-1$ and this means that the simple quadratic equation $z^2 + 1 = 0$ has no real roots. Using notation introduced by Leonhard Euler in 1777, we reserve $i$ to mean a symbol for which $i^2 = -1$ holds, and the set of complex numbers is

$$\mathbb{C} = \{ a + bi \ : \ a, b \in \mathbb{R} \} .$$

There is a one-to-one correspondence between each element $a + bi$ of $\mathbb{C}$ and the point $(a, b)$ in the set $\mathbb{R}^2$ in which real numbers $a$ correspond to the points of the form $(a, 0)$ and **imaginary numbers** $0 + bi$ correspond to the points on the $y-$axis. If the operations of addition and scalar multiplication (using real scalars) on $\mathbb{C}$ are defined "coordinatewise", namely by

$$(a_1 + b_1 i) + (a_2 + b_2 i) = (a_1 + a_2) + (b_1 + b_2)i$$

and

$$c(a + bi) = (ca) + (cb)i ,$$

In *La Géometrie* (1637) René Descartes introduced the terms real and imaginary numbers. In 1936, a US mathematician named Arnold Dresden suggested the name "imaginary" numbers be changed to "normal numbers" since the imaginary axis is normal (that is, perpendicular) to the real axis. That term never caught on, and instead has been used for something different: a real number is called **normal** if the digits of its base $b$ expansion behave in a suitably random manner for every base $b$.

the correspondence is actually a vector space isomorphism and $\mathbb{C}$ is a real vector space under these operations, and in particular $\mathbb{C}$ is an **abelian group** under addition. This means that the operation of addition on the set $\mathbb{C}$ is a commutative associative operation for which the real number 0 is the additive identity and every element of $\mathbb{C}$ has an additive identity.

How is multiplication of two complex numbers defined? If we want multiplication to be associative and commutative and also to distribute over addition, then it turns out that there's only one way to define multiplication! This is true because for any $a, b, c, d \in \mathbb{R}$,

$$\begin{aligned}
(a+bi)(c+di) &= (a+bi)c + (a+bi)(di) && \text{distributive law in } \mathbb{C}\\
&= (a+bi)c + ((a+bi)d)i && \text{associative law in } \mathbb{C}\\
&= c(a+bi) + (d(a+bi))i && \text{commutative law in } \mathbb{C}\\
&= (ca+cbi) + (da+dbi)i && \text{scalar multipication in } \mathbb{R}^2\\
&= (ac+bci) + (ad+bdi)i && \text{commutative law in } \mathbb{R}\\
&= (ac+bci) + (adi+bdi^2) && \text{distributive law in } \mathbb{C}\\
&= (ac+bci) + (-bd+adi) && i \text{ satisfies the identity } i^2=-1\\
&= (ac-bd) + (ad+bc)i && \text{definition of addition in } \mathbb{C}.
\end{aligned}$$

This shows that because $\mathbb{C}$ is a real vector space the only associative, commutative and distributive multiplication on $\mathbb{C}$ which extends the scalar multiplication by elements of $\mathbb{R}$ is

$$(a+bi)(c+di) = (ac-bd) + (ad+bc)i\,.$$

(Also note that $(a+0i)(c+0i) = ac$, which says that this multiplication is consistent with real multiplication on the subset $\mathbb{R}$.) This definition of multiplication satisfies the commutative law for multiplication and the real number 1 is the multiplicative identity. As you might already know, the set $\mathbb{C}$ with the operations of addition and multiplication defined above is the **field of complex numbers**. Also, $a$ is called the **real part** and $b$ the **imaginary part** of the complex number $a+bi$.

> On the surface, the multiplication of two complex numbers $a+bi, c+di$ seems to require the four real multiplications $ac, bd, ad, bc$. Check that the auxiliary multiplication $(a+b)(c+d)$ can be used to reduce the number to three.

Until now the geometric representation of complex numbers has been in *rectangular* coordinates. Thinking in terms of polar coordinates, we can rewrite the complex number $z = a+ib$ as $z = |z|(\cos(\theta) + i\sin(\theta))$, where $|z| = \sqrt{a^2+b^2}$ is called the **modulus** and its **argument** $\theta$ is the angle between the positive $x$-axis and the vector $a+bi$. For the moment we write $e(\theta) = \cos(\theta) + i\sin(\theta)$, and

$$z = |z|e(\theta) \text{ for some } 0 \le \theta < 2\pi$$

where $e(\pi) = \cos(\pi) = -1$ and $e(2\pi) = \cos(2\pi) = 1$.

Let's consider the function $e(\theta)$ from another point of view. The trignometric functions $\cos(\theta)$ and $\sin(\theta)$ are functions of the real variable $\theta$ whose power series are known and converge at every real number,

$$\cos(\theta) = \sum_{k\geq 0} \frac{(-1)^k}{(2k)!}\theta^{2k} \quad \text{and} \quad \sin(\theta) = \sum_{k\geq 0} \frac{(-1)^k}{(2k+1)!}\theta^{2k+1}\,.$$

Therefore, for any real number $y$ the complex number $e(y) = \cos(y) + i\sin(y)$ equals

$$e(y) = \sum_{k\geq 0} \frac{1}{(2k)!}(iy)^{2k} + \sum_{k\geq 0} \frac{1}{(2k+1)!}(iy)^{2k+1}\,,$$

since $(-1)^k = i^{2k}$. Adding these series,

$$e(y) = \sum_{k\geq 0} \frac{1}{k!}(iy)^k\,, \tag{B.1}$$

which is reminiscent of the power series expansion of $e^x = \sum_{k\geq 0} x^k/k!$. In fact, we will show that the series *can* be rearranged and so define the **complex exponential function** $e^z$ of the complex variable $z = x + iy$ as

$$e^z = e^x(\cos(y) + i\sin(y))\,. \tag{B.2}$$

Once this is done, we can drop the notation $e(\theta)$ and write $z = |z|e^{i\theta}$, where $\theta$ is the argument of $z$.

For any sequence $< a_i >$ of complex numbers and any complex number $\alpha$ the power series $\sum_{k\geq 0} a_k(z-\alpha)^k$ is said to **converge** at the complex number $z = z_0$ if the sequence of partial sums $\sum_{k=0}^{N} a_k(z_0-\alpha)^k$ converges. We will show that every power series has a **radius of convergence** $R$ (it might be infinite) which has the property that the series $\sum_{k\geq 0} |a_k(z-\alpha)^k|$ converges for all $|z-\alpha| < R$ and diverges for all $|z-\alpha| > R$. Because $e^x = \sum_{k\geq 0} x^k/k!$ converges for all real numbers, the radius of convergence of the complex power series cannot have a finite radius of convergence. This means that $e^z$ is *absolutely* convergent for all complex numbers $z$ and so can be rearranged, and $e^z$ as defined in (B.2) is a well-defined function on $\mathbb{C}$.

We will show that the radius of convergence for a power series $\sum_{k\geq 0} a_k(z-\alpha)^k$ equals the **supremum** of the set

$$\mathcal{S} = \{r \geq 0 \,:\, \text{there exists } M > 0 \text{ such that every } |a_k|r^k \leq M\}$$

by showing the sequence $\sum_{k\geq 0} |a_k(z-\alpha)^k|$ converges for all $|z-\alpha| < R$ and $\sum_{k\geq 0} a_k(z-\alpha)^k$ diverges for all $|z-\alpha| > R$.

If the complex number $z$ satisfies $r = |z-\alpha| > R$, then $r \notin \mathcal{S}$, which means $|a_k||z-\alpha|^k$ is unbounded and the series must diverge. On the other hand, for $r = |z-\alpha| < R$ we can choose $\rho$ such that $r < \rho < R$. Since $\rho \in \mathcal{S}$ there exists $M > 0$ such that all $|a_k\rho^k| < M$ and this gives

$$|a_k(z-\alpha)^k| = |a_k|r^k = |a_k|\rho^k\left(\frac{r}{\rho}\right)^k < M\left(\frac{r}{\rho}\right)^k,$$

where the ratio of the geometric series $\displaystyle\sum_{k\geq 0}\left(\frac{r}{\rho}\right)^k$ satisfies $0 \leq \dfrac{r}{\rho} < 1$. Therefore, by comparison the series $\sum_{k\geq 0} a_k(z-\alpha)^k$ is absolutely convergent on $|z-\alpha| < R$, and this proves that every complex power series has a radius of convergence.

Any power series $\sum_{k\geq 0} a_k(z-\alpha)^k$ with positive radius of convergence can be shown to be a *differentiable* function of a complex variable and its derivative is the power series $\sum_{k\geq 0} ka_k(z-\alpha)^{k-1}$ which has the *same* radius of convergence as the original series. If $R_1$ be the radius of convergence of the series, then $R_1 \leq R$ since $|ka_k| \geq |a_k|$. In particular, $R = 0$ implies $R_1 = 0$, and we may assume $R > 0$. To show $R_1 \geq R$, we'll prove that $R_1 \geq r$ for all $0 < r < R$. The construction of $R$ implies that for any $\rho$ with $r < \rho < R$ there exists $M > 0$ such that $|a_k|\rho^k \leq M$ for all $k$, and

$$|ka_k|r^{k-1} = \frac{|a_k\rho^k|}{r}\cdot k\left(\frac{r}{\rho}\right)^k \leq \frac{M}{r}\cdot k\left(\frac{r}{\rho}\right)^k.$$

Since $0 \leq r < \rho$, then $\lim_{k\mapsto\infty} k\left(\frac{r}{\rho}\right)^k = 0$ and $|ka_k|r^{k-1}$ is bounded, giving $R_1 \geq r$ for all $0 < r < R$. To summarize, for any power series $\sum_{k\geq 0} a_k(z-\alpha)^k$, $f(z) = \sum_{k\geq 0} a_k(z-\alpha)^k$ is a complex-valued function on its disk of convergence and can be proved to be differentiable on the disk with derivative $f'(z) = \sum_{k\geq 0} D(a_k(z-\alpha)^k)$ (where $D$ is the differentiation operator on the space of polynomials). Repeating this argument, $f'(z)$ must also be differentiable on the same disk and $f''(z) = \sum_{k\geq 0} D^2(a_k(z-\alpha)^k)$. From this we see that $f(z)$ is *infinitely* differentiable on its disk of convergence.

Before leaving power series, we define the **Taylor series** of $f$ about $z = \alpha$ as

$$\sum_{k\geq 0}\frac{D^k(f)(\alpha)}{k!}(z-\alpha)^k,$$

where $D^k(f)(\alpha)$ is the $k^{\text{th}}$ derivative of $f(z)$ evaluated at $z = \alpha$. It can be proved that the Taylor series of $f$ converges to $f(z)$ on some disk about $\alpha$ provided that $\sum_{n\geq 0} a_n(z-\alpha)^n$ has a nonzero radius of convergence.

Returning to the complex exponential function, from the trig formulas for the sum of two angles,

$$e^{i(\theta_1+\theta_2)} = [\cos(\theta_1)\cos(\theta_2) - \sin(\theta_1)\sin(\theta_2)] + i[\sin(\theta_1)\cos(\theta_2) + \sin(\theta_2)\cos(\theta_1)]\,.$$

The definition of muliplication can be used to unwrap this identity to get

$$e^{i(\theta_1+\theta_2)} = e^{i\theta_1}e^{i\theta_2}\,,$$

and iteration of this process for any positive integer $m$ gives

$$e^{im\theta} = e^{i\theta+i(m-1)\theta} = e^{i\theta}e^{i(m-1)\theta} = \ldots = (e^{i\theta})^m$$

and the laws of exponents

$$e^{z_1+z_2} = e^{z_1}e^{z_2} \qquad \text{and} \qquad e^{mz} = (e^z)^m\,.$$

When $\theta_0 = 2\pi/m$, this formula becomes $e^{i\theta_0 m} = e^{i2\pi} = 1$ which means that $z = e^{i\theta_0}$ is a root of the equation $z^m - 1 = 0$, and this particular root is called the **principal $\mathbf{m^{th}}$ root of unity**. The complex numbers satisfying $z^m - 1 = 0$ are called the **$\mathbf{m^{th}}$ roots of unity**. Writing the equation $z^m - 1 = 0$ in the form $z^m = 1$, we see that every power of the principal $\mathrm{m}^{\mathrm{th}}$ root of unity $e^{i\theta_0}$ is also an $\mathrm{m}^{\mathrm{th}}$ root of unity, and since $e^{i\theta} = e^{i(\theta+2\pi)}$ always holds this gives the *different* roots $e^{i\theta}$ for

$$\theta = 0\,,\ \frac{2\pi}{m}\,,\ \frac{4\pi}{m}\,,\ \ldots\,,\ \frac{(m-1)\pi}{m}\,.$$

Since these roots are equally separated complex numbers which lie on the unit circle $|z| = 1$ they are often called **cyclotomic** (or "circle dividing"). Because a polynomial equation of degree $m$ has at most $m$ roots in any *field* (refer to Exercise 8.22), these are all the complex roots of $z^m - 1 = 0$ in $\mathbb{C}$. Geometrically, they form the vertices of a regular $m$-gon inscribed in the complex unit circle with one vertex anchored at 1. Algebraically, they form a group $\mathcal{G}_m$ under multiplication, and this group is isomorphic to the integers modulo $m$ under the map

$$e^{ik2\pi/m} \mapsto k \bmod m\,.$$

Among these roots are the **primitive $\mathbf{m^{th}}$ roots of unity**, the ones which are not roots of $z^k - 1 = 0$ for any $0 < k < m$. (The principal root of unity is an example of a primitive root of unity.) The primitive roots of unity also generate the group $\mathcal{G}_m$, in the sense that for any primitive root $\zeta$ its powers

$$\zeta, \zeta^2, \ldots, \zeta^m$$

are all different and therefore every $m^{\mathrm{th}}$ root of unity is a power of $\zeta$. Also note that every $5^{\mathrm{th}}$ root of unity except 1 is primitive, whereas there are only two primitive $6^{\mathrm{th}}$ roots of unity, the principal root $e^{i2\pi/6}$ and its multiplicative inverse $e^{-i2\pi/6}$.

The fact that every polynomial of the form $f(z) = z^m - 1$ (for positive integer $m$) has $m$ complex roots is not an anomaly: *Every* polynomial of degree $m$ has $m$ complex roots (where here roots are counted according to

their multiplicity.) This statement is usually referred to as **The Fundamental Theorem of Algebra**, and is an example of a statement which is very easy to state but difficult to prove. Using the mathematical language of the time, this result was conjectured sometime in the sixteenth century. In the late seventeenth century Newton stated the Fundamental Theorem in fairly modern terms when he wrote that every non-constant polynomial with real coefficients can be factored into a product of linear and quadratic polynomials with real coefficients. The theorem was finally proved by Gauss in his 1799 dissertation, and over his lifetime Gauss published at least three other proofs of the Fundamental Theorem. Each proof is an *existence proof*, since it only proves that such roots exist and doesn't give a method for explicitly constructing any of the roots. In fact, the roots can be difficult or impossible to determine explicitly but we've ignored this important practical problem by choosing polynomials with relatively obvious factorizations.

# Appendix C
## Highlights of Linear Algebra

This appendix contains a review of the linear algebra we use in this book. It is essentially a synopsis of an introductory linear algebra course with a number of topics omitted and some new ones added.

### C.1 Vector Spaces and Subspaces

A **vector space** is a set whose elements can be added together and also multiplied by scalars, which in our case are either real or complex numbers, and $V$ is called a **real vector space** or a **complex vector space** according to whether $\mathbb{R}$ or $\mathbb{C}$ is the field of scalars for $V$. (In general, the set of scalars for a vector space must be a field as defined in Section 8.3.) In order for $V$ to be a vector space the operations of addition and scalar multiplication must satisfy a list of axioms which require that both operations be associative, addition is commutative, and scalar multiplication distributes over multiplication. In addition, there must be an additive identity and every element of $V$ has an additive inverse in $V$. The basic examples of vector spaces are $\mathbb{R}^n$ and $\mathbb{C}^n$ (with scalars from $\mathbb{R}$ and $\mathbb{C}$ respectively).

A nonempty subset $W$ of $V$ is called a **subspace** if $W$ is also a vector space under the operations of $V$. Given a set of vectors $S$ in $V$, we can form **linear combinations** of elements $v_1, \cdots, v_n$ from $S$, which are sums of the form

$$\alpha_1 v_1 + \ldots + \alpha_n v_n \text{ where the } \alpha_i \text{ are scalars .}$$

Every subspace is closed under formation of linear combinations, and it can be proved that any nonempty subset of a vector space which is closed under addition and scaling is a subspace. The subspace of all linear combinations of elements from $S$ is called the **span of** $S$, which we will denote by $\mathrm{Span}(S)$, is the smallest subspace containing $S$. A set $S$ will be called a **spanning set for** $V$ when $\mathrm{Span}(S) = V$, which means every element of $V$ can be written as a linear combination of elements from $S$.

Although we will principally concentrate on subspaces of $\mathbb{R}^n$ and $\mathbb{C}^n$, there is a more general class of vector spaces which we encounter, the vector spaces of real- or complex-valued functions defined on some set $X$. Because a sequence is a function defined on the natural numbers, the complex vector space of all sequences of complex numbers is an example of such a vector space. Both addition and scalar multiplication are defined "componentwise", in other words to add two sequences we add the $n^{\text{th}}$ term of the first sequence to the $n^{\text{th}}$ term of the second and multiplying every element of a sequence by the scalar $\alpha$ gives the scaled sequence. Considering an infinite sequence $< s_n >$ as the function $s(n) = s_n$ defined on $\mathbb{N}$, these componentwise vector space operations are the usual operations for functions,

$$(\alpha s)(n) = \alpha s(n) \text{ and } (s+t)(n) = s(n) + t(n)\,.$$

In general, the set of all functions from any set $X$ into $\mathbb{R}$ or $\mathbb{C}$ forms a vector space under the usual operations of function addition and scaling. When $X = \mathbb{Z}$ we obtain the set of all doubly infinite sequences of real or complex numbers.

## C.2 Linear Independence and Basis

How many different linear combinations yield the same vector? An nonempty set $S$ of vectors is called **linearly independent** if every element in $\mathrm{Span}(S)$ can be expressed as a linear combination of elements of $S$ in only one way. Since $\mathrm{Span}(S)$ is a subspace of $V$, it contains the zero vector. Therefore, linear independence implies that whenever

$$\alpha_1 v_1 + \ldots + \alpha_n v_n = 0 = \beta_1 v_1 + \ldots + \beta_m v_m$$

holds we have $\alpha_i = \beta_i$ holds for all $i$. Since the choice of all $\beta_i = 0$ gives the zero vector, linear independence means that

$$\alpha_1 v_1 + \ldots + \alpha_n v_n = 0 \iff \alpha_i = 0 \text{ for all } i.$$

This condition is often taken as the definition of linear independence.

A nonempty subset $\mathcal{B}$ of a vector space $V$ is called a **basis** for $V$ if it is a linearly independent spanning set for $V$. In other words, $\mathcal{B}$ is a basis

exactly when every vector can be expressed as a linear combination of elements from $\mathcal{B}$ in a unique way. When $V$ has a basis with $n$ elements, we can encode every element as an $n$-tuple. For instance, it can be checked that the set $\mathcal{B} = \{(1,2),(-1,1)\}$ is a basis for $V = \mathbb{R}^2$. The vector $v = (-2,3)$ satisfies

$$v = \frac{1}{3}(1,2) + \frac{7}{3}(-1,1),$$

and therefore its encoding in terms of the basis $\mathcal{B}$ is

$$v_{\mathcal{B}} = \begin{bmatrix} 1/3 \\ 7/3 \end{bmatrix}.$$

In general, when $\mathcal{B} = \{v_1, \ldots, v_n\}$ is a basis for $V$ and $v = \alpha_1 v_1 + \ldots + \alpha_n v_n$ we write the **coordinate vector of v relative to** $\mathcal{B}$ as $v_{\mathcal{B}} = (\alpha_1, \ldots, \alpha_n)^T$.

The **dimension** of a vector space is defined to be the number of elements in a basis. This definition makes sense because any two bases for the same vector space must have the same number of elements. The proof of this fact can be broken into two cases, depending on whether or not the vector space has a *finite* spanning set. In the first case the vector space is called **finite-dimensional**. The proof for this case proceeds by proving that if $V$ has a spanning set with $n$ elements then any subset which has more than $n$ elements must be linearly dependent. In particular, any basis can have at most $n$ elements. If $\mathcal{B}$ and $\mathcal{B}'$ are two bases for $V$, then $\mathcal{B}$ is a spanning set and the linear independence of $\mathcal{B}'$ implies that the number of elements in $\mathcal{B}'$ is bounded above by the number of elements in $\mathcal{B}$. Reversing the roles of $\mathcal{B}$ and $\mathcal{B}'$ yields the desired result, namely, that $\mathcal{B}$ and $\mathcal{B}'$ have the same size. If there is no *finite* spanning set for $V$, the vector space $V$ is called **infinite-dimensional**. The above argument is too simplistic for this case but there is a more sophisticated argument that works. We don't give that argument here.

When $V$ is a finite-dimensional vector space, any subspace of $W$ is also finite-dimensional and the dimension of any *proper* subspace $W \subsetneq V$ is strictly less than the dimension of $V$.

## C.3 Linear Transformations

A linear transformation is a map between two vector spaces (they must have the same field of scalars) which preserves addition and scalar multiplication. In other words, a function $T : V \to W$ is a **linear transformation** if

$$T(\alpha v + v') = \alpha T(v) + T(v') \text{ for all } v, v' \in v \text{ and scalars } \alpha.$$

Because of linearity, a linear transformation is defined by its action on a basis.

Linear ransformations which are functions from a vector space to itself are called **operators**. We can represent a linear operator on an $n$-dimensional vector space $V$ with respect to a basis $\mathcal{B} = \{b_1, \ldots, b_n\}$ by the $n \times n$ matrix whose $i$-th column is $T(b_i)$ as coordinatized by $\mathcal{B}$. We denote this matrix by $[T]_{\mathcal{B}}$. For example, if $T$ is the operator on $\mathbb{R}^3$ defined by $T(x, y, z) = (3x + 2y, z, -z)$ and $\mathcal{B} = \{(1, 0, 0), (1, -1, 0), (0, 0, 1)\}$ then $T(b_1) = (3, 0, 0) = 3b_1$ ; $T(b_2) = (1, 0, 0) = b_1$ ; $T(b_3) = (0, 1, -1) = b_1 - b_2 - b_3$ and

$$[T]_{\mathcal{B}} = \begin{bmatrix} 3 & 1 & 1 \\ 0 & 0 & -1 \\ 0 & 0 & -1 \end{bmatrix}.$$

Notice that there is an inherent order to the elements in the basis, and to emphasize this the term "ordered basis" is normally used. For $A = [T]_{\mathcal{B}}$ we have the helpful identity

$$T(v)_{\mathcal{B}} = Av_{\mathcal{B}} \text{ for all } v \in \mathbb{C}^n,$$

and $T$ is an invertible operator if and only if $A$ is an invertible matrix. Therefore, if we are given two pieces of data—an ordered basis $\mathcal{B}$ for an $n$-dimensional vector space and a linear operator $T$ on $V$—we can then obtain an $n \times n$ matrix $A$ with the property that a coordinate vector for $T(v)$ relative to $\mathcal{B}$ is the result of multiplying $A$ by the coordinate vector for $v$ relative to $\mathcal{B}$.

If $\mathcal{B} = \{b_1, \ldots, b_n\}, \mathcal{C} = \{c_1, \ldots, c_n\}$ are two ordered bases for $\mathbb{C}^n$, consider the matrix $P$ whose $i$-th column is the coordinatization $[c_i]_{\mathcal{B}}$. (The matrix $P$ is often called the **change of basis matrix** from $\mathcal{B}$ to $\mathcal{C}$.) Since the associated linear operator (the identity transformation) is invertible, the matrix $P$ is invertible and

$$T(v)_{\mathcal{C}} = P^{-1}APv_{\mathcal{C}};$$

that is, the representation of $T$ in the basis $\mathcal{C}$ is the matrix $P^{-1}AP$. Two matrices $A$ and $B$, which are related by a matrix equation of the form

$$B = P^{-1}AP \text{ where } P \text{ is invertible,}$$

are called **similar matrices**, and they represent the same linear transformation with respect to two different bases.

## C.4 Eigenvectors

A nonzero vector $v \in V$ is called an **eigenvector** of the linear operator $T$ associated with the **eigenvalue** $\lambda$ if $T(v) = \lambda v$. If $T$ is represented as a matrix $A$ with respect to a convenient basis $\mathcal{B}$, the eigenvectors of $T$ are

exactly the eigenvectors of $A$ and they can be found by first computing all solutions $\lambda$ to $\det(A - \lambda I) = 0$ and then for each solution $\lambda_0$ finding all (eigenvectors) $v$ satisfying $(A - \lambda_0 I)v = 0$.

Eigenvectors corresponding to different eigenvalues are linearly independent. If $V$ has a basis $\mathcal{B}_1$ of eigenvectors, the matrix representing $T$ in the basis $\mathcal{B}_1$ is a diagonal matrix (and the diagonal elements are the respective eigenvalues), which means that if $P$ is the change of basis matrix from $\mathcal{B}_1$ to $\mathcal{B}$ then $P^{-1}AP$ is a diagonal matrix. A square matrix $A$ is called **diagonalizable** if $A$ is similar to a diagonal matrix. We summarize this very nice situation in the following theorem whose proof can be found in [67, Chapter 6].

**Theorem C.4.1.** *Let $A$ be an $n \times n$ matrix with complex entries and* **characteristic polynomial** *$ch_A(x) = \det(A - Ix)$. Let $\lambda_1, \lambda_2, \ldots, \lambda_n$ be the eigenvalues of $A$, the complex roots of $ch_A(x) = 0$ repeated according to multiplicity.*

*(a) The matrix $A$ is diagonalizable if and only if $\mathbb{C}^n$ has a basis of eigenvectors.*

*(b) If all the $\lambda_i$ are distinct then $A$ is diagonalizable.*

*(c) Suppose $A$ is diagonalizable and $\{\boldsymbol{v}_1, \ldots, \boldsymbol{v}_n\}$ is a basis of eigenvectors, and form the $n \times n$ matrix $P$ whose $i^{th}$ column is $\boldsymbol{v}_i$. Then $P^{-1}AP$ is the diagonal matrix $D$ with diagonal entries $\lambda_1, \ldots, \lambda_n$.*

*(d) Real symmetric matrices are always diagonalizable.*

For us the most useful application of diagonalizability is that it allows the powers of $A$ to be computed quickly since $P^{-1}AP = D$ yields $A^n = PD^nP^{-1}$ for all $n \geq 0$. Not all square matrices are diagonalizable, but every square matrix $A$ with complex entries has a **Jordan form**. This means there exists an invertible matrix $P$ with the property that $P^{-1}AP$ is composed of **Jordan blocks** on the diagonal, where a Jordan block is a bi-diagonal matrix of the form

$$\begin{bmatrix} \lambda & 1 & 0 & \cdots & 0 \\ 0 & \lambda & 1 & \ldots & 0 \\ 0 & 0 & \lambda & \ddots & 0 \\ & \cdots & & \ddots & 1 \\ 0 & 0 & 0 & \cdots & \lambda \end{bmatrix} \text{ for some } \lambda.$$

For example, the matrix on the right is in Jordan form. It has three Jordan blocks: one of size 2 with $\lambda = 3$, one of size 1 with $\lambda = 3$, and one of size 1 with $\lambda = 2$. The Jordan form of $A$ is unique up to the order of the individual Jordan blocks, the $\lambda$ appearing in the blocks of the Jordan form are the eigenvalues of $A$, and the minimal and

$$\begin{bmatrix} \boxed{\begin{matrix} 3 & 1 \\ 0 & 3 \end{matrix}} & \begin{matrix} 0 & 0 \\ 0 & 0 \end{matrix} \\ \begin{matrix} 0 & 0 \\ 0 & 0 \end{matrix} & \begin{matrix} \boxed{3} & 0 \\ 0 & \boxed{2} \end{matrix} \end{bmatrix}$$

characteristic polynomials of $A$ can be found from $J$. The proofs of these facts and more information on Jordan form can be found in [67, Section 7.3]. Since $A^n = PJ^nP^{-1}$, powers of $A$ are relatively easy to compute because the $n^{\text{th}}$ power of a Jordan block is given by the upper triangular matrix

$$\begin{bmatrix} \lambda & 1 & 0 & \cdots & 0 \\ 0 & \lambda & 1 & \ldots & 0 \\ 0 & 0 & \lambda & \ddots & 0 \\ & \ldots & & \ddots & 1 \\ 0 & 0 & 0 & \cdots & \lambda \end{bmatrix} = \begin{bmatrix} \lambda^n & n\lambda^{n-1} & \ldots & \binom{n}{l}\lambda^{n-l} \\ 0 & \lambda^n & \ldots & \binom{n}{l-1}\lambda^{n-l+1} \\ 0 & 0 & \ddots & \vdots \\ 0 & 0 & \ldots & \lambda^n \end{bmatrix}. \tag{C.1}$$

## C.5 Characteristic and Minimal Polynomials

As in Theorem C.4.1 the characteristic polynomial of the $n \times n$ matrix $A$ is defined as $ch_A(x) = \det(A - Ix)$. For instance, for $A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$, the characteristic polynomial is $ch_A(x) = x^2 + 1$. Note that when the matrix $A$ is plugged into its characteristic polynomial,

$$ch_A(A) = A^2 + I = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix},$$

the zero matrix. This property holds in general.

**Theorem C.5.1 (The Cayley-Hamilton Theorem ).** *For any $n \times n$ complex matrix $A$, $ch_A(A)$ equals the zero matrix.*

There is another important polynomial associated with a square matrix $A$. ¿From the Cayley-Hamilton Theorem we know $ch_A(x)$ is an example of a non-zero polynomial $p(x) \in \mathbb{C}[x]$ for which $p(A)$ is the zero matrix, and the **minimal polynomial** $min_A(x)$ of $A$ is defined to be the non-zero polynomial of *smallest* degree (and leading coefficient equal to 1) such that $p(A) = 0$. Dividing $ch_A$ by $min_A$ gives polynomials $q(x), r(x)$ such that

$$ch_A(x) = q(x)min_A(x) + r(x) \text{ where } \deg(r(x)) < \deg(min_A(x)).$$

Since $ch_A(A) = 0 = min_A(A)$, then also $r(A) = 0$ and the the minimality of $\deg(min_A)$ forces the remainder $r(x)$ to be the zero polynomial. With this we've proved the useful fact that the minimal polynomial always divides the characteristic polynomial.

# Appendix D
## Roots in the Unit Circle

Our analysis of difference equations shows that the general solution of a recurrence converges to zero when all roots of the characteristic polynomial are less than 1 in absolute value. Since some roots of the characteristic polynomial might be nonreal, this condition means that all roots lie within the unit circle in the complex plane. In what follows we describe a method of Morris Marden [92, Chapter X] for calculating the number of roots of a polynomial within the unit circle. We specialize his technique to polynomials with real coefficients and recast it as an algorithm.

Before discussing the general method, consider the special example of nonnegative polynomials, which is the leading case for applications. Recall that a **nonnegative polynomial** is a polynomial $p$ of the form

$$p(x) = x^k - c_1 x^{k-1} - c_2 x^{k-2} - \ldots - c_{k-1} x - c_k \,,$$

where all $c_i \geq 0$ and $c_k > 0$. In Section 5.1 we prove some special properties of nonnegative polynomials. Among these, Theorem 5.1.3 says any nonnegative polynomial has exactly one positive real root $\lambda_0$ which is **dominant** in the sense that any other root $\lambda$ satisfies $|\lambda| \leq |\lambda_0|$. Further, from Corollary 5.1.5 when $p$ is **primitive** (that is, when $\gcd\{i \mid c_i > 0\} = 1$) $\lambda_0$ is **strictly dominant**, the only root whose modulus is the maximum. This result can be applied to construct polynomials whose roots lie inside the unit circle. For instance, if $q(x)$ is a real polynomial for which

$$p(x) = (x-1)q(x) = x^k - c_1 x^{k-1} - c_2 x^{k-2} - \ldots - c_{k-1} x - c_k$$

is nonnegative then $\lambda_0 = 1$ must be the dominant root of $p$, and when $p$ is a primitive polynomial then $\lambda_0 = 1$ a strictly dominant root of $p$ and all roots

of $q$ therefore lie inside the unit circle. For example, if $q(x) = x^2 + \frac{1}{2}x + \frac{1}{2}$, then $(x-1)q(x) = x^3 - \frac{1}{2}x^2 - \frac{1}{2}$ is a primitive nonnegative polynomial, all roots of $q$ lie inside the unit circle, and the general solution of the associated recurrence $s_n = \frac{1}{2}s_{n-1} + \frac{1}{2}s_{n-3}$ converges to zero.

Of course, in this simple example it's easy to check that $q$ has two complex roots and the absolute value of each is $1/\sqrt{2}$. For a more complicated example, consider the **generalized Fibonacci polynomial**, $p(x) = x^k - x^{k-1} - \ldots - x - 1$ for integer $k \geq 2$. Since $p(1) \leq -1 < 0 < p(2)$, the dominant root $\lambda_0$ lies in the open interval $(1, 2)$. Dividing $p(x)$ by $(x - \lambda_0)$ gives the polynomial

$$q(x) = x^{k-1} + (\lambda_0 - 1)x^{k-2} + \ldots + (\lambda_0^{k-1} - \ldots - 1)\,,$$

and set

$$\begin{aligned} P(x) := (x-1)q(x) = x^k &+ (\lambda_0 - 2)x^{k-1} + (\lambda_0^2 - 2\lambda_0)x^{k-2} \\ &+ \ldots + (\lambda_0^{k-1} - 2\lambda_0^{k-2})x - (\lambda_0^{k-1} - \ldots - 1)\,. \end{aligned}$$

From $p(\lambda_0) = 0$ we obtain $\lambda_0(\lambda_0^{k-1} - \lambda_0^{k-2} - \ldots - 1) = 1$ and so the constant term in $P$ is $-\lambda_0^{-1} < 0$. Since $1 < \lambda_0 < 2$ the other coefficents are also strictly negative and we see that $P$ is a primitive nonnegative polynomial which implies all roots of $q$ must lie inside the unit circle. Since $p(x) = (x - \lambda_0)q(x)$, from this we obtain that all roots of $p$ except $\lambda_0$ are within the unit circle.

## D.1 Marden's method

We now turn to Marden's method which can be applied to any polynomial with real coefficients. The basic idea involves a kind of pairing of $f(x) = a_0 + a_1x + \ldots + a_nx^n$ with its **reciprocal polynomial** $f^R(x) = a_0x^n + a_1x^{n-1} + \ldots + a_n$. The relationship between $f$ and $f^R$ is such that for every nonzero root $r$ of $f$ inside the unit circle there's a corresponding root $r^{-1}$ of $f^R(x)$ outside the unit circle, and conversely. This is true because

$$f(x) = a \cdot x^k \prod_{i=1}^{n-k}(x - r_i) \implies f^R(x) = A \cdot \prod_{i=1}^{n-k}(x - 1/r_i)\,,$$

where $A = a \cdot \prod_{i=1}^{n-k} r_i$.

For any $f$, Marden constructs the associated polynomial

$$\begin{aligned} G(x) = a_0 f(x) - a_n f^R(x) = \\ a_0(a_0 + a_1x + \ldots + a_nx^n) - a_n(a_0x^n + a_1x^{n-1} + \ldots + a_n), \end{aligned}$$

whose degree is at most $n-1$ and whose constant term is $a_0^2 - a_n^2$. He then streamlines some results of A. Cohn [22] and obtains Lemma 42.1 which we reword as

**Theorem D.1.1.** *Suppose $f(x) = a_0 + a_1 x + \ldots + a_n x^n$ has $p$ roots inside the unit circle. If $\delta(f) := a_0^2 - a_n^2$ is nonzero then $G(x) = a_0 f(x) - a_n f^R(x)$ has either $p$ or $n-p$ roots inside the unit circle and the sign of its constant term $\delta(f)$ determines which holds. Namely, $G$ has $p$ roots inside the unit circle if and only if $\delta(f) > 0$. Further, $f$ and $G$ have the same number of roots on the unit circle.*

Therefore, the number of roots of the constructed polynomial $G$ which lie inside the unit circle is the same as the number of roots of either $f$ or $f^R$ within the unit circle and the sign of $\delta(f)$ indicates which holds. For our current purposes we'll say $G$ **is equivalent to** $f$ when $G$ and $f$ have the same number of roots within the unit circle. (Otherwise, $G$ is equivalent to $f^R$.)

For example, let $f(x) = x^2 - \frac{1}{3}x - \frac{1}{3}$. Then $f$ has two roots inside the unit circle, and its reciprocal polynomial $f^R(x) = 1 - \frac{1}{3}x - \frac{1}{3}x^2$ has no roots inside the unit circle. The constructed polynomial is

$$\begin{aligned} G(x) &= a_0 f(x) - a_2 f^R(x) \\ &= -\frac{1}{3}(x^2 - \frac{1}{3}x - \frac{1}{3}) - 1(-\frac{1}{3}x^2 - \frac{1}{3}x + 1) \\ &= \frac{4}{9}x - \frac{8}{9} \end{aligned}$$

and $\delta(f) = -\frac{8}{9} < 0$, which means $G$ is equivalent to $f^R$. Since the only root of $G$ is $x = 2$ which lies outside the unit circle, $G$ is indeed equivalent to $f^R$. Neatly enough, since $\delta(G) \neq 0$, in order to determine if $G$ has a root inside the unit circle we could have applied the same calculation to $G$. For this, we calculate $G^R(x) = \frac{4}{9} - \frac{8}{9}x$ and construct

$$-\frac{8}{9}G(x) - \frac{4}{9}G^R(x) = \left(\frac{8}{9}\right)^2 - \left(\frac{4}{9}\right)^2 = 3\left(\frac{4}{9}\right)^2,$$

a constant polynomial which has $\delta(G) > 0$ and so is equivalent to $G$. Therefore, $G$ (and also $f^R$) has no roots inside the unit circle.

The iteration used in this last example is the basis for Marden's algorithm for counting the number of roots inside the unit circle. At the start of the procedure we set $f_0 = f$, which may have up to $\deg(f) = n$ roots within the unit circle. The degree of the constructed polynomial $f_1(x) = a_0 f_0(x) - a_n f_0^R(x)$ is at most $n-1$. Provided $\delta(f_1) \neq 0$ we can continue to use $f_1$ to construct $f_2$, and so forth.

Notice how this technique can be used to count the number of roots. When the procedure starts we have $f(x)$, which may have up to $n$ roots

within the unit circle. The constructed polynomial $a_0 f - a_n f^R$ has at most $n-1$ roots. If the constructed polynomial $f_1$ has the same number of roots as $f^R$ within the unit circle, then $f^R$ can only have at most $n-1$ roots within the circle, and hence must have at least 1 root outside the circle. But this root of $f^R$ which is outside the circle must correspond to a root of $f$ which is inside the circle. So when $f_1$ is equivalent to $f^R$ we can increase the count of $f$'s roots inside the circle by 1. On the other hand, if $f_1$ agrees with $f$ then we know that $f$ has at most $n-1$ roots within the unit circle and we can proceed with counting the roots of $f_1$ to get the count of the number of roots of $f$. One does not change the count in this situation.

As mentioned earlier, the sign of the constant term of $f_1$ indicates whether it is equivalent to $f$ or is equivalent to $f^R$. So the algorithm will have a switch, called DELTA in the version below, to keep track of whether the current polynomial is equivalent to $f$ or $f^R$. This switch will be updated when a new polynomial is constructed and will also tell the algorithm when to increment the count.

The algorithm breaks down when some $\delta(f_i)$ is zero. For this Marden uses a procedure $EQUIV(f_i)$ to obtain an equivalent polynomial whose degree is at most $\deg(f)$. Before explaining this procedure we give our implementation of Marden's algorithm.

**Marden's Algorithm**

INPUT: $f(z) = a_0 + a_1 z + \ldots + a_n z^n$,
a polynomial with real coefficients.
OUTPUT: COUNT, the number of roots of $f$
within the unit circle.

FOR $s = 0$ TO $n$
$a_s^{(0)} = a_s$; DELTA $= 1$; COUNT $= 0$
FOR $j = 0$ TO $n - 1$
IF $|a_0^{(j)}| = |a_{n-j}^{(j)}|$
THEN EQUIV$(a_0^{(j)} + \ldots + a_{n-j}^{(j)} x^{n-j})$
and update $j$ if necessary.

FOR $k = 0$ TO $n - j - 1$
$a_k^{(j+1)} = a_0^{(j)} a_k^{(j)} - a_{n-j}^{(j)} a_{n-j-k}^{(j)}$

DELTA $=$ DELTA $* sgn(a_0^{(j+1)})$
IF DELTA $< 0$ THEN COUNT $=$ COUNT $+ 1$

OUTPUT (COUNT)

As written, the time for the algorithm is $O(n^2)$ and uses $O(n^2)$ space where the space is used for storing a two-dimensional array to hold the various $a$'s. By slightly changing the order in which the $a$'s are calculated and writing over the old $a$'s with the new $a$'s, the space usage can be cut to $O(n)$. The $O(n^2)$ run time assumes that products and differences can be computed in constant time. This may not be true. If one uses "real" computer arithmetic, then there may be catastrophic loss of accuracy and extended precision may be necessary. Conversely, if rational or even integer arithmetic is used, the number of digits needed could double at each iteration, again forcing the use of extended precision.

Now let us turn to the case when we reach a stage where the constructed polynomial $f_i$ has $\delta(f_i) = 0$. In this situation we want to obtain an equivalent polynomial $EQUIV(f_i)$ [92, Section 45]. Suppose we encounter

$$g(x) := f_i(x) = a_0 + a_1 x + \ldots + a_m x^m \text{ with } \delta(g) = a_0^2 - a_m^2 = 0\,.$$

Then $a_m/a_0 = \pm 1$ which we call $u$ and the equivalent polynomial $EQUIV(g)$ will be determined according to whether or not $(a_m, \ldots, a_0)$ equals $(ua_0, \ldots, ua_m)$.

*Case* 1. If $(a_m, \ldots, a_0) = (ua_0, \ldots, ua_m)$, replace $g$ with

$$EQUIV(g) = a_1 x^{m-1} + 2a_2 x^{m-2} + \ldots + (m-1)a_{m-1} x + m\, a_m\,,$$

the reciprocal function of the derivative of $g$.

*Case* 2. If $(a_m, \ldots, a_0) \neq (ua_0, \ldots, ua_m)$, let $0 < q < m$ be the smallest subscript such that $a_{m-q} \neq ua_q$. Set $b = (a_{m-q} - ua_q)/a_m$ and

$$G(x) = (x^q + 2b/|b|)g(x) = \sum_{i=0}^{m+q} B_i x^i$$

with

$$EQUIV(g) = B_0 G(x) - B_{m+q} G^R(x)\,,$$

a polynomial whose degree can be shown to be at most $\deg(g)$.

*Example* D.1.1. Use Marden's algorithm to count the number of roots of $f(x) = -7x^3 + x^2 + 5x + 1$ inside the unit circle.
$n = 3$ DELTA=1; COUNT=0
$j = 0$

$k = 0$ $\quad a_0^{(1)} = a_0^{(0)} a_0^{(0)} - a_3^{(0)} a_3^{(0)} = 1 \cdot 1 - (-7) \cdot (-7) = -48$
$k = 1$ $\quad a_1^{(1)} = a_0^{(0)} a_1^{(0)} - a_3^{(0)} a_2^{(0)} = 1 \cdot 5 - (-7) \cdot 1 = 12$
$k = 2$ $\quad a_2^{(1)} = a_0^{(0)} a_2^{(0)} - a_3^{(0)} a_1^{(0)} = 1 \cdot 1 - (-7) \cdot 5 = 36$
DELTA $= 1 \cdot (-1) = -1$
COUNT $= 0 + 1 = 1$

$j = 1$

$k = 0$ $\quad a_0^{(2)} = a_0^{(1)} a_0^{(1)} - a_2^{(1)} a_2^{(1)} = (-48) \cdot (-48) - (36) \cdot (36) = 1008$
$k = 1$ $\quad a_1^{(2)} = a_0^{(1)} a_1^{(1)} - a_2^{(1)} a_1^{(1)} = -48 \cdot 12 - 36 \cdot 12 = -1008$
DELTA $= -1 \cdot (1) = -1$
COUNT $= 1 + 1 = 2$

$j = 2$
At this point, $|a_0^{(2)}| = |a_1^{(2)}|$, and $EQUIV(f_2)$ must be invoked. Continuing with our algorithm, Case 1 applies and gives $EQUIV(1008 - 1008x) = -1008$. Updating to $j = 3$, the FOR loop terminates and the algorithm outputs COUNT= 2 as the number of roots of $-7x^3 + x^2 + 5x + 1$ inside the unit circle. Notice that the root $x = 1$ is not *inside* the unit circle and is not counted.

*Example* D.1.2. Find the number of roots of $f(x) = x^2 + x + 1$ which lie inside the unit circle.
$n = 2$ DELTA=1; COUNT=0
$j = 0$

Since $a_0^{(0)} = a_2^{(0)}$, $EQUIV(f)$ must be found. Case 1 again applies and

$$EQUIV(x^2 + x + 1) = a_1 x + 2a_2 = x + 2\,.$$

Since the degree of this polynomial is $\deg(f) - 1$, $j$ must be increased by 1 and $a_0^{(1)} = 2$, $a_1^{(1)} = 1$.

$j = 1$

$$k = 0 \qquad a_0^{(2)} = a_0^{(1)} a_0^{(1)} - a_1^{(1)} a_1^{(1)} = 4 - 1 = 3 > 0\,.$$

DELTA remains positive and COUNT is not incremented. The FOR loop terminates and 0 is returned as the number of roots inside the unit circle for the polynomial $x^2 + x + 1$. This is correct because the two roots of this polynomial are on the unit circle.

*Example* D.1.3. We consider $f(x) = x^2 - x - 1$, the characteristic polynomial for the Fibonacci sequence. As we've noted many times, $f$ has one root inside the unit circle and one root outside the unit circle. Applying Marden's algorithm we find $a_0^{(0)} = -1$ and $a_2^{(0)} = 1$, giving $u = a_2/a_0 = -1$. But $(a_0, a_1, a_2) = (-1, -1, -1) \neq -(a_2, a_1, a_0)$ and Case 2 with $q = 1$ applies. Then $b = -2$,

$$G(x) = (x^q - 2)g(x) = (x - 2)(x^2 - x - 1) = x^3 - 3x^2 + x + 2\,,$$

and

$$EQUIV(g) = 2G(x) - G^R(x) = -7x^2 + 5x + 3\,.$$

Since $\deg(EQUIV) = \deg(f)$, $j = 0$ is unchanged.

$j = 0$

$$k = 0 \qquad a_0^{(1)} = a_0^{(0)} a_0^{(0)} - a_2^{(0)} a_2^{(0)} = 9 - 49 = -40 < 0$$
$$k = 1 \qquad a_1^{(1)} = a_0^{(0)} a_1^{(0)} - a_2^{(0)} a_1^{(0)} = 15 - (-35) = 50$$
$$\text{DELTA} = 1 \cdot (-1) = -1$$
$$\text{COUNT} = 0 + 1 = 1$$

$j = 1$

$$k = 0 \qquad a_0^{(2)} = a_0^{(1)} a_0^{(1)} - a_1^{(1)} a_1^{(1)} = (-40)(-40) - (50)(50) < 0$$
$$\text{DELTA} = -1 \cdot (-1) = 1$$
$$\text{COUNT} = 1$$

The FOR loop terminates and the algorithm correctly outputs COUNT$= 1$ for the number of roots of $x^2 - x - 1$ inside the unit circle.

*Example* D.1.4. As a final example we consider $f(x) = x^2 - 1$ which has the roots $\pm 1$. Applying Marden's algorithm, we find $\delta(f) = 0$, which means that $EQUIV(f)$ must be invoked. Since $(a_2, a_1, a_0) = -(a_0, a_1, a_2)$, Case 1 of the replacement algorithm applies and $EQUIV(f) = a_1 x + 2a_2 = -2$. This results in the degree decreasing by 2, and $j$ must be incremented by 2. But then the FOR loop terminates and the algorithm correctly outputs COUNT=0.

## D.2 Exercises

**Ex D.1.** Use Marden's algorithm to show that $5x^n - 1$ has $n$ roots inside the unit circle.

**Ex D.2.** Show that $x^{n-1} + 2x^{n-2} + 3x^{n-3} + \ldots + (n-1)x + n$ has no roots inside the unit circle by applying Marden's algorithm to $x^n + x^{n-1} + \ldots + x + 1$.

**Ex D.3.** Use Marden's algorithm twice to determine the number of roots inside the unit circle and the number of roots inside the circle of radius 2 for the polynomial $2x^3 - 3x^2 + 2x + 2$. (The actual roots are $-\frac{1}{2}$, $1 + i$, $1 - i$.)

# Index

abelian group, 216, 346
absolute convergence, 347
absolute error convergence, 111
absolute value, 54, 55, 300
adjacency matrix, 179
algorithm, 243
    analysis of, 244
    divide-and-conquer, 253
analysis of algorithms, 244
analysis, asymptotic, 6
aperiodic orbit, 293
aperiodic polynomial, 96
approximation, linear, 300
argument of complex number, 346
arithmetic model of computing, 254
Artin's conjecture, 222
asymptotic analysis, 6
asymptotic behavior, 25
asymptotic property, 292
asymptotic size, 97
asymptotically periodic matrix, 136
attractive cylce, 292
attractive fixed point, 292
basis, 352
Big-Oh, 6
Big-Omega, 6
Big-Omicron, 6
Big-Theta, 6, 245
bijection, 13
binary operation, 251
Binet's Formula, 5, 29, 65
binomial coefficient, 62
    generalized, 49
Binomial Theorem, 62, 181
    Generalized, 49
biorthogonal, 26
Birthday Problem, 235, 240
Bisection Method, 114
bit vector representation, 182
Boolean matrix, 180
butterfly effect, 293

Caesar, Julius, 253
Carmichael numbers, 233
Catalan number, 75
Cauchy sequence, 54
    of polynomials, 55
Cayley-Hamilton Theorem, 356

ceiling function, xi
change of basis matrix, 354
chaos, 293, 312
characteristic of a field, 236
characteristic polynomial
    of a matrix, 174, 355
    of a recurrence, 18
column eigenvector, 26
companion matrix, 137
    of a polynomial, 15
    of a recurrence, 15, 223
completeness, 55
complex number, xi, 346
    imaginary part, 346
    real part, 346
convergence
    absolute, 347
    absolute error, 111
    relative error, 111
convergence of power series
    radius of, 347
convergent series, 347
convolution, 36, 43, 53, 76, 254
coordinatization of vectors, 353
cost criterion
    logarithmic, 249
    uniform, 247
countable set, 296
CPU time, 244
cycle, 292
    attractive, 292
    locally stable, 299
cycle length, 292
cyclic subspace, 22
cyclotomic number, 349
cyclotomic polynomial, 311

density function, 314
depth-first search, 191
derangement, 88
Descartes' Rule of Signs, 92, 93
deterministic generators, 228
deviation of an approximation, 99
diagonalizable, 16, 355
    orthogonally, 26
diagonalization, 308
difference equation
    finite, 33
dimension, 353
    lattice, 229
directed graph, 179
    acyclic, 186
discrete dynamical system, 289
Discrete Fourier Transform, 267
distribution, invariant, 313
divide-and-conquer algorithm, 253
Division Algorithm, 237
division algorithm
    for polynomials, 69
dominant eigenvalue, 27
dominant root, 94, 357
dynamical system, discrete, 289

eigenmatrix, 140
eigenvalue, 354
    dominant, 27
    of a linear fractional system, 307
    of the recurrence, 18
    strictly dominant, 27
eigenvector, 354
    column, 26
    row, 26
equivalence class, 77
equivalence mod 1, 294
ergodic theorem, 317
Euclidean Algorithm, 132, 193, 212
    for polynomials, 70, 84
Euler Phi Function, 311
eventually periodic, 209
exponential function
    complex, 347
exponential generating function, 88
exponential power series, 56

factorization of integers, 232
fast exponentiation, 171, 189, 220, 232
Fast Fourier Transform, 79, 264

Fermat's Last Theorem, 219
Fermat's Little Theorem, 220
fertility rates, 128
Fibonacci, 2
Fibonacci numbers
    programs for, 4
Fibonacci sequence, 1, 2, 14, 63, 91
    generalized, 87, 91, 103, 113, 123
    modular, 208
field
    finite, 216
    of complex numbers, 346
finite difference equation, 33
finite field, xi, 216
finite-dimensional vector space, 353
finiteness argument, 44
fixed point, 208
    attractive, 292
    globally stable, 303
    locally stable, 297
    repelling, 292
    stable, 288
    unstable, 288
floor function, xi
forcing function, 34
formal derivative, 48
formal power series, 43
Fourier matrix, 269
    inverse, 285
Fourier Transform Method, 79
full solution, 52
function, 33
    bijective, 13
    generating, 62
    one-to-one, 13
    onto, 13
Fundamental Theorem of Algebra, 350

Galois field, 218
Gauss, Carl Friedrich, 94
gcd, 124
generalized Fibonacci polynomial, 358
generalized Fibonacci sequence, 87, 91, 103, 113
generating function, 62
    exponential, 88
    reversion of, 79
geometric power series, 48
globally stable fixed point, 303
golden section, 5
graph
    directed, 179
    directed acyclic, 186
    strongly connected, 181
greatest common divisor, 212
    computing, 194
    for polynomials, 84
group, 237
    abelian, 216, 346

Hamilton circuit problem, 192
Horner's method, 115, 141, 162
hyperplane, 42

imaginary numbers, 345
independence, linear, 28, 352
index of imprimitivity, 95, 150
infinite-dimensional vector space, 353
initial conditions, 3
initial value problem, 3, 11
input function, 34
integer, xi
integers modulo m, xi
integers, factorization of, 232
integral domain, 45, 47
interpolation
    formula, 265
    process, 265
interval map, 292
invariant distribution, 313
invariant property, 291
inverse Fourier matrix, 285
inversive generator, 229
invertible element, 212

irreducible polynomial, 218
isomorphism
    vector space, 13
iterate, 208, 292
iterative program, 4

Jacobian matrix, 300
Jordan Canonical Form, 16, 172
Julius Caesar, 253

L'Hôpital's Rule, 85
lattice dimension, 229
Lattice Test, 229
Leslie matrix, 130
Leslie normal form, 203
Leslie's Convergence Theorem, 133
Leslie's model, 129
Leverier's Lemma, 167
Liapunov, 303
Liber Abbaci, 1
linear approximation, 300
linear combination, 351
linear congruential generator, 229
linear independence, 28, 352
linear operator, 287
linear recurrence, 3, 306, 337
linear system, 306
linear transformation, 353
locally stable cycle, 299
locally stable fixed point, 297
logarithmic cost criterion, 249
Lucas numbers, 8, 52, 240
Lucas, Édouard, 8

map, 289
    interval, 292
Markov chain, 176
matrix
    0-1, 180
    asymptotically periodic, 136
    Boolean, 180
    change of basis, 354
    companion, 15
    generalized Vandermonde, 25, 29, 266
    Jacobian, 300
    Vandermonde, 19, 266
matrix form, 16
matrix multiplication, 170, 189
matrix transpose, 151
MERGESORT, 262
metric space, 55
    complete, 55
minimal polynomial, 174, 356
mod 1, 294
modular operations, 207
modulus of complex number, 346
Monte Carlo Method, 232
Montgomery multiplication, 214
multiplication
    matrix, 170, 189
    polynomial, 254, 260
    power series, 43

natural number, xi
neighborhood, 300
neutrally stable points, 306
Newton's method, 116, 262, 321
next node representation, 182
nonnegative matrix, 175
nonnegative polynomial, 92, 357
nonnegative recurrence, 91, 97
normal numbers, 345

one-to-one function, 13
onto function, 13
operator, 354
    linear, 287
orbit, 21, 208
    aperiodic, 293
    period of, 208
    periodic, 208
order of a polynomial, 54
order of a recurrence, 33
order of an element, 219, 237
order of the matrix, 223
orthogonally diagonalizable, 26
oscillate, 288

partial fraction decomposition, 69

partial sum, 44
partition, 77
path, 179
    simple, 179
path length, 179
perfect shuffle, 238
period, 80
    of an orbit, 208
periodic matrix, 135
periodic orbit, 208
periodic point, 208
periodic points, 292
periodic polynomials, 96
periodic sequence, 80
permutation, 238
    derangement, 88
permutation matrix, 151, 268
Perron-Frobenius Theorem, 175
Pisano, Leonardo, 1
plot
    time, 290
    web, 290
Pollard's Rho Method, 232
polynomial
    aperiodic, 96
    cyclotomic, 311
    dense, 264
    nonnegative, 92, 357
    primitive, 96
    reciprocal, 66, 358
    sparse, 264
polynomial multiplication, 254, 260
power series, 43
    convergence of, 347
    exponential, 56
    formal derivative of, 48
    geometric, 48
    partial sum of, 44
    quadratic formula for, 52
primitive element, 221
Primitive Element Theorem, 221
primitive matrix, 131, 175
    algorithms for, 188
primitive polynomial, 96, 357
primitive root of unity, 79
principal root of unity, 79
product rule, 48, 57
program
    iterative, 4
    recursive, 3
pseudoprime, 233
pseudorandom number generation, 228
pseudorandom numbers, 228

quadratic formula, 52
quadratic reciprocity, 226

Rabbit problem, 1
radius of convergence, 347
random algorithm, 277
Rational Canonical Form, 21
rational number, xi
real number, xi
reciprocal, 263
reciprocal polynomial, 66, 84, 358
recurrence
    linear, 3, 337
    linear homogeneous, 11
    nonnegative, 91, 97
recurrence relation, 3
recursive definition, 3
recursive program, 4
reducible polynomial, 237
relative error convergence, 111
relatively prime integers, 211
repelling, 292
Reversion Formula, 83
reversion of generating functions, 79
robustness, 250
root
    dominant, 94, 357
    strictly dominant, 96
root of unity, 79, 95, 265, 310, 349
    primitive, 79
    principal, 79, 265
Round, 91
roundability, 6
Rounding Theorem, 103

row eigenvector, 26
RSA cryptosystem, 232
running time, 244
    average case, 274
    best case, 274
    worse case, 274

Sarkovskii's Theorem, 292
search
    depth-first, 191
sequence, 2
    Cauchy, 54
    periodic, 80
shift operator, 41
sign change, 93
similar matrices, 354
simple path, 179
sink, 186
source, 186
square roots, 121, 263
stable age distribution, 133
stable fixed point, 288
steady state, 292
Stirling numbers of the second kind, 77
Strassen's method, 190, 261
strictly dominant eigenvalue, 27
strictly dominant root, 96, 357
strings, 59
    commuting, 283
    nullstring, 59
Strong Fermat Test, 233
strongly connected graph, 131, 181
Sturm's Theorem, 94
subspace, 351
    cyclic, 22
superstable point, 306
supremum of a set, 347
survival rate, 128
synthetic division, 115

Taylor series, 56, 348
time
    CPU, 244
    running, 244
    unitless, 245
    wall clock, 244
time plot, 290
Tolstoy, 287
total population, 163
Towers of Hanoi puzzle, 246
trace of a matrix, 167
Turing machines, 253, 291

unary operation, 251
uniform cost criterion, 247
unstable fixed point, 288

Vandermonde matrix, 19, 264, 266
    generalized, 25, 29, 266
vector space, 351
    complex, 351
    finite-dimensional, 353
    infinite-dimensional, 353
    real, 351

wall clock time, 244
weather prediction, 295
web plot, 290

Z-transforms, ix
zero divisor, 52