



Concerning Diagonal Similarity of Irreducible Matrices

D. J. Hartfiel

Proceedings of the American Mathematical Society, Vol. 30, No. 3. (Nov., 1971), pp. 419-425.

Stable URL:

<http://links.jstor.org/sici?sici=0002-9939%28197111%2930%3A3%3C419%3ACDSOIM%3E2.0.CO%3B2-Y>

Proceedings of the American Mathematical Society is currently published by American Mathematical Society.

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/about/terms.html>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/ams.html>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

The JSTOR Archive is a trusted digital repository providing for long-term preservation and access to leading academic journals and scholarly literature from around the world. The Archive is supported by libraries, scholarly societies, publishers, and foundations. It is an initiative of JSTOR, a not-for-profit organization with a mission to help the scholarly community take advantage of advances in technology. For more information regarding JSTOR, please contact support@jstor.org.

CONCERNING DIAGONAL SIMILARITY OF IRREDUCIBLE MATRICES

D. J. HARTFIEL

ABSTRACT. If $A = (a_{ij})$ is an $n \times n$ irreducible matrix, then there are positive numbers d_1, d_2, \dots, d_n so that $\sum_k d_i a_{ik} d_k^{-1} = \sum_k d_k a_{ki} d_i^{-1}$ for each $i \in \{1, 2, \dots, n\}$. Further, the numbers d_1, d_2, \dots, d_n are unique up to scalar multiples.

Introduction. Sinkhorn and Knopp in [4] as well as Brualdi, Parter, and Schneider in [1] have shown that if A is a fully indecomposable matrix then there are diagonal matrices D_1 and D_2 with positive main diagonals so that $D_1 A D_2$ is doubly stochastic. Further, in each paper D_1 and D_2 are shown to be unique up to scalar multiples.

In this paper we prove what is considered the analogue of the above result in terms of irreducible matrices. That is we show that if A is an $n \times n$ irreducible matrix then there is a diagonal matrix D , with positive main diagonal, so that

$$\sum_k d_i a_{ik} d_k^{-1} = \sum_k d_k a_{ki} d_i^{-1}$$

for each $i \in \{1, 2, \dots, n\}$. Further we show that D is unique up to scalar multiples.

Definitions and notation. Let $n \geq 2$ be an integer. Let $N = \{1, 2, \dots, n\}$. An $n \times n$ nonnegative matrix A is said to be reducible if there is a permutation matrix P so that

$$P A P^T = \begin{pmatrix} A_1 & 0 \\ B & A_2 \end{pmatrix}$$

where A_1 and A_2 are square. If A is not reducible we say that A is irreducible. By agreement each 1×1 matrix is irreducible. Denote

$$m(A) = \min_{\phi \neq M \subseteq N} \sum_{i \in M; j \notin M} a_{ij}, \quad M(A) = \sum_{i \in N; j \in N} a_{ij}.$$

Fiedler [2] refers to $m(A)$ as a measure of the irreducibility of A . It should be clear that A is irreducible if and only if $m(A) > 0$. Let

$$\Lambda_n = \left\{ n \times n \text{ nonnegative matrices } A \text{ with } \sum_k a_{ik} = \sum_k a_{ki} \text{ for each } i \in N \right\}$$

Received by the editors November 6, 1970.

AMS 1969 subject classifications. Primary 1530; Secondary 1560.

Key words and phrases. Irreducible, diagonally similar, matrix patterns.

Copyright © 1971, American Mathematical Society

and

$\mathfrak{D} = \{(d_1, d_2, \dots, d_n) \text{ so that } d_k > 0$
 for each $k \in N$, and at least one $d_k = 1\}$.

For A irreducible we denote by $S(A)$ a positive number so that $S(A) \cdot m(A) - M(A) > M(A)$. Further we define

$$f_A(d_1, d_2, \dots, d_n) = \max_{\phi \neq M \pm N} \left| \sum_{i \in M; j \in N} d_i a_{ij} d_j^{-1} - \sum_{i \in N; j \in M} d_i a_{ij} d_j^{-1} \right|$$

$$= \max_{\phi \neq M \pm N} \left| \sum_{i \in M; j \in M} d_i a_{ij} d_j^{-1} - \sum_{i \in M; j \in M} d_i a_{ij} d_j^{-1} \right|$$

where each $d_k > 0, k \in N$.

Results.

LEMMA 1. *If $(d_1, d_2, \dots, d_n) \in \mathfrak{D}$ and $\max_{k \in N} d_k \geq S^n(A)$, then*

$$f_A(d_1, d_2, \dots, d_n) > f_A(1, 1, \dots, 1).$$

PROOF. Reorder (d_1, d_2, \dots, d_n) to $(d'_1, d'_2, \dots, d'_n)$ where $d'_1 \geq d'_2 \geq \dots \geq d'_n$. Let s denote the smallest positive integer such that $d'_s / d'_{s+1} > S(A)$. That there is such an s follows since if for each $k \in \{1, \dots, n-1\}$, we have that $d'_k / d'_{k+1} \leq S(A)$, then $d'_1 / d'_n = \prod_{k=1}^{n-1} (d'_k / d'_{k+1}) \leq S^{n-1}(A)$. But $d'_1 \geq \dots \geq d'_n$, and some $d'_k = 1$ implies $d'_n \leq 1$. Hence $d'_1 \leq S^{n-1}(A) d'_n \leq S^{n-1}(A)$. Since $S(A) > 2M(A)/m(A) \geq 2, S^{n-1}(A) < S^n(A)$. It would therefore follow that $\max_k d_k = d'_1 < S^n(A)$, a contradiction. Let $M = \{i_R \mid d_{i_R} = d'_{i_R} \text{ for each } R \in \{1, 2, \dots, s\}\}$. Then

$$f_A(1, 1, \dots, 1) < M(A) < S(A) \cdot m(A) - M(A)$$

$$\leq (d'_s)(d'_{s+1})^{-1} \sum_{i \in M; j \in M} a_{ij} - \sum_{i \in M; j \in M} a_{ij}$$

$$\leq \sum_{i \in M; j \in M} d_i a_{ij} d_j^{-1} - \sum_{i \in M; j \in M} d_i a_{ij} d_j^{-1} \leq f_A(d_1, d_2, \dots, d_n).$$

LEMMA 2. *If $(d_1, d_2, \dots, d_n) \in \mathfrak{D}$ and $\min_{k \in N} d_k \leq S^{-n}(A)$ then*

$$f_A(d_1, d_2, \dots, d_n) > f_A(1, 1, \dots, 1).$$

PROOF. If for each $k \in \{1, \dots, n-1\}$ we have that $d'_k / d'_{k+1} \leq S(A)$, it would follow as in the proof of Lemma 1 that $\min_k d_k = d'_n > S^{-n}(A)$. Hence for this case there is also a smallest integer s such that $d'_s / d'_{s+1} > S(A)$. The rest of the proof is identical with that of Lemma 1.

LEMMA 3. $f_A(\lambda d_1, \lambda d_2, \dots, \lambda d_n) = f_A(d_1, d_2, \dots, d_n)$ for $\lambda > 0$.

LEMMA 4. $f_A(d_1, d_2, \dots, d_n)$ achieves a minimum for some $(d_1, d_2, \dots, d_n) \in \mathcal{D}$.

PROOF. Since $S(A) > 2$, $(1, 1, \dots, 1)$ belongs to the compact set $\mathcal{K} = \{(d_1, \dots, d_n) \in \mathcal{D} \mid S^{-n}(A) \leq \min_k d_k \leq \max_k d_k \leq S^n(A)\}$. By Lemmas 1 and 2, $\text{Inf}_{\mathcal{D}} f_A = \text{Inf}_{\mathcal{K}} f_A$, and the result follows.

LEMMA 5. Let $(d_1, d_2, \dots, d_n) \in \mathcal{D}$. Suppose

$$f_A(d_1, d_2, \dots, d_n) = \max_{\phi \neq M \subseteq N} \left| \sum_{i \in M; j \in M} d_i a_{ij} d_j^{-1} - \sum_{i \in M; j \in M} d_i a_{ij} d_j^{-1} \right|$$

is achieved on $M_0 = \{i_1, i_2, \dots, i_s\}$. Then

$$\left\{ \sum_k d_i a_{ik} d_k^{-1} - \sum_k d_k a_{ki} d_i^{-1} \mid i \in M_0 \right\}$$

contains no sign changes and $\left\{ \sum_k d_i a_{ik} d_k^{-1} - \sum_k d_k a_{ki} d_i^{-1} \mid i \notin M_0 \right\}$ contains no sign changes.

PROOF. Let $M'_0 = \{i \in M_0 \mid \sum_k d_i a_{ik} d_k^{-1} - \sum_k d_k a_{ki} d_i^{-1} > 0\}$ and $M''_0 = \{i \in M_0 \mid \sum_k d_i a_{ik} d_k^{-1} - \sum_k d_k a_{ki} d_i^{-1} < 0\}$. If both M'_0 and M''_0 are nonvoid, then we would have

$$\begin{aligned} & \left| \sum_{i \in M'_0; k \in N} d_i a_{ik} d_k^{-1} - \sum_{i \in M'_0; k \in N} d_k a_{ki} d_i^{-1} \right| \\ & < \max \left[\sum_{i \in M'_0; k \in N} d_i a_{ik} d_k^{-1} - \sum_{i \in M'_0; k \in N} d_k a_{ki} d_i^{-1}, \right. \\ & \qquad \qquad \qquad \left. \sum_{i \in M''_0; k \in N} d_k a_{ki} d_i^{-1} - \sum_{i \in M''_0; k \in N} d_i a_{ik} d_k^{-1} \right], \end{aligned}$$

a contradiction. The proof of the other half of the lemma is similar.

THEOREM 1. If A is irreducible, then there is a

$$D = \text{diagonal } (d_1, d_2, \dots, d_n)$$

with $(d_1, d_2, \dots, d_n) \in \mathcal{D}$ so that $DAD^{-1} \in \Lambda_n$.

PROOF. We first prove the theorem for a positive matrix A . Suppose $f_A(d_1, d_2, \dots, d_n)$ achieves a minimum at $(d_1^0, d_2^0, \dots, d_n^0) \in \mathcal{D}$. Suppose

$$f_A(d_1^0, d_2^0, \dots, d_n^0) = \left| \sum_{i \in M_0; j \in N} d_i^0 a_{ij} (d_j^0)^{-1} - \sum_{i \in N; j \in M_0} d_i^0 a_{ij} (d_j^0)^{-1} \right|.$$

We shall prove that $f_A(d_1^0, \dots, d_n^0) = 0$.

Then $\sum_{i \in M; j \in N} d_i^0 a_{ij} (d_j^0)^{-1} - \sum_{i \in N; j \in M} d_i^0 a_{ij} (d_j^0)^{-1} = 0$ for any M where $\phi \neq M \subseteq N$. The result follows by taking $M = \{k\}$, for each $k \in \{1, \dots, n\}$. Hence suppose that $f_A(d_1^0, \dots, d_n^0) > 0$. Without loss of generality suppose $d_i^0 a_{ij} (d_j^0)^{-1} = b_{ij}$ and $\sum_{k \in N} b_{ik} - \sum_{k \in N} b_{ki} > 0$ for each $i \in M_0$. Let $M_1 = \{i \mid \sum_{k \in N} b_{ik} = \sum_{k \in N} b_{ki}\}$ and $M_2 = N - (M_0 \cup M_1)$.

In view of Lemma 5, it must be that $\sum_k b_{ik} - \sum_k b_{ki} < 0$ for all $i \in M_2$. In particular, M_2 is nonempty.

Consider (d_1, d_2, \dots, d_n) defined as follows:

$$\begin{aligned} d_k &= (1 + \epsilon)^{-1} && \text{if } k \in M_0, \\ &= 1 && \text{if } k \in M_1, \\ &= (1 + \epsilon) && \text{if } k \in M_2, \text{ where } 0 < \epsilon < 1. \end{aligned}$$

Now suppose $M'_1 \subseteq M_1$. Then consider

$$\begin{aligned} g(\epsilon) &= \sum_{i \in M_0 \cup M'_1; j \in N} d_i b_{ij} d_j^{-1} - \sum_{i \in N; j \in M_0 \cup M'_1} d_i b_{ij} d_j^{-1} \\ &= \sum_{i \in M_0; j \in M_1} (1 + \epsilon)^{-1} b_{ij} + \sum_{i \in M_0; j \in M_2} (1 + \epsilon)^{-2} b_{ij} - \sum_{i \in M_1; j \in M_0} (1 + \epsilon) b_{ij} \\ &\quad - \sum_{i \in M_2; j \in M_0} (1 + \epsilon)^2 b_{ij} + \sum_{i \in M'_1; j \in M_0} (1 + \epsilon) b_{ij} + \sum_{i \in M'_1; j \in M_1} b_{ij} \\ &\quad + \sum_{i \in M'_1; j \in M_2} (1 + \epsilon)^{-1} b_{ij} - \sum_{i \in M_0; j \in M'_1} (1 + \epsilon)^{-1} b_{ij} \\ &\quad - \sum_{i \in M_1; j \in M'_1} b_{ij} - \sum_{i \in M_2; j \in M'_1} (1 + \epsilon) b_{ij}. \end{aligned}$$

Hence $g'(0) < 0$. Therefore there is a number ϵ_1 so that, if $0 < \epsilon < \epsilon_1$,

$$\max_{M'_1} \left(\sum_{i \in M_0 \cup M'_1; k \in N} d_i b_{ik} d_k^{-1} - \sum_{i \in M_0 \cup M'_1; k \in N} d_k b_{ki} d_i^{-1} \right) < f_A(d_1^0, d_2^0, \dots, d_n^0)$$

and

$$\sum_{k \in N} d_i b_{ik} d_k^{-1} - \sum_{k \in N} d_k b_{ki} d_i^{-1} > 0 \quad \text{for } i \in M_0.$$

Similarly, for $M''_1 \subseteq M_1$, consider

$$\begin{aligned}
 h(\epsilon) &= - \sum_{i \in M_2 \cup M_1'; j \in N} d_i b_{ij} d_j^{-1} + \sum_{i \in N; j \in M_2 \cup M_1''} d_i b_{ij} d_j^{-1} \\
 &= \sum_{i \in M_0; j \in M_2} (1 + \epsilon)^{-2} b_{ij} + \sum_{i \in M_1; j \in M_2} (1 + \epsilon)^{-1} b_{ij} \\
 &\quad - \sum_{i \in M_2; j \in M_0} (1 + \epsilon)^2 b_{ij} - \sum_{i \in M_2; j \in M_1} (1 + \epsilon) b_{ij} \\
 &\quad + \sum_{i \in M_0; j \in M_1'} (1 + \epsilon)^{-1} b_{ij} + \sum_{i \in M_1; j \in M_1''} b_{ij} \\
 &\quad + \sum_{i \in M_2; j \in M_1'} (1 + \epsilon) b_{ij} - \sum_{i \in M_1'; j \in M_0} (1 + \epsilon) b_{ij} \\
 &\quad - \sum_{i \in M_1'; j \in M_1} b_{ij} - \sum_{i \in M_1'; j \in M_2} (1 + \epsilon)^{-1} b_{ij}.
 \end{aligned}$$

Hence $h'(0) < 0$. Therefore there is a number ϵ_2 so that, if $0 < \epsilon < \epsilon_2$,

$$\max_{M_1'} \left(\sum_{i \in M_2 \cup M_1'; k \in N} d_k b_{ki} d_i^{-1} - \sum_{i \in M_2 \cup M_1'; k \in N} d_i b_{ik} d_k^{-1} \right) < f_A(d_1^0, d_2^0, \dots, d_n^0)$$

and

$$\sum_{k \in N} d_k b_{ki} d_i^{-1} - \sum_{k \in N} d_i b_{ik} d_k^{-1} > 0 \quad \text{for } i \in M_2.$$

Therefore for $\epsilon_3 = \min(\epsilon_1, \epsilon_2)$ and $0 < \epsilon < \epsilon_3$ we have in view of Lemma 5, any set \bar{M} for which

$$f_A(d_1, \dots, d_n) = \left| \sum_{i \in \bar{M}; j \in N} d_i a_{ij} d_j^{-1} - \sum_{i \in N; j \in \bar{M}} d_i a_{ij} d_j^{-1} \right|$$

must necessarily be one of the two forms $M_0 \cup M_1'$ or $M_2 \cup M_1''$ for some $M_1' \subseteq M_1$ or $M_1'' \subseteq M_1$. Therefore

$$f_A(d_1 d_1^0, d_2 d_2^0, \dots, d_n d_n^0) < f_A(d_1^0, d_2^0, \dots, d_n^0).$$

But $(1, d_1^{-1}(d_1^0)^{-1} d_2 d_2^0, \dots, d_1^{-1}(d_1^0)^{-1} d_n d_n^0) \in \mathfrak{D}$, and since, by Lemma 3,

$$f(1, d_1^{-1}(d_1^0)^{-1} d_2 d_2^0, \dots, d_1^{-1}(d_1^0)^{-1} d_n d_n^0) = f(d_1 d_1^0, d_2 d_2^0, \dots, d_n d_n^0),$$

f_A would not be minimal at (d_1^0, \dots, d_n^0) . This contradiction shows that $f_A(d_1^0, \dots, d_n^0) = 0$ as was to be proven.

Now suppose A is irreducible. Let A_m be a positive matrix for each positive integer m , and $\lim_{m \rightarrow \infty} A_m = A$. Then for each m there is a diagonal matrix $D_m \in \mathfrak{D}$ so that $D_m A_m D_m^{-1} \in \Lambda_n$. In the proof of Lemma 4, $S^{-n}(A_m) \leq \min_k d_k^{(m)} \leq \max_k d_k^{(m)} \leq S^n(A_m)$. Prudent choices for each

$S(A_m)$ indicate that the $d_k^{(m)}$ are bounded from above and away from zero as $m \rightarrow \infty$. Hence $\{D_m\}$ has a limit point D and D is nonsingular. Let $\{D_{m'}\}$ be a subsequence of $\{D_m\}$ so that $\lim_{m' \rightarrow \infty} D_{m'} = D$. Then $\lim_{m' \rightarrow \infty} D_{m'} A_{m'} D_{m'}^{-1} = DAD^{-1}$ and the result follows.

COROLLARY 1. *If for some permutation matrix P , P^TAP is a direct sum of irreducible matrices then there is a $D = \text{diagonal}(d_1, d_2, \dots, d_n)$ with $(d_1, d_2, \dots, d_n) \in \mathfrak{D}$ such that $DAD^{-1} \in \Lambda_n$.*

We might remark here that the result essentially characterizes irreducibility, in the sense that if A is a nonnegative $n \times n$ matrix then there exists a $D = \text{diag}(d_1, \dots, d_n)$, $(d_1, \dots, d_n) \in \mathfrak{D}$ and $DAD^{-1} \in \Lambda_n$, if and only if there exists a permutation matrix P such that P^TAP is a direct sum of irreducible matrices. The converse part of this statement follows readily from the normal form of a reducible matrix [3, p. 74].

THEOREM 2. *If A is irreducible, $A \in \Lambda_n$, $B \in \Lambda_n$ and D is a diagonal matrix with positive main diagonal so that $DAD^{-1} = B$, then $A = B$ and $D = \lambda I$ for some $\lambda > 0$.*

PROOF. Without loss of generality we may assume

$$D = \text{diagonal}(d_1, d_2, \dots, d_n) \quad \text{where } d_1 \geq d_2 \geq \dots \geq d_n.$$

If $d_1 = d_n$ then we are through. If $d_1 \neq d_n$, then let s denote the smallest integer so that $d_s > d_{s+1}$. Let $M = \{1, 2, \dots, s\}$. If $i \in M$, $k \notin M$, $d_i > d_k$ and so $d_i a_{ik} d_k^{-1} \geq a_{ik}$, the last inequality is strict for some $i \in M$, $k \notin M$ due to the irreducibility of A . Thus

$$\sum_{i \in M; k \notin M} d_i a_{ik} d_k^{-1} > \sum_{i \in M; k \notin M} a_{ik} = \sum_{i \notin M; k \in M} a_{ik}.$$

Similarly $\sum_{i \notin M; k \in M} a_{ik} > \sum_{i \notin M; k \in M} d_i a_{ik} d_k^{-1}$ and, therefore, $\sum_{i \in M; k \notin M} d_i a_{ik} d_k^{-1} > \sum_{i \notin M; k \in M} d_i a_{ik} d_k^{-1}$, i.e.

$$\sum_{i \in M; k \notin M} d_i a_{ik} d_k^{-1} > \sum_{i \in N; k \in M} d_i a_{ik} d_k^{-1}$$

which gives us a contradiction. Therefore $d_1 = d_n$ and $D = \lambda I$ for some $\lambda > 0$.

COROLLARY 2. *If A is irreducible and D a diagonal matrix with $DAD^{-1} \in \Lambda_n$, then D is unique up to a scalar multiple.*

ACKNOWLEDGEMENT. The author is grateful for the referee's suggestions concerning the presentation of this paper.

BIBLIOGRAPHY

1. R. A. Brualdi, S. V. Parter, and H. Schneider, *The diagonal equivalence of a nonnegative matrix to a stochastic matrix*, J. Math. Anal. Appl. **16** (1966), 31–50. MR **34** #5844.
2. M. Fiedler, *Bounds for eigenvalues of doubly stochastic matrices* (submitted).
3. F. R. Gantmaher, *The theory of matrices*, GITTL, Moscow, 1953; English transl., Vol. 2, Chelsea, New York, 1959. MR **16**, 438.
4. R. Sinkhorn and P. Knopp, *Concerning nonnegative matrices and doubly stochastic matrices*, Pacific J. Math. **21** (1967), 343–348. MR **35** #1617.

TEXAS A&M UNIVERSITY, COLLEGE STATION, TEXAS 77843