

The Repulsive Lattice Gas, the Independent-Set Polynomial, and the Lovász Local Lemma

Alexander D. Scott
Department of Mathematics
University College London
London WC1E 6BT, England
scott@math.ucl.ac.uk

Alan D. Sokal
Department of Physics
New York University
4 Washington Place
New York, NY 10003 USA
sokal@nyu.edu

September 15, 2003
revised September 8, 2004

Abstract

We elucidate the close connection between the repulsive lattice gas in equilibrium statistical mechanics and the Lovász local lemma in probabilistic combinatorics. We show that the conclusion of the Lovász local lemma holds for dependency graph G and probabilities $\{p_x\}$ if and only if the independent-set polynomial for G is nonvanishing in the polydisc of radii $\{p_x\}$. Furthermore, we show that the usual proof of the Lovász local lemma — which provides a sufficient condition for this to occur — corresponds to a simple inductive argument for the nonvanishing of the independent-set polynomial in a polydisc, which was discovered implicitly by Shearer [98] and explicitly by Dobrushin [37, 38]. We also present some refinements and extensions of both arguments, including a generalization of the Lovász local lemma that allows for “soft” dependencies. In addition, we prove some general properties of the partition function of a repulsive lattice gas, most of which are consequences of the alternating-sign property for the Mayer coefficients. We conclude with a brief discussion of the repulsive lattice gas on countably infinite graphs.

Key Words: Graph, lattice gas, hard-core interaction, independent-set polynomial, polymer expansion, cluster expansion, Mayer expansion, Lovász local lemma, probabilistic method.

Contents

1	Introduction	3
1.1	The repulsive lattice gas	4
1.2	The Lovász local lemma	6
1.3	Plan of this paper	8
2	The lattice-gas partition function	8
2.1	Elementary identities	9
2.2	The Mayer expansion	10
2.3	The fundamental theorem	17
2.4	Properties of the set $\mathcal{R}(W)$	22
2.5	Further consequences of the alternating-sign property	28
2.6	Algebraic irreducibility of $Z_W(\mathbf{w})$	33
2.7	Convexity of $\log Z$ at nonnegative fugacity	35
3	The lattice gas with hard-core self-repulsion	36
3.1	The fundamental identity	36
3.2	Examples	38
3.3	Reduction formulae	42
3.4	When G is a tree	43
3.5	Upper bounds on $\mathcal{R}(W)$ when G_W is a tree	45
4	The Lovász local lemma	49
4.1	Hard-core version	49
4.2	Soft-core version	53
5	Sufficient conditions for $Z_W \neq 0$ in a polydisc	56
5.1	Basic bound	57
5.2	Improved bound	61
5.3	Optimal bound	64
6	Tree interpretation	66
6.1	Tree interpretation of Theorem 5.6	66
6.2	Tree interpretation of the “optimal” bound: hard-core case	68
6.3	Tree interpretation of the “optimal” bound: general case	70
7	Unfolding	71
8	Infinite graphs	73
8.1	General properties	74
8.2	Infinite trees	75
8.3	Regular lattices	77
8.4	Quantitative bounds for the lattice \mathbb{Z}^d	85
8.4.1	Upper bounds on λ_c	86
8.4.2	Lower bounds on λ_c	88

1 Introduction

The lattice gas with repulsive pair interactions is an important model in equilibrium statistical mechanics [36, 46, 59, 57, 60, 95, 6, 7, 12, 55, 26, 110, 61, 50, 102, 70, 71, 43]. In the special case of a hard-core self-repulsion and hard-core nearest-neighbor exclusion (i.e. no site can be multiply occupied and no pair of adjacent sites can be simultaneously occupied), the partition function of the lattice gas coincides with the independent-set polynomial in combinatorics (also known as the independence polynomial or the stable-set polynomial) [4, 41, 42, 56, 51, 52, 53, 63, 27, 28, 64, 77, 35]. Moreover, the hard-core lattice gas is the *universal* statistical-mechanical model in the sense that any statistical-mechanical model living on a vertex set V_0 can be mapped onto a gas of nonoverlapping “polymers” on V_0 , i.e. a hard-core lattice gas on the intersection graph of V_0 [100, Section 5.7]. This construction, which is termed the “polymer expansion” or “cluster expansion”, is an important tool in mathematical statistical mechanics [97, 29, 47, 31, 24], and much effort has been devoted to finding complex polydiscs in which the polymer expansion is convergent, i.e. in which the (polymer-)lattice-gas partition function is nonvanishing [88, 33, 97, 29, 72, 30, 32, 100, 37, 38, 89, 90, 31, 82, 25, 101, 102, 104]. One goal of this paper is to make a modest contribution to this line of development.

The Lovász local lemma [39, 40, 105, 106] is an important tool in probabilistic existence proofs in combinatorics. It provides a lower bound on the probability that none of a collection of “bad” events occurs, when those events are subject to a set of “local” dependencies, controlled by a “dependency graph”. The Lovász local lemma (and its algorithmic versions [8, 3]) has found a significant variety of applications, especially in graph coloring, Ramsey theory, and related algorithmic problems [2, 23, 83].

In this paper we would like to elucidate the close relation between these two apparently disparate subjects. Following a seminal (but apparently little-known) paper of Shearer [98], we shall show that the conclusion of the Lovász local lemma holds for dependency graph G and probabilities $\{p_x\}$ if and only if the independent-set polynomial for G is nonvanishing in the polydisc of radii $\{p_x\}$. Moreover, we shall show that the usual proof of the Lovász local lemma — which provides a *sufficient* condition for this to occur — corresponds to a simple inductive argument for the nonvanishing of the independent-set polynomial in a polydisc, which was discovered implicitly by Shearer [98] and explicitly by Dobrushin [37, 38]. Finally, we shall present some refinements and extensions of both arguments.

We have tried hard to make this paper comprehensible to the union (not the intersection!) of combinatorialists, probabilists, classical analysts and mathematical physicists. We apologize in advance to experts in each of these fields for boring them every now and then with overly detailed explanations of elementary facts.

1.1 The repulsive lattice gas

In statistical mechanics, a “grand-canonical gas” is defined by a *single-particle state space* X (here a nonempty finite set), a *fugacity vector* $\mathbf{w} = \{w_x\}_{x \in X} \in \mathbb{C}^X$, and a *two-particle Boltzmann factor* $W: X \times X \rightarrow \mathbb{C}$ with $W(x, y) = W(y, x)$. The (grand) partition function $Z_W(\mathbf{w})$ is then defined to be the sum over ways of placing $n \geq 0$ “particles” on “sites” $x_1, \dots, x_n \in X$, with each configuration assigned a “Boltzmann weight” given by the product of the corresponding factors w_{x_i} and $W(x_i, x_j)$:

$$Z_W(\mathbf{w}) = \sum_{n=0}^{\infty} \frac{1}{n!} \sum_{x_1, \dots, x_n \in X} \left(\prod_{i=1}^n w_{x_i} \right) \left(\prod_{1 \leq i < j \leq n} W(x_i, x_j) \right) \quad (1.1a)$$

$$= \sum_{\mathbf{n}} \left(\prod_{x \in X} \frac{w_x^{n_x} W(x, x)^{n_x(n_x-1)/2}}{n_x!} \right) \left(\prod_{\{x, y\} \subseteq X} W(x, y)^{n_x n_y} \right) \quad (1.1b)$$

where in (1.1b) the sum runs over all multi-indices $\mathbf{n} = \{n_x\}_{x \in X}$ of nonnegative integers, and the product runs over all two-element subsets $\{x, y\} \subseteq X$ ($x \neq y$). In this paper we shall limit attention to the *repulsive* case in which $0 \leq W(x, y) \leq 1$ for all x, y . From this assumption it follows immediately that $Z_W(\mathbf{w})$ is an entire analytic function of \mathbf{w} satisfying $|Z_W(\mathbf{w})| \leq \exp(\sum_{x \in X} |w_x|)$.

If $W(x, x) = 0$ for all $x \in X$ — in statistical mechanics this is called a *hard-core self-repulsion* — then the only nonvanishing terms in (1.1b) have $n_x = 0$ or 1 for all x (i.e. each site can be occupied by at most one particle), so that $Z_W(\mathbf{w})$ can be written as a sum over subsets:

$$Z_W(\mathbf{w}) = \sum_{X' \subseteq X} \left(\prod_{x \in X'} w_x \right) \left(\prod_{\{x, y\} \subseteq X'} W(x, y) \right). \quad (1.2)$$

In this case $Z_W(\mathbf{w})$ is a multiaffine polynomial, i.e. of degree 1 in each w_x separately. Combinatorially, $Z_W(\mathbf{w})$ is the generating polynomial for *induced subgraphs* of the complete graph, in which each vertex x gets weight w_x and each edge xy gets weight $W(x, y)$.

If, in addition to hard-core self-repulsion, we have $W(x, y) = 0$ or 1 for each pair $x \neq y$ — in statistical mechanics this is called a *hard-core pair interaction* — then we can define a (simple loopless) graph $G = (X, E)$ by setting $xy \in E$ whenever $W(x, y) = 0$ and $x \neq y$, so that $Z_W(\mathbf{w})$ is precisely the *independent-set polynomial* for G :

$$Z_G(\mathbf{w}) = \sum_{\substack{X' \subseteq X \\ X' \text{ independent}}} \prod_{x \in X'} w_x. \quad (1.3)$$

Traditionally the independent-set polynomial is defined as a univariate polynomial $Z_G(w)$ in which w_x is set equal to the same value w for all vertices x [4, 41, 42, 56, 51, 52, 53, 63, 27, 28, 64, 77]. But one of our main contentions in this paper is that Z_G is

more naturally understood as a multivariate polynomial; this allows us, in particular, to exploit the fact that Z_G is multiaffine.¹

More generally, given any W satisfying $0 \leq W(x, y) \leq 1$ for all x, y , let us define a simple loopless graph $G = G_W$ (the *support graph* of W) by setting $xy \in E(G)$ if and only if $W(x, y) \neq 1$ and $x \neq y$.² The partition function $Z_W(\mathbf{w})$ can be thought of as a “soft” version of the independent-set polynomial for G , in which an edge $xy \in E(G)$ has “strength” $1 - W(x, y) \in (0, 1]$.

In summary, we shall consider in this paper three levels of generality:

- (a) The general repulsive lattice gas (1.1), in which multiple occupation of a site may be permitted.
- (b) The lattice gas with hard-core self-repulsion (1.2), in which multiple occupation of a site is forbidden, but in which the interactions between adjacent sites may be “soft”.
- (c) The lattice gas with hard-core self-repulsion and hard-core pair interaction (1.3), which is simply the independent-set polynomial for the graph G .

Let us now return to the general case of a repulsive lattice gas (1.1). Since $Z_W(\mathbf{w})$ is an entire function of \mathbf{w} satisfying $Z_W(\mathbf{0}) = 1$, its logarithm is analytic in some neighborhood of $\mathbf{w} = \mathbf{0}$, and so can be expanded in a convergent Taylor series:

$$\log Z_W(\mathbf{w}) = \sum_{\mathbf{n}} c_{\mathbf{n}}(W) \mathbf{w}^{\mathbf{n}}, \quad (1.4)$$

where we have used the notation $\mathbf{w}^{\mathbf{n}} = \prod_{x \in X} w_x^{n_x}$, and of course $c_{\mathbf{0}} = 0$. In statistical mechanics, (1.4) is called the *Mayer expansion* [112], and there is a beautiful formula for the coefficients $c_{\mathbf{n}}(W)$, whose derivation we will review briefly (Proposition 2.1). As a corollary of this formula, we will show (Proposition 2.8) that the Taylor series (1.4) has alternating signs whenever the lattice gas is repulsive:

$$(-1)^{|\mathbf{n}|-1} c_{\mathbf{n}}(W) \geq 0 \quad (1.5)$$

for $0 \leq W \leq 1$. (Here $|\mathbf{n}| = \sum_{x \in X} n_x$.) Moreover, in this interval, $|c_{\mathbf{n}}(W)| = (-1)^{|\mathbf{n}|-1} c_{\mathbf{n}}(W)$ is a decreasing function of each $W(x, y)$, i.e. an increasing function of the “interaction strength” $1 - W(x, y)$.

¹More generally, let $G = (X, E)$ be a graph, and suppose that for $x \neq y$ we have $W(x, y) = 0$ or 1 according as $xy \in E$ or $xy \notin E$; but let $0 \leq W(x, x) \leq 1$ be arbitrary. Then we have the amusing identity $Z_W(\mathbf{w}) = Z_G(\tilde{\mathbf{w}})$ where

$$\tilde{w}_x = \sum_{n=1}^{\infty} \frac{W(x, x)^{n(n-1)/2}}{n!} w_x^n.$$

²Strictly speaking, G_W ought to be called the (simple) support graph of $1 - W$, since $xy \in E(G)$ if and only if $1 - W(x, y) \neq 0$ and $x \neq y$. In particular, in the case of a hard-core self-repulsion and hard-core pair interaction, W is the adjacency matrix of the complementary graph \bar{G} . Please note also that G_W does not contain any information about the diagonal weights $W(x, x)$.

As a simple consequence of the alternating-sign property for $\log Z_W$, we will then prove (Theorem 2.10) the equivalence of a number of conditions for the nonvanishing of Z_W in a closed polydisc $\bar{D}_{\mathbf{R}} = \{\mathbf{w}: |w_x| \leq R_x \text{ for all } x\}$. These equivalent conditions will play a central role in our study of the Lovász local lemma (Section 4).

Finally, we will provide (Section 5) some sufficient conditions for the nonvanishing of Z_W in a closed polydisc $\bar{D}_{\mathbf{R}}$, based on “local” properties of the interaction W (or of the graph G). Results of this type have traditionally been proven [88, 33, 97, 29, 30, 32, 100, 31, 104] by explicitly bounding the terms in the Mayer expansion (1.4); this requires some rather nontrivial combinatorics (for example, Proposition 2.4 below together with the counting of trees). Once this is done, an immediate consequence is that Z_W is nonvanishing in any polydisc where the series for $\log Z_W$ is convergent. Dobrushin’s brilliant idea [37, 38] was to prove these two results in the opposite order. First one proves, by an elementary induction on the cardinality of the state space, that Z_W is nonvanishing in a suitable polydisc (Theorem 5.1); it then follows immediately that $\log Z_W$ is analytic in that polydisc, and hence that its Taylor series (1.4) is convergent there. In Section 5 we will prove some refinements of this result and investigate their sharpness; and in Sections 6 and 7 we will provide some complementary results that give additional insight into the nature of these bounds. Let us remark that the Dobrushin–Shearer inductive method employed in Section 5 is limited, at present, to models with hard-core self-repulsion (1.2), for which Z_W is a multiaffine polynomial. It is an interesting open question to know whether this approach can be made to work without the assumption of hard-core self-repulsion.

1.2 The Lovász local lemma

In combinatorics we are frequently faced with a finite family $(A_x)_{x \in X}$ of “bad” events in some probability space, having probabilities $p_x = \mathbb{P}(A_x)$, and we want to show that there is a positive probability that none of the events A_x occurs. Under what conditions can we do this? In general all we can say is that $\mathbb{P}(\bigcap_{x \in X} \bar{A}_x) \geq 1 - \sum_x p_x$, since in the worst case the events A_x could be disjoint; so we would need $\sum_x p_x < 1$ to ensure that $\mathbb{P}(\bigcap_{x \in X} \bar{A}_x) > 0$. On the other hand, if the events $(A_x)_{x \in X}$ are independent, then $\mathbb{P}(\bigcap_{x \in X} \bar{A}_x) = \prod_x (1 - p_x)$, which is positive as soon as $p_x < 1$ for all x . This suggests that if the $(A_x)_{x \in X}$ are in some sense “not too strongly dependent”, then it might be possible to prove $\mathbb{P}(\bigcap_{x \in X} \bar{A}_x) > 0$ under relatively mild conditions on the $\{p_x\}$. This is the situation addressed by the Lovász local lemma: we allow strong dependence among *some* subsets of the $(A_x)_{x \in X}$, but insist that *most* of these events are independent. Specifically, the local lemma applies to collections of events in which the dependencies are controlled by a dependency graph, so that each event is independent from the events that are not adjacent to it.

More precisely, let G be a graph with vertex set X . We say that G is a *dependency graph* for the family $(A_x)_{x \in X}$ if, for each $x \in X$, the event A_x is independent from the σ -algebra generated by the events $\{A_y: y \in X \setminus \Gamma^*(x)\}$. [Here we have used the notation $\Gamma^*(x) = \Gamma(x) \cup \{x\}$, where $\Gamma(x)$ is the set of vertices of G adjacent to x .] Erdős and Lovász [39] proved the following fundamental result:

Theorem 1.1 (Lovász local lemma) *Let G be a dependency graph for the family of events $(A_x)_{x \in X}$, and suppose that $(r_x)_{x \in X}$ are real numbers in $[0, 1)$ such that, for each x ,*

$$\mathbb{P}(A_x) \leq r_x \prod_{y \in \Gamma(x)} (1 - r_y). \quad (1.6)$$

Then $\mathbb{P}(\bigcap_{x \in X} \bar{A}_x) \geq \prod_{x \in X} (1 - r_x) > 0$.

Erdős and Spencer [40] (see also [2, 83]) later noted that the same conclusion holds even if A_x and $\sigma(A_y: y \in X \setminus \Gamma^*(x))$ are not independent, provided that the “harmful” conditional probabilities are suitably bounded. More precisely:

Theorem 1.2 (Lopsided Lovász local lemma) *Let $(A_x)_{x \in X}$ be a family of events on some probability space, and let G be a graph with vertex set X . Suppose that $(r_x)_{x \in X}$ are real numbers in $[0, 1)$ such that, for each x and each $Y \subseteq X \setminus \Gamma^*(x)$, we have*

$$\mathbb{P}(A_x | \bigcap_{y \in Y} \bar{A}_y) \leq r_x \prod_{y \in \Gamma(x)} (1 - r_y). \quad (1.7)$$

Then $\mathbb{P}(\bigcap_{x \in X} \bar{A}_x) \geq \prod_{x \in X} (1 - r_x) > 0$.

In fact, the arguments of [39, 40] (see also [105, 106]) show that in Theorems 1.1 and 1.2 a slightly stronger conclusion holds: for all pairs Y, Z of subsets of X we have

$$\mathbb{P}\left(\bigcap_{x \in Y} \bar{A}_x \mid \bigcap_{x \in Z} \bar{A}_x\right) \geq \prod_{x \in Y \setminus Z} (1 - r_x). \quad (1.8)$$

In this paper we shall (following Shearer [98]) approach the problem by dividing our discussion into two parts:

- 1) A best-possible condition to have $\mathbb{P}(\bigcap_{x \in X} \bar{A}_x) > 0$, in terms of the independent-set polynomial $Z_G(-\mathbf{p})$; and
- 2) A *sufficient* condition to have $Z_G(-\mathbf{p}) > 0$, along the lines of Lovász, Dobrushin and Shearer.

We shall treat the first problem in Section 4 and the second problem in Sections 5–7. We shall also prove a generalization of the Lovász local lemma that allows for “soft” dependencies (Theorems 4.2 and 5.4).

Let us remark that we have been able to relate the Lovász local lemma to a combinatorial polynomial (namely, the independent-set polynomial) only in the case of an *undirected* dependency graph G . Although the local lemma can be formulated quite naturally for a dependency *digraph* [2, 23, 83], we do not know whether the digraph Lovász problem can be related to any combinatorial polynomial. (Clearly the independent-set polynomial cannot be the right object in the digraph context, since exclusion of simultaneous occupation is manifestly a symmetric condition.)

1.3 Plan of this paper

The plan of this paper is as follows: In Section 2 we prove some general properties of the partition function of a repulsive lattice gas. In Section 3 we study the additional properties that arise in the case of a hard-core self-repulsion (1.2), for which Z_W is a multiaffine polynomial. We also calculate some simple examples; in particular, we show that when the support graph $G = G_W$ is a tree, the partition function Z_W can easily be calculated by working upwards from the leaves. In Section 4 we prove our lower bound on $\mathbb{P}(\bigcap_{x \in X} \bar{A}_x) > 0$ in terms of the independent-set polynomial $Z_G(-\mathbf{p})$. We also prove a “soft-core” generalization of this result. In Section 5 we give some sufficient conditions for the nonvanishing of Z_W in a polydisc, and investigate their sharpness. In Section 6 we show how the bounds of Section 5 can be interpreted in terms of a lattice gas on either the “self-avoiding-walk tree” or the “pruned self-avoiding-walk tree” of G . In Section 7 we show how this tree bound can be understood as arising from the repeated application of a single “unfolding” step. Finally, in Section 8 we study the repulsive lattice gas on an infinite graph.

The reader primarily interested in the Lovász local lemma can read the statement of Theorem 2.10 (skipping the proof) and Definition 2.14, quickly read Section 3.1, and then jump directly to Section 4. The reader primarily interested in the lattice gas can skip Section 4.

2 The lattice-gas partition function

In this section we prove some general properties of the partition function of a repulsive lattice gas. We begin with some elementary identities (Section 2.1). In Section 2.2 we derive the Mayer expansion and prove some of its properties, notably the alternating-sign property (Proposition 2.8). In Section 2.3 we state and prove the “fundamental theorem” (Theorem 2.10), which sets forth a number of equivalent conditions defining the set that we shall call $\mathcal{R}(W)$ [Definition 2.14] and that plays a central role in the remainder of this paper. In Section 2.4 we derive the principal properties of the set $\mathcal{R}(W)$. In Section 2.5 we derive some further consequences of the alternating-sign property. In Section 2.6 (which is a digression from the main thread of the paper and can be omitted on a first reading) we discuss the algebraic irreducibility of the multivariate partition function $Z_W(\mathbf{w})$. Finally, in Section 2.7 we make a brief remark concerning the convexity of $\log Z_W(\mathbf{w})$ at nonnegative fugacity \mathbf{w} .

Nearly all of the results in this section are valid for an arbitrary repulsive lattice gas (1.1), in which multiple occupation of a site is permitted. A few of the results are restricted to the case of a hard-core self-repulsion (1.2), in which multiple occupation of a site is forbidden.

2.1 Elementary identities

Consider a general lattice gas (1.1) defined on a finite set X . For any subset $\Lambda \subseteq X$, let us define the restricted partition function

$$Z_{W,\Lambda}(\mathbf{w}) = Z_W(\mathbf{w} \mathbf{1}_\Lambda) \quad (2.1)$$

where

$$(\mathbf{w} \mathbf{1}_\Lambda)_x = \begin{cases} w_x & \text{if } x \in \Lambda \\ 0 & \text{otherwise} \end{cases} \quad (2.2)$$

This simply forces the sites in $X \setminus \Lambda$ to be unoccupied. Since W will be fixed throughout this subsection, we shall often omit it from the notation and write simply Z_Λ .

For any $x \in \Lambda$, we can expand the partition function (1.1b) in terms of the occupation number n_x : we obtain the *fundamental identity*

$$Z_\Lambda(\mathbf{w}) = \sum_{n=0}^{\infty} \frac{w_x^n}{n!} W(x, x)^{n(n-1)/2} Z_{\Lambda \setminus x}(W(x, \cdot)^n \mathbf{w}) \quad (2.3)$$

where

$$[W(x, \cdot)^n \mathbf{w}]_y = W(x, y)^n w_y. \quad (2.4)$$

In the special case of a hard-core self-repulsion at site x [i.e. $W(x, x) = 0$], only the terms $n = 0, 1$ appear in this sum; this case will play a key role in this paper starting in Section 3.1. (Analogous identities can be derived where we expand in the occupation numbers at an arbitrary set of sites $S \subseteq \Lambda$; for example, a two-site identity for the hard-core case will play a major role in Section 7.)

From (2.3) we can deduce the *differentiation identity*

$$\frac{\partial Z_\Lambda(\mathbf{w})}{\partial w_x} = \sum_{n=1}^{\infty} \frac{w_x^{n-1}}{(n-1)!} W(x, x)^{n(n-1)/2} Z_{\Lambda \setminus x}(W(x, \cdot)^n \mathbf{w}) \quad (2.5a)$$

$$= \sum_{n=0}^{\infty} \frac{w_x^n}{n!} W(x, x)^{n(n+1)/2} Z_{\Lambda \setminus x}(W(x, \cdot)^{n+1} \mathbf{w}) \quad (2.5b)$$

$$= \sum_{n=0}^{\infty} \frac{[W(x, x)w_x]^n}{n!} W(x, x)^{n(n-1)/2} Z_{\Lambda \setminus x}(W(x, \cdot)^n W(x, \cdot) \mathbf{w}) \quad (2.5c)$$

$$= Z_\Lambda(W(x, \cdot) \mathbf{w}). \quad (2.5d)$$

Repeated application of (2.5) yields the *multiple differentiation identity*

$$\frac{\partial^n Z_\Lambda(\mathbf{w})}{\partial w_{x_1} \cdots \partial w_{x_n}} = \left(\prod_{1 \leq i < j \leq n} W(x_i, x_j) \right) Z_\Lambda \left(\prod_{i=1}^n W(x_i, \cdot) \mathbf{w} \right), \quad (2.6)$$

where $\left(\prod_{i=1}^n W(x_i, \cdot) \mathbf{w} \right)_x = w_x \prod_{i=1}^n W(x_i, x)$.

2.2 The Mayer expansion

We begin by reviewing the derivation of the Mayer expansion (1.4). The first step is to trivially rewrite the partition function (1.1a) as

$$Z_W(\mathbf{w}) = \sum_{n=0}^{\infty} \frac{1}{n!} \sum_{x_1, \dots, x_n \in X} \left(\prod_{i=1}^n w_{x_i} \right) \sum_{G \in \mathcal{G}_n} \prod_{ij \in E(G)} F(x_i, x_j), \quad (2.7)$$

where \mathcal{G}_n is the set of all (simple loopless undirected) graphs on the vertex set $\{1, \dots, n\}$, and

$$F(x, y) = W(x, y) - 1 \quad (2.8)$$

is called the *two-particle Mayer factor*. This is of the form

$$Z_W(\mathbf{w}) = \sum_{n=0}^{\infty} \frac{1}{n!} \sum_{G \in \mathcal{G}_n} \mathcal{W}(G) \quad (2.9)$$

where the weights

$$\mathcal{W}(G) = \sum_{x_1, \dots, x_n \in X} \left(\prod_{i=1}^n w_{x_i} \right) \prod_{ij \in E(G)} F(x_i, x_j) \quad (2.10)$$

satisfy

- (a) $\mathcal{W}(\emptyset) = 1$;
- (b) $\mathcal{W}(G) = \mathcal{W}(G')$ whenever $G \cong G'$ (i.e. whenever G and G' differ only by a relabelling of vertices); and
- (c) $\mathcal{W}(G) = \mathcal{W}(G_1) \mathcal{W}(G_2)$ whenever G is isomorphic to the disjoint union of G_1 and G_2 .

It then follows from the exponential formula [112, 115, 13, 108] that

$$\log Z_W(\mathbf{w}) = \sum_{n=0}^{\infty} \frac{1}{n!} \sum_{G \in \mathcal{C}_n} \mathcal{W}(G) \quad (2.11a)$$

$$= \sum_{n=0}^{\infty} \frac{1}{n!} \sum_{x_1, \dots, x_n \in X} \left(\prod_{i=1}^n w_{x_i} \right) \sum_{G \in \mathcal{C}_n} \prod_{ij \in E(G)} F(x_i, x_j), \quad (2.11b)$$

at least in the sense of formal power series in \mathbf{w} , where $\mathcal{C}_n \subseteq \mathcal{G}_n$ is the set of *connected* graphs on $\{1, \dots, n\}$. Therefore:

Proposition 2.1 *The coefficients $c_{\mathbf{n}}(W)$ of the Mayer expansion (1.4) are given by*

$$c_{\mathbf{n}}(W) = \frac{1}{n!} \sum_{\substack{x_1, \dots, x_n \in X \\ \#\{i: x_i = x\} = n_x \forall x}} \sum_{G \in \mathcal{C}_n} \prod_{ij \in E(G)} F(x_i, x_j) \quad (2.12a)$$

$$= \sum_{\substack{(S_x)_{x \in X} \\ \bigsqcup_{x \in X} S_x = \{1, \dots, n\} \\ |S_x| = n_x \forall x}} \sum_{G \in \mathcal{C}_n} \prod_{\{x, y\} \subseteq X} F(x, y)^{e_G(S_x, S_y)} \prod_{x \in X} F(x, x)^{e_G(S_x)} \quad (2.12b)$$

where $n = |\mathbf{n}|$, the first sum in (2.12b) runs over all partitions of $\{1, \dots, n\}$ into disjoint subsets $(S_x)_{x \in X}$ with the specified cardinalities, and $e_G(S_x, S_y)$ [resp. $e_G(S_x)$] denotes the number of edges of G connecting S_x to S_y [resp. within S_x].

Remark. Even in the simple case $X = \{x\}$ and $W(x, x) = 0$, for which $\log Z_W(w) = \log(1 + w) = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} w^n$, the Mayer expansion (2.11) implies the nontrivial identity

$$\sum_{G \in \mathcal{C}_n} (-1)^{|E(G)|} = (-1)^{n-1} (n-1)! \quad (2.13)$$

for the generating function of connected spanning subgraphs of the complete graph K_n . We leave it as an exercise for the reader to find a direct combinatorial proof of (2.13).

In order to analyze the Mayer coefficients $c_{\mathbf{n}}(W)$, it is convenient to proceed in a bit more generality. So let $H = (V, E)$ be a graph — we allow loops and multiple edges — and let $\mathbf{z} = \{z_e\}_{e \in E}$ be a family of complex edge weights for H . [Later we will specialize to $H = K_n$.] Define the generating function of connected spanning subgraphs of H (or “connected sum” for short),

$$C_H(\mathbf{z}) = \sum_{\substack{E' \subseteq E \\ (V, E') \text{ connected}}} \prod_{e \in E'} z_e. \quad (2.14)$$

It is easy to see that C_H satisfies the deletion-contraction relation

$$C_H(\mathbf{z}) = C_{H \setminus e}(\mathbf{z}_{\neq e}) + z_e C_{H/e}(\mathbf{z}_{\neq e}) \quad (2.15)$$

for any edge $e \in E$: here $H \setminus e$ is the graph H with edge e deleted, H/e is the graph H with edge e contracted (note that we do *not* delete any loops or multiple edges that may be formed), and $\mathbf{z}_{\neq e}$ denotes the family $\{z_f\}_{f \in E \setminus e}$. Please note that if e is a loop, then $H/e = H \setminus e$ and hence

$$C_H(\mathbf{z}) = C_{H \setminus e}(\mathbf{z}_{\neq e}) + z_e C_{H/e}(\mathbf{z}_{\neq e}) = (1 + z_e) C_{H \setminus e}(\mathbf{z}_{\neq e}). \quad (2.16)$$

Let \mathcal{C} (resp. \mathcal{T}) be the set of subsets $E' \subseteq E$ such that (V, E') is connected (resp. is a tree). Clearly \mathcal{C} is an increasing family of subsets of E with respect to set-theoretic inclusion, and the minimal elements of \mathcal{C} are precisely those of \mathcal{T} (i.e. the

spanning trees). It is a nontrivial but well-known fact [17, Sections 7.2 and 7.3] [117, Section 8.3] that the (anti-)complex \mathcal{C} is *partitionable*: that is, there exists a map $R: \mathcal{T} \rightarrow \mathcal{C}$ such that $R(T) \supseteq T$ for all $T \in \mathcal{T}$ and $\mathcal{C} = \bigsqcup_{T \in \mathcal{T}} [T, R(T)]$ (disjoint union), where $[E_1, E_2]$ denotes the Boolean interval $\{E': E_1 \subseteq E' \subseteq E_2\}$. In fact, many alternative choices of R are available [17, Sections 7.2 and 7.3] [45, Sections 2 and 6] [15, Proposition 13.7 et seq.] [101, Proposition 4.1], and most of our arguments will not depend on any specific choice of R . However, at one point (Proposition 2.6) we shall need the existence of an R with the following additional property:

Lemma 2.2 *Let $H = (V, E)$ be a connected graph, and fix any $T_0 \in \mathcal{T}$. Then there exists a map $R: \mathcal{T} \rightarrow \mathcal{C}$ such that*

- (a) $R(T) \supseteq T$ for all $T \in \mathcal{T}$;
- (b) \mathcal{C} is the disjoint union of the Boolean intervals $[T, R(T)]$, $T \in \mathcal{T}$; and
- (c) $R(T_0) = T_0$.

PROOF. If H has one vertex and no edges, then $\mathcal{T} = \mathcal{C} = \{\emptyset\}$ and the result holds trivially; so assume henceforth that $E \neq \emptyset$. Assign arbitrary weights $w_e > 0$ chosen so that no two spanning trees have equal weight (for example, one can choose the w_e to be linearly independent over the rationals). For each $E' \in \mathcal{C}$, let $S(E')$ be the (unique) minimum-weight spanning tree contained in E' . (This can be constructed by a greedy algorithm, i.e. start from \emptyset and keep adding the lowest-weight edge in E' that does not create a cycle. See, for instance, [22, Section I.2].) We then define $R(T)$ to be the union of all E' that have $S(E') = T$. To verify that this works, we need to show that if $S(E_1) = S(E_2) = T$, then $S(E_1 \cup E_2) = T$; but this follows easily from the validity of the greedy algorithm. [Note that this construction includes, as a special case, the lexicographically minimum spanning tree for any ordering of E : it suffices to take $w_n = 2^n$.]

By choosing the weights so that $w_e > w_f$ whenever $e \in T_0$ and $f \in E \setminus T_0$, we can ensure that for $E' \neq T_0$ the first edge chosen by the greedy algorithm will not belong to T_0 . Therefore, $R(T_0) = T_0$. ■

Remark. The greedy algorithm works, more generally, for an arbitrary matroid [86, Section 1.8]. It follows that the independent-set complex of a matroid (or, dually, the spanning-set anti-complex) is partitionable [17, Sections 7.2 and 7.3].

Given the existence of R , we have the following simple but fundamental identity:

Proposition 2.3 (partitionability identity) *Let $R: \mathcal{T} \rightarrow \mathcal{C}$ be any map satisfying $R(T) \supseteq T$ for all $T \in \mathcal{T}$ and \mathcal{C} is the disjoint union of the Boolean intervals $[T, R(T)]$, for $T \in \mathcal{T}$. Then*

$$C_H(\mathbf{z}) = \sum_{\substack{T \subseteq E \\ (V, T) \text{ tree}}} \prod_{e \in T} z_e \sum_{T \subseteq E' \subseteq R(T)} \prod_{e \in E' \setminus T} z_e$$

$$= \sum_{\substack{T \subseteq E \\ (V, T) \text{ tree}}} \prod_{e \in T} z_e \prod_{e \in R(T) \setminus T} (1 + z_e). \quad (2.17)$$

This identity (for one specific choice of R) is due originally to Penrose [88].

One immediate consequence of the identity (2.17) is the following inequality valid in the “complex repulsive” regime $|1 + z_e| \leq 1$:

Proposition 2.4 (Penrose [88]) *Let $H = (V, E)$ be a finite undirected graph equipped with complex edge weights $\{z_e\}_{e \in E}$ satisfying $|1 + z_e| \leq 1$ for all e . Then*

$$|C_H(\mathbf{z})| \leq \sum_{\substack{T \subseteq E \\ (V, T) \text{ tree}}} \prod_{e \in T} |z_e|. \quad (2.18)$$

This bound plays a major role in several traditional (pre-Dobrushin) proofs of convergence of the Mayer expansion [88, 33, 97, 29, 100, 31, 104], as well as in a recent bound on the zeros of chromatic polynomials [101].

Let us now define the “generalized connected sum”

$$C_H(\lambda; \mathbf{z}) = \sum_{\substack{E' \subseteq E \\ (V, E') \text{ connected}}} \lambda^{c(E')} \prod_{e \in E'} z_e \quad (2.19a)$$

$$= \lambda^{-(|V|-1)} C_H(\lambda \mathbf{z}) \quad (2.19b)$$

where $c(E') = |E'| - |V| + 1$ is the cyclomatic number of the connected subgraph (V, E') [more generally, for a subgraph (V, E') with $k(E')$ components, the cyclomatic number is $c(E') = |E'| - |V| + k(E')$]. Of course, (2.19b) shows that $C_H(\lambda; \mathbf{z})$ contains no more information than $C_H(\mathbf{z})$; it is just a convenient way of scaling all the variables z_e simultaneously. The function $C_H(\lambda; \mathbf{z})$ interpolates between the tree sum ($\lambda = 0$) and the connected sum ($\lambda = 1$); and Proposition 2.4 can be rephrased as saying that $|C_H(1; \mathbf{z})| \leq C_H(0; |\mathbf{z}|)$. One of us has conjectured that the absolute-value signs can in fact be put outside the sum, i.e. $|C_H(1; \mathbf{z})| \leq |C_H(0; \mathbf{z})|$ [101, Remark 2 in Section 4.1]. In general this is still an open problem; but if the z_e are *real* and lie in the interval $[-1, 0]$ — which corresponds to the physical repulsive regime $0 \leq W \leq 1$ — then a vastly *stronger* result is true:

Proposition 2.5 (Sokal [101]) *Let $H = (V, E)$ be a finite undirected graph equipped with real edge weights $\{z_e\}_{e \in E}$ satisfying $-1 \leq z_e \leq 0$ for all e . Then*

$$(-1)^{k+|V|-1} \frac{d^k}{d\lambda^k} C_H(\lambda; \mathbf{z}) \geq 0 \quad (2.20)$$

on $0 \leq \lambda \leq 1$, for all integers $k \geq 0$. In particular, setting $k = 0$ and $\lambda = 1$ we have

$$(-1)^{|V|-1} C_H(\mathbf{z}) \geq 0. \quad (2.21)$$

PROOF. By (2.17) and (2.19b), we have

$$C_H(\lambda; \mathbf{z}) = \sum_{\substack{T \subseteq E \\ (V, T) \text{ tree}}} \prod_{e \in T} z_e \prod_{e \in \mathbf{R}(T) \setminus T} (1 + \lambda z_e) \quad (2.22)$$

and hence

$$\frac{d^k}{d\lambda^k} C_H(\lambda; \mathbf{z}) = k! \sum_{\substack{T \subseteq E \\ (V, T) \text{ tree}}} \sum_{\substack{\tilde{T} \subseteq \mathbf{R}(T) \setminus T \\ |\tilde{T}| = k}} \prod_{e \in T \cup \tilde{T}} z_e \prod_{e \in \mathbf{R}(T) \setminus (T \cup \tilde{T})} (1 + \lambda z_e), \quad (2.23)$$

which has the claimed sign whenever $0 \leq \lambda \leq 1$ and $-1 \leq z_e \leq 0$ for all e .³ ■

We shall also need a variant of Proposition 2.5 with strict inequality for the case $k = 0$:

Proposition 2.6 *Let $H = (V, E)$ be a finite undirected graph equipped with real edge weights $\{z_e\}_{e \in E}$ satisfying $-1 \leq z_e \leq 0$ for all e . Assume furthermore that the subgraph $(V, \text{supp } \mathbf{z})$ is connected [here $\text{supp } \mathbf{z} = \{e \in E: z_e \neq 0\}$]. Then*

$$(-1)^{|V|-1} C_H(\mathbf{z}) > 0. \quad (2.24)$$

PROOF. Choose $T_0 \subseteq \text{supp } \mathbf{z}$ such that (V, T_0) is a spanning tree, and use Lemma 2.2 to choose \mathbf{R} so that $\mathbf{R}(T_0) = T_0$. Then, in the sum (2.22), the term $T = T_0$ is nonzero with sign $(-1)^{|V|-1}$, and all the other terms either have the sign $(-1)^{|V|-1}$ or else are zero. ■

We can also prove some inequalities on the partial derivatives of $C_H(\mathbf{z})$ with respect to individual weights z_e . Note first that the deletion-contraction identity (2.15) implies that

$$\frac{\partial C_H(\mathbf{z})}{\partial z_e} = C_{H/e}(\mathbf{z}_{\neq e}). \quad (2.25)$$

The graph H/e has $|V|$ vertices if e is a loop, and $|V| - 1$ vertices if e is not a loop. Repeated application of these facts together with (2.21) yields the following result:

Proposition 2.7 *Let $H = (V, E)$ be a finite undirected graph equipped with real edge weights $\{z_e\}_{e \in E}$ satisfying $-1 \leq z_e \leq 0$ for all e . Let $k \geq 1$ and $e_1, \dots, e_k \in E$.*

(a) *If e_1, \dots, e_k are not all distinct, we have*

$$\frac{\partial C_H(\mathbf{z})}{\partial z_{e_1} \cdots \partial z_{e_k}} = 0. \quad (2.26)$$

³Equation (4.7) of [101] inadvertently omitted the prefactor $k!$ in (2.23).

(b) If e_1, \dots, e_k are all distinct and form a subgraph with cyclomatic number c , we have

$$(-1)^{|V|-k+c-1} \frac{\partial C_H(\mathbf{z})}{\partial z_{e_1} \cdots \partial z_{e_k}} \geq 0. \quad (2.27)$$

Let us now specialize these results to the Mayer expansion (2.12a) by taking $H = K_n$ (where $n = |\mathbf{n}|$ and K_n denotes the complete graph on n vertices), $z_{ij} = F(x_i, x_j) = W(x_i, x_j) - 1$ and then summing over $x_1, \dots, x_n \in X$ with the specified cardinalities: we get

$$c_{\mathbf{n}}(W) = \frac{1}{n!} \sum_{\substack{x_1, \dots, x_n \in X \\ \#\{i: x_i = x\} = n_x \forall x}} C_{K_n}(\mathbf{z}(\mathbf{x})) \quad (2.28)$$

where $\mathbf{x} = (x_1, \dots, x_n)$ and $\mathbf{z}(\mathbf{x})_{ij} = F(x_i, x_j)$. Since any subgraph of K_n with 0, 1 or 2 edges has cyclomatic number 0, we can use Propositions 2.5 and 2.7 to deduce the following:

Proposition 2.8 (signs of Mayer coefficients) *Suppose that the lattice gas is repulsive, i.e. $0 \leq W(x, y) \leq 1$ for all $x, y \in X$. Then, for all $x, y, x', y' \in X$, the Mayer coefficients $c_{\mathbf{n}}(W)$ satisfy*

$$(-1)^{|\mathbf{n}|-1} c_{\mathbf{n}}(W) \geq 0 \quad (2.29)$$

$$(-1)^{|\mathbf{n}|-1} \frac{\partial c_{\mathbf{n}}(W)}{\partial W(x, y)} \leq 0 \quad (2.30)$$

$$(-1)^{|\mathbf{n}|-1} \frac{\partial^2 c_{\mathbf{n}}(W)}{\partial W(x, y) \partial W(x', y')} \geq 0 \quad (2.31)$$

PROOF. (2.29) is an immediate consequence of (2.28) and (2.21). For (2.30), note first that for any fixed $x_1, \dots, x_n \in X$, we have

$$\frac{\partial C_{K_n}(\mathbf{z}(\mathbf{x}))}{\partial W(x, y)} = \sum_{\{i, j\} \in E(x, y; x_1, \dots, x_n)} \left. \frac{\partial C_{K_n}(\mathbf{z})}{\partial z_{ij}} \right|_{\mathbf{z}=\mathbf{z}(\mathbf{x})}, \quad (2.32)$$

where $E(x, y; x_1, \dots, x_n)$ is the set of unordered pairs $\{i, j\}$ ($i \neq j$) such that $\{x_i, x_j\} = \{x, y\}$. Since any subgraph of K_n with one edge has cyclomatic number 0, the inequality (2.30) now follows from (2.27). Similarly, the left-hand side of (2.31) gives rise to a double sum over pairs $\{i, j\} \in E(x, y; x_1, \dots, x_n)$ and $\{i', j'\} \in E(x', y'; x_1, \dots, x_n)$. The diagonal terms (if any) in this sum vanish by (2.26), and the other terms have the claimed sign by (2.27). ■

Please note that (2.31) can be nonzero even when $x = x'$ and $y = y'$, because distinct edges of K_n could correspond to the same pair x, y . Note also that the sign of the third and higher derivatives cannot be controlled by this method, because a

subgraph of K_n ($n \geq 4$) containing three or more edges could have cyclomatic number of either parity. Indeed, for the simple case $X = \{x\}$ and $W(x, x) = W$ we have

$$Z_W(w) = \sum_{n=0}^{\infty} \frac{w^n W^{n(n-1)/2}}{n!} \quad (2.33)$$

and hence

$$\begin{aligned} \log Z_W(w) &= w + \frac{F}{2}w^2 + \frac{3F^2 + F^3}{6}w^3 \\ &\quad + \frac{16F^3 + 15F^4 + 6F^5 + F^6}{24}w^4 + O(w^5) \end{aligned} \quad (2.34a)$$

$$\begin{aligned} &= w + \frac{W-1}{2}w^2 + \frac{(W+2)(W-1)^2}{6}w^3 \\ &\quad + \frac{(W^3 + 3W^2 + 6W + 6)(W-1)^3}{24}w^4 + O(w^5), \end{aligned} \quad (2.34b)$$

so that

$$\frac{\partial^3 c_4(W)}{\partial W^3} = 5W^3 - 1 \quad (2.35)$$

has no fixed sign on $0 \leq W \leq 1$.

We can also determine the precise conditions under which the inequality (2.29) is strict:

Proposition 2.9 (condition for Mayer coefficients to be nonzero) *Suppose that the lattice gas is repulsive, i.e. $0 \leq W(x, y) \leq 1$ for all $x, y \in X$. Let $\mathbf{n} = (n_x)_{x \in X}$ be a multi-index with support $\text{supp } \mathbf{n} = \{x \in X: n_x \neq 0\}$. Then $(-1)^{|\mathbf{n}|-1} c_{\mathbf{n}}(W) > 0$ if and only if one of the following is true:*

- (a) $|\mathbf{n}| = 1$;
- (b) $|\mathbf{n}| \geq 2$, $\text{supp } \mathbf{n} = \{x\}$ and $W(x, x) \neq 1$; or
- (c) $|\text{supp } \mathbf{n}| \geq 2$ and the induced subgraph $G_W[\text{supp } \mathbf{n}]$ is connected.

PROOF. In case (a), $c_{\mathbf{n}}(W) = 1 > 0$; in case (b) or (c), for each term in (2.28) [actually they are all the same!] the subgraph of K_n with edge set $\text{supp } \mathbf{z}(\mathbf{x})$ is connected, so that $(-1)^{|\mathbf{n}|-1} C_{K_n}(\mathbf{z}(\mathbf{x})) > 0$ by Proposition 2.6.

There are only two other possibilities:

- (d) $\mathbf{n} = 0$;
- (e) $|\text{supp } \mathbf{n}| \geq 2$ and $G_W[\text{supp } \mathbf{n}]$ is disconnected.

In case (d), clearly $c_{\mathbf{n}}(W) = 0$; and in case (e), for each term in (2.28) the subgraph of K_n with edge set $\text{supp } \mathbf{z}(x)$ is disconnected, so that $(-1)^{|\mathbf{n}|-1} C_{K_n}(\mathbf{z}(x)) = 0$. ■

Historical remark. The alternating-sign property (2.29) for the Mayer coefficients of a repulsive gas has been known in the physics literature for over 40 years: see Groeneveld [48] for a brief sketch of the proof, which uses methods quite different from ours.⁴ Nevertheless, this result does not seem to be as well known as it should be. To the best of our knowledge, the inequalities (2.30) and (2.31) are new. We think that the Mayer coefficients $c_{\mathbf{n}}(W)$, and more generally the “connected sums” $C_H(\mathbf{z})$, merit further study from a combinatorial point of view; we would not be surprised if new identities or inequalities were waiting to be discovered.

2.3 The fundamental theorem

Let us now state the principal result of this section:

Theorem 2.10 (The fundamental theorem) *Consider any repulsive lattice gas, and let $\mathbf{R} = \{R_x\}_{x \in X} \geq 0$. Then the following are equivalent:*

- (a) *There exists a connected set $C \subseteq (-\infty, 0]^X$ that contains both $\mathbf{0}$ and $-\mathbf{R}$, such that $Z_W(\mathbf{w}) > 0$ for all $\mathbf{w} \in C$. [Equivalently, $-\mathbf{R}$ belongs to the connected component of $Z_W^{-1}(0, \infty) \cap (-\infty, 0]^X$ containing $\mathbf{0}$.]*
- (b) *$Z_W(\mathbf{w}) > 0$ for all \mathbf{w} satisfying $-\mathbf{R} \leq \mathbf{w} \leq \mathbf{0}$.*
- (c) *$Z_W(\mathbf{w}) \neq 0$ for all \mathbf{w} satisfying $|\mathbf{w}| \leq \mathbf{R}$.⁵*
- (d) *The Taylor series for $\log Z_W(\mathbf{w})$ around $\mathbf{0}$ is convergent at $\mathbf{w} = -\mathbf{R}$.*
- (e) *The Taylor series for $\log Z_W(\mathbf{w})$ around $\mathbf{0}$ is absolutely convergent for $|\mathbf{w}| \leq \mathbf{R}$.*

Moreover, when these conditions hold, we have $|Z_W(\mathbf{w})| \geq Z_W(-\mathbf{R}) > 0$ for all \mathbf{w} satisfying $|\mathbf{w}| \leq \mathbf{R}$.

In the case of hard-core self-repulsion, (a)–(e) are also equivalent to

- (b') *$Z_W(-\mathbf{R} \mathbf{1}_S) > 0$ for all $S \subseteq X$, where*

$$(\mathbf{R} \mathbf{1}_S)_x = \begin{cases} R_x & \text{if } x \in S \\ 0 & \text{otherwise} \end{cases} \quad (2.36)$$

⁴Groeneveld [48] writes that “a more detailed account of this work, containing also the corresponding results for the pressure and the distribution functions will be published in *Physica*”, but to our knowledge that more detailed paper never appeared. See also Penrose [87, eq. (8.1)] for a related result.

⁵Here we use the notation $|\mathbf{w}| = \{|w_x|\}_{x \in X}$. This conflicts slightly with our notation $|\mathbf{n}| = \sum_{x \in X} n_x$ for multi-indices, but we trust that it will not lead to any confusion.

(f) $Z_W(-\mathbf{R}) > 0$, and $(-1)^{|S|} Z_W(-\mathbf{R}; S) \geq 0$ for all $S \subseteq X$, where

$$Z_W(\mathbf{w}; S) = \sum_{S \subseteq X' \subseteq X} \left(\prod_{x \in X'} w_x \right) \left(\prod_{\{x,y\} \subseteq X'} W(x,y) \right). \quad (2.37)$$

(g) There exists a probability measure P on 2^X satisfying $P(\emptyset) > 0$ and

$$\sum_{T \supseteq S} P(T) = \left(\prod_{x \in S} R_x \right) \left(\prod_{\{x,y\} \subseteq S} W(x,y) \right) \quad (2.38)$$

for all $S \subseteq X$. [This probability measure is unique and is given by $P(S) = (-1)^{|S|} Z_W(-\mathbf{R}; S)$. In particular, $P(\emptyset) = Z_W(-\mathbf{R}) > 0$.]

Remarks. 1. The conditions (b'), (f) and (g) are inspired in part by Shearer [98, Theorem 1].

2. Suppose that the univariate entire function $Z_W(w)$, defined by setting $w_x = w$ for all x , is strictly positive whenever $-R \leq w \leq 0$. Then in fact $Z_W(\mathbf{w}) > 0$ whenever $-R \leq w_x \leq 0$ for all x : this follows from (a) \implies (b) by taking C to be the segment $[-R, 0]$ of the diagonal.

The proof of Theorem 2.10 will hinge on the alternating-sign property (2.29) for the Taylor coefficients of $\log Z_W$. In preparation for this proof, let us recall the Vivanti–Pringsheim theorem in the theory of analytic functions of a single complex variable [62, Theorem 5.7.1]: if a power series $f(z) = \sum_{n=0}^{\infty} a_n z^n$ with *nonnegative* coefficients has a finite nonzero radius of convergence, then the point of the circle of convergence lying on the positive real axis is a singular point of the function f . Otherwise put, if f is a function whose Taylor series at 0 has all nonnegative coefficients and f is analytic on some complex neighborhood of the real interval $[0, R)$, then f is in fact analytic on the open disc of radius R centered at the origin and its Taylor series is absolutely convergent there. Here we will need the following multidimensional generalization of the Vivanti–Pringsheim theorem:

Proposition 2.11 (multidimensional Vivanti–Pringsheim theorem) *Let C be a connected subset of $[0, \infty)^n$ containing $\mathbf{0}$, let U be an open neighborhood of C in \mathbb{C}^n , and let f be a function analytic on U whose Taylor series around $\mathbf{0}$ has all nonnegative coefficients. Then the Taylor series of f around $\mathbf{0}$ converges absolutely on the set $\text{hull}(C) \equiv \bigcup_{\mathbf{R} \in C} \bar{D}_{\mathbf{R}}$, where $\bar{D}_{\mathbf{R}}$ denotes the closed polydisc $\{\mathbf{w} \in \mathbb{C}^n: |w_i| \leq R_i \text{ for all } i\}$, and it defines a function that is continuous on $\text{hull}(C)$ and analytic on its interior.*

Remark. We could equally well start from an open neighborhood U of $\mathbf{0}$ in \mathbb{C}^n , and define C to be the connected component of $U \cap [0, \infty)^n$ containing $\mathbf{0}$. This is the

maximal set C compatible with the given U ; note that it is open in $[0, \infty)^n$, and that $\text{hull}(C)$ is then open in \mathbb{C}^n .

PROOF. Let $f(\mathbf{z}) = \sum_{\mathbf{n}} a_{\mathbf{n}} \mathbf{z}^{\mathbf{n}}$ be the Taylor series of f around $\mathbf{0}$. And let us define

$$S = \{\mathbf{R} \in [0, \infty)^n: \sum_{\mathbf{n}} a_{\mathbf{n}} \mathbf{z}^{\mathbf{n}} \text{ converges absolutely at } \mathbf{z} = \mathbf{R}\} \quad (2.39a)$$

$$= \{\mathbf{R} \in [0, \infty)^n: \sum_{\mathbf{n}} a_{\mathbf{n}} \mathbf{z}^{\mathbf{n}} \text{ converges absolutely for } \mathbf{z} \in \bar{D}_{\mathbf{R}}\} \quad (2.39b)$$

Note that S is a down-set (that is, $\mathbf{R} \in S$ and $\mathbf{0} \leq \mathbf{R}' \leq \mathbf{R}$ imply $\mathbf{R}' \in S$) and that $\mathbf{0} \in S$; we shall show that $C \subset S$.

Consider any point $\mathbf{z}_0 \in \bar{S} \cap C$ (here \bar{S} denotes the closure of S). Choose $\epsilon > 0$ such that the closed polydisc of radius ϵ in each direction around \mathbf{z}_0 — call it $\bar{D}(\mathbf{z}_0, \epsilon)$ — is contained in U . Then choose $\mathbf{z}_1 \in S \cap \bar{D}(\mathbf{z}_0, \epsilon/5)$, and choose $\mathbf{z}_2 \in \bar{D}(\mathbf{z}_1, \epsilon/5)$ such that $\mathbf{0} \leq \mathbf{z}_2 \leq \mathbf{z}_1$ and $(\mathbf{z}_2)_i < (\mathbf{z}_1)_i$ for all coordinates i with $(\mathbf{z}_1)_i > 0$. It follows that $\bar{D}(\mathbf{z}_2, 3\epsilon/5) \subset U$. Now, since $\mathbf{z}_1 \in S$ and \mathbf{z}_2 is strictly below \mathbf{z}_1 in all nonzero coordinates, the Taylor series around $\mathbf{0}$ for f and all its derivatives converge absolutely at \mathbf{z}_2 . And the Taylor series for f around \mathbf{z}_2 converges absolutely in $\bar{D}(\mathbf{z}_2, 3\epsilon/5)$. If $\mathbf{z}_3 \in \bar{D}(\mathbf{z}_2, 3\epsilon/5)$ with $\mathbf{z}_3 - \mathbf{z}_2 \geq \mathbf{0}$, we can write

$$f(\mathbf{z}_3) = \sum_{\mathbf{k}} \frac{f^{(\mathbf{k})}(\mathbf{z}_2)}{\mathbf{k}!} (\mathbf{z}_3 - \mathbf{z}_2)^{\mathbf{k}} \quad (2.40a)$$

$$= \sum_{\mathbf{k}} (\mathbf{z}_3 - \mathbf{z}_2)^{\mathbf{k}} \sum_{\mathbf{n} \geq \mathbf{k}} \binom{\mathbf{n}}{\mathbf{k}} a_{\mathbf{n}} \mathbf{z}_2^{\mathbf{n}-\mathbf{k}} \quad (2.40b)$$

$$= \sum_{\mathbf{n}} a_{\mathbf{n}} \sum_{\mathbf{0} \leq \mathbf{k} \leq \mathbf{n}} \binom{\mathbf{n}}{\mathbf{k}} (\mathbf{z}_3 - \mathbf{z}_2)^{\mathbf{k}} \mathbf{z}_2^{\mathbf{n}-\mathbf{k}} \quad (2.40c)$$

$$= \sum_{\mathbf{n}} a_{\mathbf{n}} \mathbf{z}_3^{\mathbf{n}}, \quad (2.40d)$$

where the rearrangements are justified because all the terms are nonnegative, so that the convergence of the iterated sum (2.40b) implies the absolute convergence of the corresponding double sum. It follows that $\mathbf{z}_3 \in S$. In particular we can choose $\mathbf{z}_3 = \mathbf{z}_2 + (3\epsilon/5, \dots, 3\epsilon/5)$. Since S is a down-set, and since the real points of the polydisc $\bar{D}(\mathbf{z}_0, \epsilon/5)$ all lie below \mathbf{z}_3 , we conclude that $\bar{D}(\mathbf{z}_0, \epsilon/5) \cap [0, \infty)^n \subset S$. Hence $\bar{D}(\mathbf{z}_0, \epsilon/5) \cap C \subset S \cap C$. Since \mathbf{z}_0 is an arbitrary point of $\bar{S} \cap C$, it follows that $S \cap C$ is both closed and open in C (and nonempty since $\mathbf{0} \in S \cap C$). Since C is connected, we conclude that $S \cap C = C$. ■

For the proof of Theorem 2.10, we need an elementary topological lemma:

Lemma 2.12 *Let K be a convex set in \mathbb{R}^n , let C be an open connected subset of K (in the relative topology), let U be an open neighborhood of C in \mathbb{R}^n , and let $x_1, x_2 \in C$. Then there exists a finite polygonal path $P \subset C$ running from x_1 to x_2 and a simply connected open set U' in \mathbb{R}^n satisfying $P \subset U' \subset U$.*

PROOF. It is a well-known fact that any two points in C can be connected by a finite polygonal path P lying in C . [Sketch of proof: Define an equivalence relation \sim on C by setting $x \sim y$ iff there exists a finite polygonal path in C connecting x to y . Because K is convex and C is open in K , the equivalence classes of \sim are (relatively) open subsets of C . Since C is connected, there is just one equivalence class.] By removing loops, we can assume that P is non-self-intersecting.

Then we can take U' to be a sufficiently small tube centered on P . [Let δ be the minimal distance between non-adjacent segments of P , and let δ' be the distance from P to the complement of U ; we have $\delta' > 0$ by compactness of P . Then let U' be the set of all points whose distance from P is less than $\min(\delta/3, \delta'/2)$. It is not hard to see that U' is simply connected.⁶] ■

We shall also make use of the following elementary result:

Lemma 2.13 *Let F be a function on 2^X , and define*

$$F_-(S) = \sum_{X' \subseteq S} F(X') \quad (2.41)$$

$$F_+(S) = \sum_{X' \supseteq S} F(X') \quad (2.42)$$

Then

$$F_-(S) = \sum_{Y \subseteq S^c} (-1)^{|Y|} F_+(Y) \quad (2.43)$$

where $S^c \equiv X \setminus S$.

PROOF. This is a straightforward application of inclusion-exclusion (see e.g. [107, Section 2.1] or [1, Chapter IV]). We have

$$F(X') = \sum_{Y \supseteq X'} (-1)^{|Y \setminus X'|} F_+(Y). \quad (2.44)$$

Hence

$$F_-(S) = \sum_{X' \subseteq S} F(X') = \sum_{X' \subseteq S} \sum_{Y \supseteq X'} (-1)^{|Y \setminus X'|} F_+(Y) \quad (2.45a)$$

$$= \sum_Y F_+(Y) \sum_{X' \subseteq Y \cap S} (-1)^{|Y \setminus X'|} \quad (2.45b)$$

⁶It suffices to use repeatedly the following fact: If U_1 and U_2 are simply connected open subsets of \mathbb{R}^n , and $U_1 \cap U_2$ is nonempty and connected, then $U_1 \cup U_2$ is simply connected. For a proof of this lemma, see e.g. [11, pp. 46–47]. This is a special case of the Seifert–van Kampen theorem, which gives a recipe for computing the fundamental group (first homotopy group) of $U_1 \cup U_2$ in terms of the fundamental groups of U_1 , U_2 and $U_1 \cap U_2$ [79, Chapter 4].

$$= \sum_Y F_+(Y) (-1)^{|Y \cap S^c|} \sum_{X' \subseteq Y \cap S} (-1)^{|(Y \cap S) \setminus X'|} \quad (2.45c)$$

$$= \sum_Y F_+(Y) (-1)^{|Y \cap S^c|} \delta_{Y \cap S, \emptyset} \quad (2.45d)$$

$$= \sum_{Y \subseteq S^c} (-1)^{|Y|} F_+(Y). \quad (2.45e)$$

■

PROOF OF THEOREM 2.10. (c) \implies (b) \implies (a) is trivial.

(e) \implies (d) is trivial, while (d) \implies (e) follows from the alternating-sign property (2.29).

(e) implies that the sum of the Taylor series for $\log Z_W(\mathbf{w})$ defines an analytic function on the open polydisc $D_{\mathbf{R}}$ and a continuous function on the closed polydisc $\bar{D}_{\mathbf{R}}$. Its exponential equals $Z_W(\mathbf{w})$ on $D_{\mathbf{R}}$ and hence by continuity also on $\bar{D}_{\mathbf{R}}$. Therefore (e) \implies (c).

Finally, assume (a). Since Z_W is continuous on \mathbb{C}^X (and has real coefficients), we can find an open connected neighborhood C' of C in $(-\infty, 0]^X$ on which $Z_W > 0$, and an open neighborhood U of C' in $\mathbb{C}^X \simeq \mathbb{R}^{2|X|}$ on which $Z_W \neq 0$. Applying Lemma 2.12 (with $n = 2|X|$), we can find a finite polygonal path $P \subset C'$ from 0 to $-\mathbf{R}$ and a simply connected open set U' in \mathbb{C}^X satisfying $P \subset U' \subset U$. Then $\log Z_W$ is a well-defined single-valued analytic function on U' , once we specify $\log Z_W(\mathbf{0}) = 0$. Applying Proposition 2.11 to $\log Z_W$ on P and U' [using the alternating-sign property (2.29)], we conclude that the Taylor series for $\log Z_W$ around $\mathbf{0}$ is absolutely convergent on $\bar{D}_{\mathbf{R}}$. Therefore (a) \implies (e).

The bound $|Z_W(\mathbf{w})| \geq Z_W(-\mathbf{R})$ for $|\mathbf{w}| \leq \mathbf{R}$, which is equivalent to $\operatorname{Re} \log Z_W(\mathbf{w}) \geq \log Z_W(-\mathbf{R})$, is an immediate consequence of the alternating-sign property (2.29).

Now consider the special case of a hard-core self-repulsion. (b) \implies (b') is trivial, and (b') \implies (b) follows from the fact that Z_W is multiaffine (i.e. of degree ≤ 1 in each w_x separately) because the value of Z_W at any point \mathbf{w} in the rectangle $-\mathbf{R} \leq \mathbf{w} \leq \mathbf{0}$ is a convex combination of the values at the vertices.

To show that (b) \implies (f), note that

$$Z_W(\mathbf{w}; S) = \left(\prod_{x \in S} w_x \right) \left(\prod_{\{x, y\} \subseteq S} W(x, y) \right) Z_W(W(S, \cdot) \mathbf{w}) \quad (2.46)$$

where we have defined

$$[W(S, \cdot) \mathbf{w}]_y = \left(\prod_{x \in S} W(x, y) \right) w_y \quad (2.47)$$

(note in particular that this vanishes whenever $y \in S$). Hence

$$(-1)^{|S|} Z_W(-\mathbf{R}; S) = \left(\prod_{x \in S} R_x \right) \left(\prod_{\{x, y\} \subseteq S} W(x, y) \right) Z_W(-W(S, \cdot) \mathbf{R}) \geq 0 \quad (2.48)$$

since $-\mathbf{R} \leq -W(S, \cdot)\mathbf{R} \leq 0$, with strict inequality when $|S| = 0$ or 1 [since the product over $W(x, y)$ is in that case empty].

To show that (f) \implies (b'), use Lemma 2.13 applied to the set function

$$F(S) = \left(\prod_{x \in S} -R_x \right) \left(\prod_{\{x, y\} \subseteq S} W(x, y) \right). \quad (2.49)$$

We have

$$F_-(S) = Z_W(-\mathbf{R} \mathbf{1}_S) \quad (2.50)$$

$$F_+(S) = Z_W(-\mathbf{R}; S) \quad (2.51)$$

so that Lemma 2.13 asserts the identity

$$Z_W(-\mathbf{R} \mathbf{1}_S) = \sum_{Y \subseteq S^c} (-1)^{|Y|} Z_W(-\mathbf{R}; Y). \quad (2.52)$$

By (f), the $Y = \emptyset$ term is > 0 and the other terms are ≥ 0 , so $Z_W(-\mathbf{R} \mathbf{1}_S) > 0$ for all S .

Finally, let us show that (f) \iff (g). By inclusion-exclusion, there are unique numbers $P(T)$ satisfying (2.38), namely $P(T) = (-1)^{|T|} Z_W(-\mathbf{R}; T)$. Moreover, taking $S = \emptyset$ in (2.38) we see that $\sum_T P(T) = 1$. Therefore, P is a probability measure if and only if $(-1)^{|T|} Z_W(-\mathbf{R}; T) \geq 0$ for all T ; and $P(\emptyset) > 0$ if and only if $Z_W(-\mathbf{R}; \emptyset) = Z_W(-\mathbf{R}) > 0$. ■

2.4 Properties of the set $\mathcal{R}(W)$

The following definition plays a central role in the remainder of this paper:

Definition 2.14 (definition of $\mathcal{R}(W)$) *We define $\mathcal{R}(W)$ to be the set of all vectors $\mathbf{R} \geq 0$ satisfying the equivalent conditions (a)–(e) of Theorem 2.10. When W is the hard-core pair interaction for a graph G ,*

$$W(x, y) = \begin{cases} 0 & \text{if } x = y \text{ or } xy \in E(G) \\ 1 & \text{if } x \neq y \text{ and } xy \notin E(G) \end{cases} \quad (2.53)$$

we also write $\mathcal{R}(G)$.

Proposition 2.15 (elementary properties of $\mathcal{R}(W)$) *For any repulsive lattice gas, the set $\mathcal{R}(W)$ is*

(a) *open in $[0, \infty)^X$*

(b) *a down-set [i.e. $\mathbf{R} \in \mathcal{R}(W)$ and $0 \leq \mathbf{R}' \leq \mathbf{R}$ imply $\mathbf{R}' \in \mathcal{R}(W)$]*

(c) *logarithmically convex* [i.e. $\mathbf{R}, \mathbf{R}' \in \mathcal{R}(W)$ and $0 \leq \lambda \leq 1$ imply $\mathbf{R}^\lambda (\mathbf{R}')^{1-\lambda} \in \mathcal{R}(W)$].

Moreover, the set $\mathcal{R}(W)$ is an increasing function of each $W(x, y)$ on $0 \leq W \leq 1$.

PROOF. It is obvious from Theorem 2.10(b) or (c) that $\mathcal{R}(W)$ is open in $[0, \infty)^X$ and is a down-set.

Given any formal power series $f(\mathbf{z}) = \sum_{\mathbf{n}} a_{\mathbf{n}} \mathbf{z}^{\mathbf{n}}$ with complex coefficients, let us define its set of radii of absolute convergence by

$$\text{cvg}(f) = \{ \mathbf{R} \geq 0: \sum_{\mathbf{n}} |a_{\mathbf{n}}| \mathbf{R}^{\mathbf{n}} < \infty \}. \quad (2.54)$$

It then follows immediately from Hölder's inequality for infinite series that $\text{cvg}(f)$ is logarithmically convex.⁷ By Theorem 2.10(e) we have $\mathcal{R}(W) = \text{cvg}(\log Z_W)$, so $\mathcal{R}(W)$ is logarithmically convex.

Finally, from (2.29)/(2.30) we see that $|c_{\mathbf{n}}(W)|$ is a decreasing function of each $W(x, y)$ on $0 \leq W \leq 1$, so $\mathcal{R}(W) = \text{cvg}(\log Z_W)$ is an increasing function of each $W(x, y)$ on $0 \leq W \leq 1$. ■

See Corollary 2.19 below for a strict monotonicity that strengthens Proposition 2.15(b); and see Corollary 2.29 for an additional convexity property in the special case of a hard-core self-repulsion.

Remark. If Λ is any nonempty subset of X , we can obviously define a lattice gas on Λ with interaction $W \upharpoonright \Lambda$, and there will be a corresponding set $\mathcal{R}(W \upharpoonright \Lambda) \subseteq [0, \infty)^\Lambda$. Now, for any $\mathbf{w} \in \mathbb{C}^\Lambda$, we trivially have $Z_{W \upharpoonright \Lambda}(\mathbf{w}) = Z_W(\mathbf{w}, \mathbf{0})$ where $\mathbf{0} \in \mathbb{C}^{X \setminus \Lambda}$; hence $\mathbf{R} \in \mathcal{R}(W \upharpoonright \Lambda)$ if and only if $(\mathbf{R}, \mathbf{0}) \in \mathcal{R}(W)$. So the sets $\mathcal{R}(W \upharpoonright \Lambda)$ are the sections through $\mathbf{0}$ of $\mathcal{R}(W)$.

Next let us show that the sets $\mathcal{R}(W)$ are bounded in a suitable sense. In the case of hard-core self-repulsion this is immediate, because $\mathcal{R}(W) \subseteq [0, 1]^X$. [This containment is trivial if $|X| = 1$, and otherwise follows from Proposition 2.15(b). Alternatively, it is trivial if $W(x, y) = 1$ for all $x \neq y$, and otherwise follows from the monotonicity statement in the last sentence of Proposition 2.15.] However, in the general case the set $\mathcal{R}(W)$ can be unbounded, as is shown by the following two examples:

- (a) Let $X = \{x\}$ and $W(x, x) = 1$; then $Z_W(w) = e^w$ and $\mathcal{R}(W) = [0, \infty)$.
- (b) Let $X = \{x, y\}$, $W(x, x) = W(y, y) = 1$ and $W(x, y) = 0$; then $Z_W(w_x, w_y) = e^{w_x} + e^{w_y} - 1$ and $\mathcal{R}(W) = \{(R_x, R_y): R_x, R_y \geq 0 \text{ and } e^{-R_x} + e^{-R_y} > 1\}$. This set is unbounded, since R_x can go to ∞ as $R_y \rightarrow 0$ (and vice versa). [When $R_y = 0$ this reduces to example (a).]

⁷See [49, Chapters B and G], [73, Sections 2.3 and 3.4.3], [92, Section II.3.8] and [114, pp. 116–122] for further discussion of related questions in the theory of analytic functions of several complex variables.

We now claim that these are essentially the only ways in which $\mathcal{R}(W)$ can be unbounded:

Proposition 2.16 (boundedness of $\mathcal{R}(W)$) *Consider any repulsive lattice gas, and let $x \in X$.*

- (a) *If $W(x, x) < 1$, then there exists $C_x < \infty$ such that $\mathcal{R}(W) \subseteq [0, C_x) \times [0, \infty)^{X \setminus x}$. [In particular, if $W(x, x) < 1$ for all $x \in X$, then $\mathcal{R}(W)$ is bounded.]*
- (b) *Suppose that $W(x, x) = 1$, and let $\mathbf{a} \in [0, \infty)^{X \setminus x}$. If there exists at least one $y \in X$ such that $W(x, y) < 1$ and $a_y > 0$, then the section $\{R_x: (R_x, \mathbf{a}) \in \mathcal{R}(W)\}$ is bounded. Moreover, the converse is true provided that $\mathbf{a} \in \mathcal{R}(W \upharpoonright (X \setminus x))$.*

Remark. Since $\mathcal{R}(W)$ is a down-set, this proposition can alternatively be formulated as follows: Let $\mathbf{a}, \mathbf{b} \in [0, \infty)^X$; then the set $\{\lambda \geq 0: \mathbf{a} + \lambda \mathbf{b} \in \mathcal{R}(W)\}$ is bounded if there exists $x \in \text{supp } \mathbf{b}$ and $y \in \text{supp } \mathbf{a} \cup \text{supp } \mathbf{b}$ such that $W(x, y) < 1$; and the converse is true provided that $\mathbf{a} \in \mathcal{R}(W)$.

PROOF. (a) Using the fact that $\mathcal{R}(W)$ is a down-set, it suffices to prove the claim for the single-site partition function

$$Z_W(w) = \sum_{n=0}^{\infty} \frac{W^{n(n-1)/2}}{n!} w^n \quad (2.55)$$

where $0 \leq W < 1$. If $W = 0$, we have $Z_W(w) = 1 + w$, so that $\mathcal{R}(W) = [0, 1)$. If $0 < W < 1$, then Z_W is a nonpolynomial entire function of order 0 [76, Theorem I.2]; and by the Hadamard factorization theorem [76, Theorem I.13], any such function must have infinitely many zeros. In particular, $\mathcal{R}(W) = [0, \alpha)$ where α is the smallest absolute value of a zero of Z_W .⁸

(b) If there does not exist $y \in X$ such that $W(x, y) < 1$ and $a_y > 0$, then $Z_W(\mathbf{w}) = e^{w_x} Z_{W, X \setminus x}(\mathbf{w}_{\neq x})$ whenever $\text{supp } \mathbf{w} \subseteq \text{supp } \mathbf{a} \cup \{x\}$. It follows that the section $\{R_x: (R_x, \mathbf{a}) \in \mathcal{R}(W)\}$ is either empty [in case $\mathbf{a} \notin \mathcal{R}(W \upharpoonright (X \setminus x))$] or all of $[0, \infty)$ [in case $\mathbf{a} \in \mathcal{R}(W \upharpoonright (X \setminus x))$].

Now suppose that there does exist such a y . Using the fact that $\mathcal{R}(W)$ is a down-set, it suffices to prove the claim for the two-site partition function with $\Lambda = \{x, y\}$. Moreover, by the monotonicity statement in the last sentence of Proposition 2.15, we may assume that $W(y, y) = 1$. Writing $W \equiv W(x, y) \in [0, 1)$, we need to treat

$$Z_W(w_x, w_y) = \sum_{n_x, n_y=0}^{\infty} \frac{W^{n_x n_y}}{n_x! n_y!} w_x^{n_x} w_y^{n_y} \quad (2.56a)$$

$$= \sum_{n_x=0}^{\infty} \frac{w_x^{n_x} \exp(w_y W^{n_x})}{n_x!} \quad (2.56b)$$

⁸It turns out (though we do not need this fact here) that all the zeros of Z_W are negative real numbers [85, 66].

If $w_y \neq 0$, this is an entire function of order 1 in w_x that is not of the form $e^{\alpha w_x} P(w_x)$ for any $\alpha \in \mathbb{C}$ and polynomial P .⁹ It again follows from the Hadamard factorization theorem that $Z_W(\cdot, w_y)$ has infinitely many zeros. Choosing any $w_y \in \mathbb{C}$ with $|w_y| = a_y$ (which by hypothesis is nonzero), we conclude from Theorem 2.10(c) that the section $\{R_x: (R_x, a_y) \in \mathcal{R}(W)\}$ is bounded. ■

Let $\overline{\mathcal{R}(W)}$ be the closure of $\mathcal{R}(W)$, and let $\partial\mathcal{R}(W) = \overline{\mathcal{R}(W)} \setminus \mathcal{R}(W)$. [Note that this is the boundary of $\mathcal{R}(W)$ in $[0, \infty)^X$, not in \mathbb{R}^X .] Since $\mathcal{R}(W)$ is a down-set, it is obvious that $\overline{\mathcal{R}(W)}$ is a down-set as well.

Proposition 2.17 (properties of closure and boundary of $\mathcal{R}(W)$) *For any repulsive lattice gas:*

(a) *If $-\mathbf{w} \in \overline{\mathcal{R}(W)}$, then*

$$\frac{\partial^n Z_W(\mathbf{w})}{\partial w_{x_1} \cdots \partial w_{x_n}} \geq 0 \quad (2.57)$$

for all $n \geq 0$ and all $x_1, \dots, x_n \in X$.

(b) *If $-\mathbf{w} \in \partial\mathcal{R}(W)$, then $Z_W(\mathbf{w}) = 0$.*

PROOF. (a) We have $Z_W(\mathbf{w}) > 0$ for $-\mathbf{w} \in \mathcal{R}(W)$, so by continuity $Z_W(\mathbf{w}) \geq 0$ for $-\mathbf{w} \in \overline{\mathcal{R}(W)}$. The general claim (2.57) then follows from the multiple differentiation identity (2.6), using the facts that $0 \leq W \leq 1$ and that $\overline{\mathcal{R}(W)}$ is a down-set.

(b) If $Z_W(\mathbf{w})$ were strictly positive, then $-\mathcal{R}(W) \cup \{\mathbf{w}\}$ would be a connected subset of $(-\infty, 0]^X$ on which Z_W is strictly positive, so by Theorem 2.10(a) we would have $-\mathbf{w} \in \mathcal{R}(W)$. ■

⁹PROOF. If $P(z) = \sum_{k=0}^K a_k z^k$, then

$$\begin{aligned} e^{\alpha z} P(z) &= \sum_{n=0}^{\infty} \frac{(\alpha z)^n}{n!} \left[a_0 + n \frac{a_1}{\alpha} + n(n-1) \frac{a_2}{\alpha^2} + \dots + n(n-1) \cdots (n-K+1) \frac{a_K}{\alpha^K} \right] \\ &\equiv \sum_{n=0}^{\infty} \frac{\alpha^n Q(n)}{n!} z^n \end{aligned}$$

where Q is a polynomial. But $F(n) = \exp(w_y W^n)$ is not of the form $\alpha^n Q(n)$ for any $\alpha \in \mathbb{C}$ and polynomial Q .

ALTERNATE PROOF. We have

$$Z_W(w_x, w_y) - e^{w_x} = \sum_{n_x=0}^{\infty} \frac{w_x^{n_x}}{n_x!} [\exp(w_y W^{n_x}) - 1]$$

Since w_y is fixed and $0 \leq W \leq 1$, we have $|\exp(w_y W^{n_x}) - 1| \leq C W^{n_x}$ for some constant $C < \infty$, from which it follows that

$$|Z_W(w_x, w_y) - e^{w_x}| \leq C e^{W|w_x|}.$$

If $Z_W(w_x, w_y)$ were equal to $e^{\alpha w_x} P(w_x)$, then by taking $w_x \rightarrow +\infty$ we conclude that we would have to have $\alpha = 1$ and $P \equiv 1$ (since $W < 1$); but Z_W is not in fact of this form when $w_y \neq 0$.

Under an appropriate connectivity condition, we can prove that the inequality (2.57) is strict for $n = 1$:

Proposition 2.18 *Consider any repulsive lattice gas. Let $\mathbf{R} \in \overline{\mathcal{R}(W)}$, and suppose that the induced subgraph $G_W[\text{supp } \mathbf{R}]$ is connected. Then*

$$\left. \frac{\partial Z_W(\mathbf{w})}{\partial w_x} \right|_{\mathbf{w}=-\mathbf{R}} > 0 \quad \text{for all } x \in \text{supp } \mathbf{R}. \quad (2.58)$$

Here the connectedness hypothesis is essential, as is shown by taking G to be the edgeless graph \bar{K}_n , for which $Z_G(\mathbf{w}) = \prod_{i=1}^n (1 + w_i)$, and taking $\mathbf{R} = (1, \dots, 1)$. It is also essential to consider the induced subgraph $G_W[\text{supp } \mathbf{R}]$, since a site can be effectively eliminated by setting its fugacity to zero: for example, when G is the path 123, we have $Z_G(\mathbf{w}) = (1 + w_1)(1 + w_3) + w_2$, and a counterexample to (2.58) is obtained by taking $R_2 = 0$ and $R_1 = R_3 = 1$.

Before proving Proposition 2.18, let us state and prove some corollaries.

Corollary 2.19 (strict monotonicity) *Consider any repulsive lattice gas. Let $\mathbf{R} \in \overline{\mathcal{R}(W)}$, and suppose that the induced subgraph $G_W[\text{supp } \mathbf{R}]$ is connected. If $\mathbf{0} \leq \mathbf{R}' \leq \mathbf{R}$ with $R'_x < R_x$ for at least one x , then $\mathbf{R}' \in \mathcal{R}(W)$.*

PROOF. Since $Z_W(-\mathbf{R}) \geq 0$ and $\partial Z_W(\mathbf{w})/\partial w_x|_{\mathbf{w}=-\mathbf{R}} > 0$ by Propositions 2.17(a) and 2.18, we have $Z_W(-(\mathbf{R} - \epsilon \boldsymbol{\delta}_x)) > 0$ for all sufficiently small $\epsilon > 0$ [here $\boldsymbol{\delta}_x$ is the vector with x th coordinate 1 and all other coordinates 0]. Since $\overline{\mathcal{R}(W)}$ is a down-set, we have $\mathbf{R} - \epsilon \boldsymbol{\delta}_x \in \overline{\mathcal{R}(W)}$; but by Proposition 2.17(b), $\mathbf{R} - \epsilon \boldsymbol{\delta}_x$ cannot lie in $\partial \mathcal{R}(W)$, so we must have $\mathbf{R} - \epsilon \boldsymbol{\delta}_x \in \mathcal{R}(W)$. Now pick ϵ small enough so that $\mathbf{R}' \leq \mathbf{R} - \epsilon \boldsymbol{\delta}_x$ and use the fact that $\mathcal{R}(W)$ is a down-set. ■

Corollary 2.20 *Consider any repulsive lattice gas.*

- (a) *If $\mathbf{R} \in \partial \mathcal{R}(W)$, then every neighborhood of $-\mathbf{R}$ in $(-\infty, 0]^X$ contains points $\mathbf{w} \neq -\mathbf{R}$ with $\text{supp } \mathbf{w} = \text{supp } \mathbf{R}$ where $Z_W > 0$, where $Z_W = 0$ and where $Z_W < 0$.*
- (b) *$\mathbf{R} \in \overline{\mathcal{R}(W)}$ if and only if $Z_W(\mathbf{w}) \geq 0$ for all \mathbf{w} satisfying $-\mathbf{R} \leq \mathbf{w} \leq \mathbf{0}$.*

PROOF. (a) Let X_1, \dots, X_k be the vertex sets of the components of $G_W[\text{supp } \mathbf{R}]$. For any \mathbf{w} with $\text{supp } \mathbf{w} \subseteq \text{supp } \mathbf{R}$, we have $Z_W(\mathbf{w}) = \prod_{i=1}^k Z_W(\mathbf{w} \mathbf{1}_{X_i})$. Pick a vertex x_i in each X_i , and let $\mathbf{w}_{\pm} = -\mathbf{R} \pm \epsilon \boldsymbol{\delta}_{x_1} + \epsilon \sum_{i=2}^k \boldsymbol{\delta}_{x_i}$. By Proposition 2.18, for all sufficiently small $\epsilon > 0$ we have $\pm Z_W(\mathbf{w} \mathbf{1}_{X_1}) > 0$ and $Z_W(\mathbf{w} \mathbf{1}_{X_i}) > 0$ for $2 \leq i \leq k$. This proves the existence of points near $-\mathbf{R}$ with $Z_W > 0$ and $Z_W < 0$. The existence of points $\mathbf{w} \neq -\mathbf{R}$ with $Z_W = 0$ then follows by the intermediate value theorem.

(b) “Only if” is obvious from Proposition 2.17(a) together with the fact that $\overline{\mathcal{R}(W)}$ is a down-set. To prove “if”, suppose that $Z_W(\mathbf{w}) \geq 0$ for all \mathbf{w} satisfying $-\mathbf{R} \leq \mathbf{w} \leq \mathbf{0}$. Consider the line segment $\{\lambda \mathbf{R}\}_{0 \leq \lambda \leq 1}$, and let $\lambda_{\max} = \sup\{\lambda \in [0, 1]: \lambda \mathbf{R} \in$

$\mathcal{R}(W)\}$. If $\lambda_{\max} = 1$, then $\mathbf{R} \in \overline{\mathcal{R}(W)}$. If $\lambda_{\max} < 1$, then $\lambda_{\max}\mathbf{R} \in \partial\mathcal{R}(W)$, so by part (a) we can choose \mathbf{w} arbitrarily close to $-\lambda_{\max}\mathbf{R}$ with $\text{supp } \mathbf{w} = \text{supp } \mathbf{R}$ (and in particular satisfying $-\mathbf{R} \leq \mathbf{w} \leq \mathbf{0}$) such that $Z_W(\mathbf{w}) < 0$ — a contradiction. ■

Remark. Corollary 2.20(b) is an analogue of Theorem 2.10(b) for the closure of $\mathcal{R}(W)$. It is worth noting that the analogue of Theorem 2.10(a) is *false*: for example, for $G = K_2$ we have $Z_G(\mathbf{w}) = (1+w_1)(1+w_2)$, so that the set $\{\mathbf{R} \geq \mathbf{0}: Z_G(-\mathbf{R}) \geq 0\}$ is connected and equals $[0, 1]^2 \cup [1, \infty)^2$, while $\overline{\mathcal{R}(W)} = [0, 1]^2$ only.

PROOF OF PROPOSITION 2.18. If $\mathbf{R} \in \mathcal{R}(W)$, then $W(x, \cdot)\mathbf{R} \in \mathcal{R}(W)$ as well [since $0 \leq W \leq 1$ and $\mathcal{R}(W)$ is a down-set], so (2.58) follows from the differentiation identity (2.5). So we can assume henceforth that $\mathbf{R} \in \partial\mathcal{R}(W)$. By deleting sites where $R_x = 0$, we can also assume without loss of generality that $\text{supp } \mathbf{R} = X$. Let us define

$$S = \{x \in X: \left. \frac{\partial Z_W(\mathbf{w})}{\partial w_x} \right|_{\mathbf{w}=-\mathbf{R}} = 0\} \quad (2.59)$$

$$S' = \{x \in X: Z_W(w_x, -\mathbf{R}_{\neq x}) = 0 \text{ for all } w_x \in \mathbb{C}\} \quad (2.60)$$

Clearly $S' \subseteq S \subseteq X$. The Proposition will then be an immediate consequence of Lemmas 2.21 and 2.22 below. ■

Lemma 2.21 *Let $\mathbf{R} \in \partial\mathcal{R}(W)$, and let S, S' be defined as in (2.59)/(2.60). If $\text{supp } \mathbf{R} = X$ and G_W is connected, then either $S' = S = \emptyset$ or else $S' = S = X$.*

Lemma 2.22 *Suppose $\mathbf{R} \in \overline{\mathcal{R}(W)}$ and for all $x \in X$ and all $w_x \in \mathbb{C}$, we have $Z_W(w_x, -\mathbf{R}_{\neq x}) = 0$. Then $G_W[\text{supp } \mathbf{R}]$ is disconnected.*

PROOF OF LEMMA 2.21. Suppose $S \neq \emptyset$, and consider any $x \in S$. Then $Z_W(-\mathbf{R}) = 0$ and $Z_W(-W(x, \cdot)\mathbf{R}) = \left. \frac{\partial Z_W(\mathbf{w})}{\partial w_x} \right|_{\mathbf{w}=-\mathbf{R}} = 0$ [using the differentiation identity (2.5)]. By monotonicity of Z_W in $\overline{\mathcal{R}(W)}$, we have $Z_W(-\mathbf{R}') = 0$ whenever $-\mathbf{R} \leq -\mathbf{R}' \leq -W(x, \cdot)\mathbf{R}$. In particular, if y is adjacent to x in G_W [i.e. $W(x, y) < 1$], then $Z_W(w_y, -\mathbf{R}_{\neq y}) = 0$ for all $w_y \in [-R_y, -W(x, y)R_y]$. By analyticity it follows that $Z_W(w_y, -\mathbf{R}_{\neq y}) = 0$ for all $w_y \in \mathbb{C}$, i.e. $y \in S' \subseteq S$. Since G_W is connected, it easily follows that $S' = S = X$. ■

PROOF OF LEMMA 2.22. As before, we may assume without loss of generality that $\text{supp } \mathbf{R} = X$. The proof is by induction on $|X|$. If $|X| = 1$, the hypothesis is impossible, since $Z_W(0) = 1 \neq 0$. So let $|X| > 1$ and suppose that G_W is connected. Choose a vertex $x \in X$ such that $G_W \setminus x$ is connected [e.g. let x be any endvertex of a spanning tree in G_W]. By the fundamental identity (2.3), we have

$$\begin{aligned} Z_W(w_x, -\mathbf{R}_{\neq x}) &= Z_W(0, -\mathbf{R}_{\neq x}) + w_x Z_W(0, -W(x, \cdot)\mathbf{R}_{\neq x}) \\ &\quad + \text{terms of order } w_x^2 \text{ and higher} \end{aligned} \quad (2.61)$$

Since $Z_W(\cdot, -\mathbf{R}_{\neq x})$ is identically zero, all coefficients in the Taylor expansion (2.61) must vanish; in particular we must have

$$Z_W(0, -\mathbf{R}_{\neq x}) = 0 \quad (2.62a)$$

$$Z_W(0, -W(x, \cdot)\mathbf{R}_{\neq x}) = 0 \quad (2.62b)$$

Pick $y \neq x$ with $W(x, y) < 1$ [which is possible since x is not an isolated vertex of G_W]. Since $R_y > 0$ and $Z_W(0, w_y, -\mathbf{R}_{\neq x, y})$ is monotonic in w_y for $w_y \in [-R_y, 0]$, it must be vanishing on the nontrivial interval $[-R_y, -W(x, y)R_y]$. By analyticity of Z_W , it follows that $Z_W(0, w_y, -\mathbf{R}_{\neq x, y}) = 0$ for all $w_y \in \mathbb{C}$, so in particular

$$Z_W(0, -R_y, -\mathbf{R}_{\neq x, y}) = 0 \quad (2.63a)$$

$$\left. \frac{\partial Z_W(0, w_y, -\mathbf{R}_{\neq x, y})}{\partial w_y} \right|_{w_y = -R_y} = 0 \quad (2.63b)$$

Notice that $\mathbf{R} \equiv (R_x, \mathbf{R}_{\neq x}) \in \overline{\mathcal{R}(W)}$ implies $(0, \mathbf{R}_{\neq x}) \in \overline{\mathcal{R}(W)}$ [since $\overline{\mathcal{R}(W)}$ is a down-set], or in other words $\mathbf{R}_{\neq x} \in \overline{\mathcal{R}(W \upharpoonright (X \setminus x))}$. Moreover, by (2.63a) $\mathbf{R}_{\neq x}$ must lie in $\partial\mathcal{R}(W \upharpoonright (X \setminus x))$. So we can apply Lemma 2.21 to $X \setminus x$ to conclude that $S' = X \setminus x$; therefore, by the inductive hypothesis, $G_W \setminus x$ is disconnected, which contradicts the choice of x . ■

Remark. One might think that Lemma 2.22 would hold not only for $\mathbf{R} \in \overline{\mathcal{R}(W)}$ but for arbitrary $\mathbf{w}^{(0)} \in \mathbb{C}^X$. However, this is false: Consider the star $K_{1,4}$ with center x and endvertices y_1, \dots, y_4 with interactions $W \in (0, 1)$ along the edges [and $W(x, x) = W(y_i, y_i) = 0$]. Then

$$Z_W(w_x, w_{y_1}, \dots, w_{y_4}) = \prod_{i=1}^4 (1 + w_{y_i}) + w_x \prod_{i=1}^4 (1 + W w_{y_i}). \quad (2.64)$$

Setting $w_{y_1}^{(0)} = w_{y_2}^{(0)} = -1$ and $w_{y_3}^{(0)} = w_{y_4}^{(0)} = -1/W$ [here $w_x^{(0)}$ is arbitrary], we see that $Z_W(w_z, \mathbf{w}_{\neq z}^{(0)}) = 0$ for every vertex z and every $w_z \in \mathbb{C}$.

2.5 Further consequences of the alternating-sign property

Let us now exploit systematically the alternating-sign property (2.29) for the Taylor coefficients of $\log Z_W$. The general context is the following:

Definition 2.23 (absolute monotonicity) *Let $U \subset \mathbb{C}^n$ be a union of open polydiscs centered at $\mathbf{0}$ (a “complete Reinhardt domain”). We say that f is absolutely monotone in U if it is analytic in U and all the Taylor coefficients of f at $\mathbf{0}$ are nonnegative.*

We will use the following essentially trivial result:

Lemma 2.24 (elementary consequences of absolute monotonicity) *Let f be absolutely monotone in a union U of open polydiscs centered at $\mathbf{0} \in \mathbb{C}^n$. Then:*

(a) *All the derivatives $D^\alpha f$ (where α is a multi-index) are absolutely monotone in U .*

(b) *For every $\lambda \in [0, 1]^n$, the function*

$$f_\lambda(\mathbf{z}) = f(\mathbf{z}) - f(\lambda\mathbf{z}) \quad (2.65)$$

is absolutely monotone in U . [Here $\lambda\mathbf{z}$ is the pointwise product, so that $(\lambda\mathbf{z})_x = \lambda_x z_x$.]

(c) *For all multi-indices α , we have*

$$|(D^\alpha f)(\mathbf{z})| \leq (D^\alpha f)(\mathbf{R}) \quad (2.66)$$

whenever $|\mathbf{z}| \leq \mathbf{R} \in U$. In particular, $(D^\alpha f)(\mathbf{R}) \geq 0$ whenever $0 \leq \mathbf{R} \in U$.

Now, the alternating-sign property (2.29) is precisely the statement that

$$f(\mathbf{z}) = -\log Z_W(-\mathbf{z}) \quad (2.67)$$

is absolutely monotone in the domain $D_{\mathcal{R}(W)} = \bigcup_{\mathbf{R} \in \mathcal{R}(W)} D_{\mathbf{R}} = \bigcup_{\mathbf{R} \in \mathcal{R}(W)} \bar{D}_{\mathbf{R}}$. Applying Lemma 2.24(c) to f , we obtain:

Proposition 2.25 *For any repulsive lattice gas:*

(a) *When $-\mathbf{w} \in \mathcal{R}(W)$, we have*

$$(-1)^{n-1} \frac{\partial^n \log Z_W(\mathbf{w})}{\partial w_{x_1} \cdots \partial w_{x_n}} \geq 0 \quad (2.68)$$

for all $n \geq 0$ and all $x_1, \dots, x_n \in X$.

(b) *When $|\mathbf{w}| \leq \mathbf{R} \in \mathcal{R}(W)$, we have*

$$\left| \frac{\partial^n \log Z_W(\mathbf{w})}{\partial w_{x_1} \cdots \partial w_{x_n}} \right| \leq (-1)^{n-1} \frac{\partial^n \log Z_W(\mathbf{w}')}{\partial w'_{x_1} \cdots \partial w'_{x_n}} \Big|_{\mathbf{w}' = -\mathbf{R}} \quad (2.69)$$

for all $n \geq 0$ and all $x_1, \dots, x_n \in X$.

PROOF. This is an immediate consequence of the alternating-sign property (2.29) together with Lemma 2.24(c) and Theorem 2.10(d). ■

Let us now define

$$Y_W(\mathbf{w}; \lambda) = \frac{Z_W(\lambda\mathbf{w})}{Z_W(\mathbf{w})}. \quad (2.70)$$

By Lemma 2.24(b), the function

$$f_{\boldsymbol{\lambda}}(\mathbf{z}) = \log Y_W(-\mathbf{z}; \boldsymbol{\lambda}) = \log \frac{Z_W(-\boldsymbol{\lambda}\mathbf{z})}{Z_W(-\mathbf{z})} \quad (2.71)$$

is absolutely monotone on $D_{\mathcal{R}(W)}$ whenever $\boldsymbol{\lambda} \in [0, 1]^X$. Applying Lemma 2.24(c) to $f_{\boldsymbol{\lambda}}$, we obtain:

Proposition 2.26 *For any repulsive lattice gas and any $\boldsymbol{\lambda} \in [0, 1]^X$:*

(a) *When $-\mathbf{w} \in \mathcal{R}(W)$, we have*

$$(-1)^n \frac{\partial^n \log Y_W(\mathbf{w}; \boldsymbol{\lambda})}{\partial w_{x_1} \cdots \partial w_{x_n}} \geq 0 \quad (2.72)$$

for all $n \geq 0$ and all $x_1, \dots, x_n \in X$.

(b) *When $|\mathbf{w}| \leq \mathbf{R} \in \mathcal{R}(W)$, we have*

$$\left| \frac{\partial^n \log Y_W(\mathbf{w}; \boldsymbol{\lambda})}{\partial w_{x_1} \cdots \partial w_{x_n}} \right| \leq (-1)^n \frac{\partial^n \log Y_W(\mathbf{w}'; \boldsymbol{\lambda})}{\partial w'_{x_1} \cdots \partial w'_{x_n}} \Big|_{\mathbf{w}' = -\mathbf{R}} \quad (2.73)$$

for all $n \geq 0$ and all $x_1, \dots, x_n \in X$. In particular, setting $n = 0$ we have

$$\left| \frac{Z_W(\boldsymbol{\lambda}\mathbf{w})}{Z_W(\mathbf{w})} \right| \leq \frac{Z_W(-\boldsymbol{\lambda}\mathbf{R})}{Z_W(-\mathbf{R})} < \infty \quad (2.74)$$

for all $\mathbf{w} \in \bar{D}_{\mathbf{R}}$. That is, the quantity $|Z_W(\boldsymbol{\lambda}\mathbf{w})/Z_W(\mathbf{w})|$ takes its maximum on the polydisc $\bar{D}_{\mathbf{R}}$ at the point $\mathbf{w} = -\mathbf{R}$, and this maximum value $Z_W(-\boldsymbol{\lambda}\mathbf{R})/Z_W(-\mathbf{R})$ is therefore an increasing function of $\mathbf{R} \in \mathcal{R}(W)$.

Let us define the function $\mathcal{Z}_W: [0, \infty)^X \rightarrow [0, \infty)$ by

$$\mathcal{Z}_W(\mathbf{R}) = \inf_{-\mathbf{R} \leq \mathbf{w} \leq 0} \max[Z_W(\mathbf{w}), 0] \quad (2.75a)$$

$$= \inf_{\mathbf{w} \in \bar{D}_{\mathbf{R}}} |Z_W(\mathbf{w})| \quad (2.75b)$$

$$= \begin{cases} Z_W(-\mathbf{R}) & \text{for } \mathbf{R} \in \mathcal{R}(W) \\ 0 & \text{for } \mathbf{R} \notin \mathcal{R}(W) \end{cases} \quad (2.75c)$$

where the equivalence of these three expressions is an immediate consequence of Theorem 2.10. We then have:

Corollary 2.27 *For any repulsive lattice gas, the function \mathcal{Z}_W is*

(a) *continuous*

(b) *decreasing*

- (c) satisfies $\mathcal{Z}_W(\boldsymbol{\lambda}_1 \boldsymbol{\lambda}_2 \mathbf{R}) \mathcal{Z}_W(\mathbf{R}) \leq \mathcal{Z}_W(\boldsymbol{\lambda}_1 \mathbf{R}) \mathcal{Z}_W(\boldsymbol{\lambda}_2 \mathbf{R})$ for all $\boldsymbol{\lambda}_1, \boldsymbol{\lambda}_2 \in [0, 1]^X$ and $\mathbf{R} \in [0, \infty)^X$
- (d) log submodular [i.e. $\mathcal{Z}_W(\mathbf{R}_1 \wedge \mathbf{R}_2) \mathcal{Z}_W(\mathbf{R}_1 \vee \mathbf{R}_2) \leq \mathcal{Z}_W(\mathbf{R}_1) \mathcal{Z}_W(\mathbf{R}_2)$, where \wedge (resp. \vee) denotes the elementwise min (resp. max) of vectors]

PROOF. (a) The continuity of \mathcal{Z}_W follows easily from (2.75a) or (2.75b) and the continuity of Z_W . [For any $r < \infty$, the function Z_W is uniformly continuous on the compact ball $B = \{\mathbf{w} \in \mathbb{C}: |\mathbf{w}| \leq r\}$. If $\mathbf{R}, \mathbf{R}' \in B \cap [0, \infty)^X$ with $|\mathbf{R} - \mathbf{R}'| \leq \delta$, then $|\mathcal{Z}_W(\mathbf{R}) - \mathcal{Z}_W(\mathbf{R}')| \leq \sup_{\substack{\mathbf{w}, \mathbf{w}' \in B \\ |\mathbf{w} - \mathbf{w}'| \leq \delta}} |Z_W(\mathbf{w}) - Z_W(\mathbf{w}')|$. Uniform continuity of Z_W on

B therefore implies uniform continuity of \mathcal{Z}_W on $B \cap [0, \infty)^X$.]

(b) The decreasing property is an immediate consequence of (2.75a) or (2.75b).

(c) This is trivial if $\mathbf{R} \notin \mathcal{R}(W)$, so assume $\mathbf{R} \in \mathcal{R}(W)$. Then $Y_W(-\mathbf{R}; \boldsymbol{\lambda}_2) = \mathcal{Z}_W(\boldsymbol{\lambda}_2 \mathbf{R}) / \mathcal{Z}_W(\mathbf{R})$, while $Y_W(-\boldsymbol{\lambda}_1 \mathbf{R}; \boldsymbol{\lambda}_2) = \mathcal{Z}_W(\boldsymbol{\lambda}_1 \boldsymbol{\lambda}_2 \mathbf{R}) / \mathcal{Z}_W(\boldsymbol{\lambda}_1 \mathbf{R})$, so that the claim follows from Proposition 2.26 with $n = 1$.

(d) The log submodularity follows immediately from (c) by setting $\mathbf{R} = \mathbf{R}_1 \vee \mathbf{R}_2$ and $\boldsymbol{\lambda}_i = \mathbf{R}_i / \mathbf{R}$ for $i = 1, 2$ (setting $0/0 = 1$ where needed). [Alternatively, for $\mathbf{R}_1 \vee \mathbf{R}_2 \in \mathcal{R}(W)$ it follows by integrating (2.68) with $n = 2$ and $x_1 \neq x_2$.] ■

We can also prove a strict version of the cases $n = 0$ of Propositions 2.25(a) and 2.26(a) as well as Corollary 2.27(c). Note first that

$$-\log Z_W(-\mathbf{R}) = \sum_{\mathbf{n}} (-1)^{|\mathbf{n}|-1} c_{\mathbf{n}}(W) \mathbf{R}^{\mathbf{n}} \quad (2.76)$$

$$-\log Z_W(-\mathbf{R}) + \log Z_W(-\boldsymbol{\lambda} \mathbf{R}) = \sum_{\mathbf{n}} (-1)^{|\mathbf{n}|-1} c_{\mathbf{n}}(W) (1 - \boldsymbol{\lambda}^{\mathbf{n}}) \mathbf{R}^{\mathbf{n}} \quad (2.77)$$

$$\begin{aligned} & -\log Z_W(-\mathbf{R}) + \log Z_W(-\boldsymbol{\lambda}_1 \mathbf{R}) + \log Z_W(-\boldsymbol{\lambda}_2 \mathbf{R}) - \log Z_W(-\boldsymbol{\lambda}_1 \boldsymbol{\lambda}_2 \mathbf{R}) \\ &= \sum_{\mathbf{n}} (-1)^{|\mathbf{n}|-1} c_{\mathbf{n}}(W) (1 - \boldsymbol{\lambda}_1^{\mathbf{n}})(1 - \boldsymbol{\lambda}_2^{\mathbf{n}}) \mathbf{R}^{\mathbf{n}} \end{aligned} \quad (2.78)$$

By Proposition 2.8, all three quantities are nonnegative whenever $\mathbf{R} \in \mathcal{R}(W)$ and $\boldsymbol{\lambda}, \boldsymbol{\lambda}_1, \boldsymbol{\lambda}_2 \in [0, 1]^X$. We can determine when they are strictly positive:

Proposition 2.28 *Consider any repulsive lattice gas, and let $\mathbf{R} \in \mathcal{R}(W)$ and $\boldsymbol{\lambda}, \boldsymbol{\lambda}_1, \boldsymbol{\lambda}_2 \in [0, 1]^X$. Then:*

- (a) $-\log Z_W(-\mathbf{R}) > 0$ if and only if $\mathbf{R} \neq \mathbf{0}$.
- (b) $-\log Z_W(-\mathbf{R}) + \log Z_W(-\boldsymbol{\lambda} \mathbf{R}) > 0$ if and only if $\text{supp } \mathbf{R} \cap \text{supp}(\mathbf{1} - \boldsymbol{\lambda}) \neq \emptyset$.
- (c) $-\log Z_W(-\mathbf{R}) + \log Z_W(-\boldsymbol{\lambda}_1 \mathbf{R}) + \log Z_W(-\boldsymbol{\lambda}_2 \mathbf{R}) - \log Z_W(-\boldsymbol{\lambda}_1 \boldsymbol{\lambda}_2 \mathbf{R}) > 0$ if and only there exists a component of $G_W[\text{supp } \mathbf{R}]$ that meets both $\text{supp}(\mathbf{1} - \boldsymbol{\lambda}_1)$ and $\text{supp}(\mathbf{1} - \boldsymbol{\lambda}_2)$.

PROOF. We examine the terms in (2.76)–(2.78) and determine (for some of them) when they are nonzero.

(a) “Only if” is trivial; for “if”, just consider the term $\mathbf{n} = \boldsymbol{\delta}_x$ for some $x \in \text{supp } \mathbf{R}$, and use Proposition 2.9(a).

(b) “Only if” is again trivial; for “if”, just consider $\mathbf{n} = \boldsymbol{\delta}_x$ for some $x \in \text{supp } \mathbf{R} \cap \text{supp}(\mathbf{1} - \boldsymbol{\lambda})$, and again use Proposition 2.9(a).

(c) For “if”, let S be the vertex set of a component of $G_W[\text{supp } \mathbf{R}]$ that meets both $\text{supp}(\mathbf{1} - \boldsymbol{\lambda}_1)$ and $\text{supp}(\mathbf{1} - \boldsymbol{\lambda}_2)$, take $\mathbf{n} = \mathbf{1}_S$, and use Proposition 2.9(a) or 2.9(c). For “only if”, suppose that $c_{\mathbf{n}}(W) \neq 0$. Then, by Proposition 2.9, either $\mathbf{n} = \boldsymbol{\delta}_x$ for some $x \in X$, or $\mathbf{n} = k\boldsymbol{\delta}_x$ for some $k \geq 2$ and some $x \in X$ having $W(x, x) \neq 1$, or $|\text{supp } \mathbf{n}| \geq 2$ and $G_W[\text{supp } \mathbf{n}]$ is connected. In all three cases, $G_W[\text{supp } \mathbf{n}]$ is connected. Furthermore, to make a nonzero contribution to (2.78), $\text{supp } \mathbf{n}$ has to meet both $\text{supp}(\mathbf{1} - \boldsymbol{\lambda}_1)$ and $\text{supp}(\mathbf{1} - \boldsymbol{\lambda}_2)$ and be contained in $\text{supp } \mathbf{R}$. Therefore, there is a component of $G_W[\text{supp } \mathbf{R}]$ (namely, the one containing $\text{supp } \mathbf{n}$) that meets both $\text{supp}(\mathbf{1} - \boldsymbol{\lambda}_1)$ and $\text{supp}(\mathbf{1} - \boldsymbol{\lambda}_2)$. ■

For a repulsive lattice gas with hard-core self-repulsion [i.e. $W(x, x) = 0$ for all $x \in X$], the set $\mathcal{R}(W)$ has an additional convexity property:

Corollary 2.29 (convexity of two-dimensional sections) *For any repulsive lattice gas with hard-core self-repulsion, the two-dimensional sections of $\mathcal{R}(W)$ parallel to the coordinate axes are convex and (when nonempty) are bounded either by a hyperbola or by a straight line.*

PROOF. For $x, y \in X$ ($x \neq y$) and a nonnegative vector $\boldsymbol{\alpha} = (\alpha_z)_{z \in X \setminus \{x, y\}}$, consider the section

$$P_{x, y, \boldsymbol{\alpha}} \mathcal{R}(W) = \{(R_x, R_y) : (R_x, R_y, \boldsymbol{\alpha}) \in \mathcal{R}(W)\}. \quad (2.79)$$

If $(0, 0, \boldsymbol{\alpha}) \notin \mathcal{R}(W)$, then clearly the section is empty; so let us assume henceforth that $(0, 0, \boldsymbol{\alpha}) \in \mathcal{R}(W)$. Let $Z_W(w_x, w_y; -\boldsymbol{\alpha})$ be the polynomial obtained from $Z_W(\mathbf{w})$ by setting $w_z = -\alpha_z$ for $z \neq x, y$. Then, as Z_W is multiaffine, we have

$$Z_W(w_x, w_y; -\boldsymbol{\alpha}) = A + Bw_x + Cw_y + Dw_xw_y \quad (2.80)$$

where

$$A = Z_W(0, 0; -\boldsymbol{\alpha}) \quad (2.81a)$$

$$B = Z_W(0, 0; -W(x, \cdot)\boldsymbol{\alpha}) \quad (2.81b)$$

$$C = Z_W(0, 0; -W(y, \cdot)\boldsymbol{\alpha}) \quad (2.81c)$$

$$D = W(x, y) Z_W(0, 0; -W(x, \cdot)W(y, \cdot)\boldsymbol{\alpha}) \quad (2.81d)$$

as a consequence of the multiple differentiation identity (2.6). Since $(0, 0, \boldsymbol{\alpha}) \in \mathcal{R}(W)$, $0 \leq W(x, y) \leq 1$ for all x, y , and $\mathcal{R}(W)$ is a down-set, we have $A, B, C > 0$ and $D \geq 0$ [with $D = 0$ if and only if $W(x, y) = 0$]. Moreover, by Corollary 2.27(c) we have

$AD \leq BC$.¹⁰ It follows from Theorem 2.10(a) that $P_{x,y,\alpha}\mathcal{R}(W)$ is the component of $[0, \infty)^2 \cap \{(R_x, R_y): Z_W(-R_x, -R_y; -\alpha) > 0\}$ containing $(0, 0)$ [here we have used the fact that $(0, 0, \alpha) \in \mathcal{R}(W)$]. Thus,

$$P_{x,y,\alpha}\mathcal{R}(W) = \{(R_x, R_y): R_x, R_y \geq 0 \text{ and } BR_x + CR_y < A\} \quad (2.82)$$

in case $D = 0$, and

$$P_{x,y,\alpha}\mathcal{R}(W) = \{(R_x, R_y): 0 \leq R_x \leq C/D \text{ and } 0 \leq R_y \leq B/D \text{ and } (C/D - R_x)(B/D - R_y) > (BC - AD)/D^2\} \quad (2.83)$$

in case $D > 0$. ■

Remarks. 1. The proof shows that the boundary of $P_{x,y,\alpha}\mathcal{R}(W)$ is a straight line when $W(x, y) = 0$ and is a hyperbola when $0 < W(x, y) \leq 1$.

2. Consider $\mathbf{w} = (w_x, w_y)$ and $\mathbf{w}' = (w'_x, w'_y)$, and let $0 \leq \lambda \leq 1$; then

$$\begin{aligned} Z_W(\lambda\mathbf{w} + (1-\lambda)\mathbf{w}'; -\alpha) &- [\lambda Z_W(\mathbf{w}; -\alpha) + (1-\lambda)Z_W(\mathbf{w}'; -\alpha)] \\ &= -D\lambda(1-\lambda)(w_x - w'_x)(w_y - w'_y), \end{aligned} \quad (2.84)$$

so that $Z_W(\cdot; -\alpha)$ is concave along line segments of negative slope in the (w_x, w_y) -plane. This provides an alternate proof that the projected section $P_{x,y,\alpha}\mathcal{R}(W)$ is convex [because $\mathcal{R}(W)$ is a down-set, only line segments of negative slope need be considered].

3. The convexity of two-dimensional sections is false without the hypothesis of hard-core self-repulsion: consider $X = \{1, 2\}$, $W(1, 1) = W(2, 2) = 1$ and $W(1, 2) = 0$. Then $Z_W(\mathbf{w}) = e^{w_1} + e^{w_2} - 1$, so that $\mathcal{R}(W) = \{(R_1, R_2): R_1, R_2 \geq 0 \text{ and } e^{-R_1} + e^{-R_2} > 1\}$, which is not convex.

4. Even in the case of hard-core self-repulsion, $\mathcal{R}(W)$ itself is not in general convex. To see this, let $X = \{1, 2, 3\}$ and let G be the path 123; its independent-set polynomial is $Z_G(w) = w_2 + (1 + w_1)(1 + w_3)$. The diagonal section of $\mathcal{R}(G)$ given by $R_1 = R_3 = x$ and $R_2 = y$ is the region of $[0, 1]^2$ where $y < (1 - x)^2$, which is not convex.

2.6 Algebraic irreducibility of $Z_W(\mathbf{w})$

In this subsection — which is a digression from the main thread of the paper and can be omitted on a first reading — we discuss the algebraic irreducibility of the multivariate partition function $Z_W(\mathbf{w})$. We restrict attention to the case of hard-core self-repulsion, in which Z_W is a multi-affine polynomial.

Let R be a commutative ring with identity, and let $W: X \times X \rightarrow R$ be symmetric and satisfy $W(x, x) = 0$ for all $x \in X$. Define the support graph G_W by setting

¹⁰In the case of a hard-core pair interaction, this follows alternatively from Corollary 2.27(d).

$xy \in E(G)$ if and only if $W(x, y) \neq 1$ and $x \neq y$. Define the polynomial $Z_W \in R[\mathbf{w}]$ by

$$Z_W = \sum_{X' \subseteq X} \left(\prod_{\{x, y\} \subseteq X'} W(x, y) \right) \left(\prod_{x \in X'} w_x \right). \quad (2.85)$$

Proposition 2.30 *Suppose that R is an integral domain. Then Z_W is irreducible over R if and only if G_W is connected.*

The proof is based on an easy but crucial lemma [34, Lemma 4.7]:

Lemma 2.31 *Let P_1 and P_2 be nonzero polynomials in the indeterminates $\{w_x\}_{x \in X}$ with coefficients in an integral domain R . Suppose that $P_1 P_2$ is multiaffine (i.e. of degree 0 or 1 in each variable separately). Then:*

- (a) *There exist disjoint subsets $X_1, X_2 \subseteq X$ such that P_i uses only the variables $\{w_x\}_{x \in X_i}$ ($i = 1, 2$).*
- (b) *P_1 and P_2 are both multiaffine.*

PROOF. Suppose there exists $x \in X$ such that both P_1 and P_2 use the variable w_x . For $i = 1, 2$, let $d_i \geq 1$ be the degree of P_i in the variable w_x , and let $Q_i \neq 0$ be the coefficient of $w_x^{d_i}$ in P_i , considered as an element of the polynomial ring $R[\mathbf{w}_{\neq x}]$. Then $Q_1 Q_2 \neq 0$ because $R[\mathbf{w}_{\neq x}]$ is an integral domain [65, Theorem III.5.1 and Corollary III.5.7]. But this shows that the coefficient of $w_x^{d_1 + d_2}$ in $P_1 P_2$ is nonzero, contradicting the hypothesis that $P_1 P_2$ is multiaffine (since $d_1 + d_2 \geq 2$). This proves (a); and (b) is an easy consequence. ■

PROOF OF PROPOSITION 2.30. If G_W is disconnected, it is obvious that Z_W is reducible. To prove the converse, suppose that Z_W is reducible, i.e. $Z_W = P_1 P_2$ where P_1 and P_2 are nonconstant polynomials over R . It follows from Lemma 2.31 that there exist disjoint subsets $X_1, X_2 \subseteq X$ such that P_i uses only the variables $\{w_x\}_{x \in X_i}$ (we can assume without loss of generality that $X_1 \cup X_2 = X$); moreover, neither X_1 nor X_2 can be empty (since P_1 and P_2 are nonconstant). Let $a_i \in R$ be the constant term of P_i ($i = 1, 2$). Since Z_W has constant term 1, we have $a_1 a_2 = 1$. Replacing P_1 by $a_2 P_1$ and P_2 by $a_1 P_2$, we can assume without loss of generality that $a_1 = a_2 = 1$. For $x \in X_1$ (resp. $x \in X_2$), let b_x be the coefficient of the linear term w_x in P_1 (resp. in P_2). Since the linear term w_x has coefficient 1 in Z_W and $a_1 = a_2 = 1$, we have $b_x = 1$ for all $x \in X$. It follows that, for $x \in X_1$ and $y \in X_2$, the term $w_x w_y$ in $P_1 P_2$ has coefficient 1; since this term has coefficient $W(x, y)$ in $Z_W = P_1 P_2$, we have $W(x, y) = 1$ whenever $x \in X_1$ and $y \in X_2$. Hence G_W is disconnected. ■

Remarks. 1. The *univariate* polynomial $Z_W(w)$ is obviously reducible over \mathbb{C} whenever $|X| \geq 2$; and it is in many cases reducible over \mathbb{R} as well. Indeed, it

can be reducible over the integers: e.g. for $G = P_4$ (the 4-vertex path) we have $Z_G(w) = 1 + 4w + 3w^2 = (3w + 1)(w + 1)$.

2. Results analogous to Proposition 2.30 hold for some other combinatorial polynomials. For example, the irreducibility over \mathbb{C} for the (bivariate) Tutte polynomial $Z_M(x, y)$ of a connected matroid M was proven recently by Merino, de Mier and Noy [80]. Likewise, the irreducibility over any integral domain R for the *multivariate* basis generating polynomial of a connected matroid M — or more generally, for any multiaffine polynomial whose support is exactly the collection of bases of M — was proven recently by Choe, Oxley, Sokal and Wagner [34].

3. It would be interesting to know whether analogous results can be obtained without the hypothesis of hard-core self-repulsion. The difficulty is to find the right ring in which to work. Irreducibility in the ring $R[[\mathbf{w}]]$ of formal power series is too trivial, because *every* formal power series with constant term 1 is invertible. On the other hand, it might conceivably be possible to prove irreducibility in the ring of entire functions (or entire functions of order 1) on \mathbb{C}^X — but only for $|X| \geq 2$, since it is manifestly false if $|X| = 1$ and $0 < W(x, x) < 1$ (a nonpolynomial entire function of order 0 has infinitely many zeros [76, Theorem I.13]).

2.7 Convexity of $\log Z$ at nonnegative fugacity

In this paper we are primarily concerned with the behavior of $Z_W(\mathbf{w})$ for *complex* fugacities \mathbf{w} . However, the regime of *nonnegative* fugacities \mathbf{w} is of particular interest to probabilists and statistical mechanicians, since the Boltzmann weights [cf. (1.1)] are there nonnegative and so can be interpreted, after normalization, as a probability measure on lattice-gas configurations. In this “probabilistic regime”, the logarithm of the partition function is, in very great generality, a convex function of all the interaction energies [67, p. 12]. Let us prove the specialization of this statement to our model:

Let $0 \leq W, W' \leq 1$ and $\mathbf{w}, \mathbf{w}' \geq 0$. Then, for $0 \leq \lambda \leq 1$, define $W_\lambda(x, y) = W(x, y)^\lambda W'(x, y)^{1-\lambda}$ and $(w_\lambda)_x = w_x^\lambda (w'_x)^{1-\lambda}$. By Hölder’s inequality applied to (1.1a) or (1.1b), we obtain:

Lemma 2.32 *Suppose that $0 \leq W, W' \leq 1$ and $\mathbf{w}, \mathbf{w}' \geq 0$. Then, for $0 \leq \lambda \leq 1$,*

$$Z_{W_\lambda}(\mathbf{w}_\lambda) \leq Z_W(\mathbf{w})^\lambda Z_{W'}(\mathbf{w}')^{1-\lambda}. \quad (2.86)$$

That is, $\log Z_{W_\lambda}(\mathbf{w}_\lambda)$ is a convex function of λ ; or in other words, $\log Z_W(\mathbf{w})$ is a convex function of the vector $\langle \{\log W(x, y)\}_{x, y \in X}, \{\log w_x\}_{x \in X} \rangle$.

3 The lattice gas with hard-core self-repulsion

We now restrict attention to the case of a repulsive lattice gas with hard-core self-repulsion, i.e. $0 \leq W(x, y) \leq 1$ for all x, y and $W(x, x) = 0$ for all x . This means, as noted in the Introduction, that $Z_W(\mathbf{w})$ can be written as a sum over subsets:

$$Z_W(\mathbf{w}) = \sum_{X' \subseteq X} \prod_{x \in X'} w_x \prod_{\{x, y\} \subseteq X'} W(x, y). \quad (3.1)$$

Here we shall exploit this special structure, which implies that Z_W is a *multiaffine polynomial*, i.e. a polynomial of degree 1 in each w_x separately. Since W will be fixed throughout, we shall henceforth often omit it from the notation.

3.1 The fundamental identity

Let us define, for each subset $\Lambda \subseteq X$, the restricted partition function

$$Z_\Lambda(\mathbf{w}) = \sum_{X' \subseteq \Lambda} \prod_{x \in X'} w_x \prod_{\{x, y\} \subseteq X'} W(x, y). \quad (3.2)$$

Of course this notation is redundant, since the same effect can be obtained by setting $w_x = 0$ for $x \in X \setminus \Lambda$, but it is useful for the purpose of inductive computations and proofs. We have, for any $x \in \Lambda$, the *fundamental identity*

$$Z_\Lambda(\mathbf{w}) = Z_{\Lambda \setminus x}(\mathbf{w}) + w_x Z_{\Lambda \setminus x}(W(x, \cdot)\mathbf{w}) \quad (3.3)$$

where

$$[W(x, \cdot)\mathbf{w}]_y = W(x, y) w_y; \quad (3.4)$$

here the first term on the right-hand side of (3.3) covers the summands in (3.2) with $X' \not\ni x$, while the second covers $X' \ni x$. [Note that (3.3) is a special case of (2.3).] In the special case of a hard-core interaction (= independent-set polynomial) for a graph G , (3.3) reduces to

$$Z_\Lambda(\mathbf{w}) = Z_{\Lambda \setminus x}(\mathbf{w}) + w_x Z_{\Lambda \setminus \Gamma^*(x)}(\mathbf{w}), \quad (3.5)$$

where we have used the notation $\Gamma^*(x) = \Gamma(x) \cup \{x\}$ [here $\Gamma(x)$ denotes the set of vertices of G adjacent to x]. The fundamental identity (3.3)/(3.5) will play an important role both in the inductive proof of the Lovász local lemma [cf. (4.10) and (4.22)] and in the Dobrushin–Shearer inductive argument for the nonvanishing of Z_W in a polydisc (Section 5).

Let us also remark that because $Z_{\Lambda \setminus x}$ is multiaffine, the last term in (3.3) can be expanded to rewrite $Z_{\Lambda \setminus x}(W(x, \cdot)\mathbf{w})$ as a linear combination of values at the vertices

of a rectangle:¹¹

$$Z_\Lambda(\mathbf{w}) = Z_{\Lambda \setminus x}(\mathbf{w}) + w_x \sum_{Y \subseteq \Lambda \setminus x} \left(\prod_{y \in Y} W(x, y) \right) \left(\prod_{y \in (\Lambda \setminus x) \setminus Y} [1 - W(x, y)] \right) Z_Y(\mathbf{w}). \quad (3.6)$$

In the hard-core case, the only term with a nonzero coefficient is $Y = \Lambda \setminus \Gamma^*(x)$, so that (3.6) reduces to (3.5).

Remark. Repeated use of (3.3) obviously gives an algorithm to compute $Z_W(\mathbf{w})$. But this algorithm takes in general exponential time. In fact, calculating $Z_G(\mathbf{w})$ for general graphs G (or even for cubic planar graphs) is NP-hard (as noted by Shearer [98]), since even calculating the *degree* of $Z_G(\mathbf{w})$ — that is, the maximum size of an independent set — is NP-hard [44, pp. 194–195]. Therefore, if $P \neq NP$ it is impossible to calculate $Z_G(\mathbf{w})$ for general graphs in polynomial time.

A key role in (3.3) is manifestly played by the rational function

$$\mathcal{K}_{x,\Lambda}(\mathbf{w}) \equiv \frac{Z_{\Lambda \setminus x}(W(x, \cdot)\mathbf{w})}{Z_{\Lambda \setminus x}(\mathbf{w})}. \quad (3.7)$$

Note that $\mathcal{K}_{x,\Lambda}(\mathbf{w})$ depends only on $\{w_y\}_{y \in \Lambda \setminus x}$; in particular, it does not depend on w_x . We shall sometimes write it as $\mathcal{K}_{x,\Lambda}(\mathbf{w}_{\neq x})$ to emphasize this fact. Please note also that $Z_\Lambda(\mathbf{w})$ is nonvanishing in the region $|w_x| < 1/|\mathcal{K}_{x,\Lambda}(\mathbf{w}_{\neq x})|$ and vanishes at $w_x = -1/\mathcal{K}_{x,\Lambda}(\mathbf{w}_{\neq x})$. We have

$$\frac{\partial \log Z_\Lambda(\mathbf{w})}{\partial w_x} = \frac{\mathcal{K}_{x,\Lambda}(\mathbf{w}_{\neq x})}{1 + \mathcal{K}_{x,\Lambda}(\mathbf{w}_{\neq x})w_x}. \quad (3.8)$$

Since $\mathcal{K}_{x,\Lambda}(\mathbf{w})$ is a special case of the function $Y_W(\mathbf{w}; \boldsymbol{\lambda})$ defined in (2.70), we obtain as an immediate corollary of Proposition 2.26(b):

Proposition 3.1 *Consider any repulsive lattice gas with hard-core self-repulsion. If $\mathbf{R} \in \mathcal{R}(W)$, then*

$$|\mathcal{K}_{x,\Lambda}(\mathbf{w})| \leq \mathcal{K}_{x,\Lambda}(-\mathbf{R}) < \infty \quad (3.9)$$

for all $\mathbf{w} \in \bar{D}_{\mathbf{R}}$. That is, the maximum of $|\mathcal{K}_{x,\Lambda}(\mathbf{w})|$ over $\bar{D}_{\mathbf{R}}$ is attained at $\mathbf{w} = -\mathbf{R}$, and this maximum value $\mathcal{K}_{x,\Lambda}(-\mathbf{R})$ is an increasing function of $\mathbf{R} \in \mathcal{R}(W)$.

¹¹If $F: \mathbb{C}^X \rightarrow \mathbb{C}$ is multiaffine, then F can be reconstructed from its values at the corners of any rectangle with positive volume. For example, if the rectangle is $[0, 1]^X$, we have

$$F(\mathbf{z}) = \sum_{Y \subseteq X} \left(\prod_{i \in Y} z_i \right) \left(\prod_{i \in X \setminus Y} (1 - z_i) \right) F(\mathbf{1}_Y).$$

This is clear when $\mathbf{z} \in \{0, 1\}^X$, and equality for all \mathbf{z} then follows because both sides are multiaffine.

The function $\mathcal{K}_{x,\Lambda}(-\mathbf{R})$ will play an important role in Section 5.

We can translate the fundamental identity (3.3) into a recursion for the rational functions $\mathcal{K}_{x,\Lambda}(\mathbf{w})$. Let us order arbitrarily the sites $\Lambda \setminus x = \{y_1, \dots, y_k\}$. Then we write (3.7) as a telescoping product:

$$\mathcal{K}_{x,\Lambda}(\mathbf{w}) = \prod_{i=1}^k \frac{Z_{\Lambda \setminus x}(\tilde{\mathbf{w}}^{(i)})}{Z_{\Lambda \setminus x}(\tilde{\mathbf{w}}^{(i-1)})} \quad (3.10)$$

where the vectors $\tilde{\mathbf{w}}^{(i)}$ are defined by

$$(\tilde{\mathbf{w}}^{(i)})_y = \begin{cases} W(x, y)w_y & \text{if } y = y_j \text{ for some } j \leq i \\ w_y & \text{otherwise} \end{cases} \quad (3.11)$$

[Note that it is irrelevant how we define $(\tilde{\mathbf{w}}^{(i)})_y$ for $y = x$, since we set this weight to zero anyway by considering $Z_{\Lambda \setminus x}$.] Applying the fundamental identity (3.3) to y_i , we obtain

$$\frac{Z_{\Lambda \setminus x}(\tilde{\mathbf{w}}^{(i)})}{Z_{\Lambda \setminus x}(\tilde{\mathbf{w}}^{(i-1)})} = \frac{1 + W(x, y_i)w_{y_i}\mathcal{K}_{y_i, \Lambda \setminus x}(\tilde{\mathbf{w}}^{(i-1)})}{1 + w_{y_i}\mathcal{K}_{y_i, \Lambda \setminus x}(\tilde{\mathbf{w}}^{(i-1)})} \quad (3.12)$$

and hence

$$\mathcal{K}_{x,\Lambda}(\mathbf{w}) = \prod_{i=1}^k \frac{1 + W(x, y_i)w_{y_i}\mathcal{K}_{y_i, \Lambda \setminus x}(\tilde{\mathbf{w}}^{(i-1)})}{1 + w_{y_i}\mathcal{K}_{y_i, \Lambda \setminus x}(\tilde{\mathbf{w}}^{(i-1)})}. \quad (3.13)$$

This identity will play a central role at the end of Section 5. In the special case of a hard-core pair interaction, (3.13) becomes

$$\mathcal{K}_{x,\Lambda}(\mathbf{w}) = \prod_{i=1}^l \frac{1}{1 + w_{y_i}\mathcal{K}_{y_i, \Lambda \setminus \{y_1, \dots, y_{i-1}\}}(\mathbf{w})} \quad (3.14)$$

where $\{y_1, \dots, y_l\}$ is an ordering of $\Lambda \cap \Gamma(x)$.

Note, finally, that the partition functions $Z_\Lambda(\mathbf{w})$ can be reconstructed from the rational functions $\mathcal{K}_{x,\Lambda}(\mathbf{w})$. For, by the fundamental identity (3.3) and the definition (3.7), we have

$$\frac{Z_\Lambda(\mathbf{w})}{Z_{\Lambda \setminus x}(\mathbf{w})} = 1 + w_x \mathcal{K}_{x,\Lambda}(\mathbf{w}_{\neq x}); \quad (3.15)$$

and hence, setting $\Lambda = \{x_1, \dots, x_n\}$ (in arbitrary order) we can write

$$Z_\Lambda(\mathbf{w}) = \prod_{i=1}^n \frac{Z_{\{x_1, \dots, x_i\}}(\mathbf{w})}{Z_{\{x_1, \dots, x_{i-1}\}}(\mathbf{w})} = \prod_{i=1}^n [1 + w_{x_i} \mathcal{K}_{x_i, \{x_1, \dots, x_i\}}(\mathbf{w})]. \quad (3.16)$$

3.2 Examples

Let us take a moment to compute a few examples of partition functions $Z_W(\mathbf{w})$. We shall assume hard-core self-repulsion [i.e. $W(x, x) = 0$ for all x] throughout, and shall mostly restrict attention to the case of a hard-core pair interaction, i.e.

the independent-set polynomial $Z_G(\mathbf{w})$ for a graph G . In some cases we shall, for simplicity, compute only the univariate polynomial $Z_G(w)$ obtained by setting $w_x = w$ for all vertices x .

Example 3.1. *The complete graph K_n .* Clearly $Z_{K_n}(\mathbf{w}) = 1 + w_1 + \dots + w_n$. In particular, $\mathcal{R}(K_n) = \{\mathbf{R}: R_1 + \dots + R_n < 1\}$.

Example 3.2. *The n -vertex path P_n .* We limit attention to the univariate independent-set polynomial. Applying the fundamental identity (3.5) to an end-vertex, we obtain the recursion relation

$$Z_{P_n}(w) = Z_{P_{n-1}}(w) + wZ_{P_{n-2}}(w), \quad (3.17)$$

which is valid for all $n \geq 0$ if we define $Z_{P_0} \equiv Z_{P_{-1}} \equiv 1$ and $Z_{P_{-2}} \equiv 0$. The solution of (3.17) is

$$Z_{P_n}(w) = \frac{1}{\sqrt{1+4w}} (\lambda_+^{n+2} - \lambda_-^{n+2}) \quad (3.18)$$

where

$$\lambda_{\pm} = \frac{1 \pm \sqrt{1+4w}}{2}. \quad (3.19)$$

(An alternative method of obtaining this result is given in Example 3.2' below.) The zeros of Z_{P_n} are located at

$$w = -\frac{1}{4 \cos^2 \frac{\pi k}{n+2}} \quad (3.20)$$

for $k = 1, 2, \dots, \lfloor \frac{n+1}{2} \rfloor$. The zero nearest the origin converges from below to $w = -1/4$ as $n \rightarrow \infty$. Elementary counting arguments give also the explicit formula

$$Z_{P_n}(w) = \sum_{k=0}^{\lfloor \frac{n+1}{2} \rfloor} \binom{n+1-k}{k} w^k. \quad (3.21)$$

Example 3.3. *The n -vertex cycle C_n .* We again limit attention to the univariate independent-set polynomial. Applying the fundamental identity (3.5), we find

$$Z_{C_n}(w) = Z_{P_{n-1}}(w) + wZ_{P_{n-3}}(w), \quad (3.22)$$

valid for $n \geq 2$. Inserting (3.18)/(3.19), we obtain

$$Z_{C_n}(w) = \lambda_+^n + \lambda_-^n. \quad (3.23)$$

(An alternative method of obtaining this result is given in Example 3.3' below.) The zeros of Z_{C_n} are located at

$$w = -\frac{1}{4 \cos^2 \frac{\pi(k+\frac{1}{2})}{n}} \quad (3.24)$$

for $k = 0, 1, \dots, \lfloor \frac{n}{2} \rfloor - 1$. The zero nearest the origin converges from below to $w = -1/4$ as $n \rightarrow \infty$. Elementary counting arguments [or (3.21)/(3.22)] give also the explicit formula

$$Z_{C_n}(w) = \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \frac{n}{n-k} \binom{n-k}{k} w^k. \quad (3.25)$$

Remark. There is a one-to-one correspondence between matchings on a graph G and independent sets on the line graph $L(G)$. Since $L(P_n) = P_{n-1}$ and $L(C_n) = C_n$, these formulae for Z_{P_n} and Z_{C_n} can also be found in the work of Heilmann and Lieb on matching polynomials [58, pp. 196–197]. Heilmann and Lieb also noted that Z_{P_n} and Z_{C_n} can be written as Chebyshev polynomials.

Example 3.3'. *The “soft” n -vertex cycle C_n .* Let us now generalize Example 3.3 by considering the cycle C_n with edge weights

$$W(x, y) = \begin{cases} W & \text{if } xy \text{ is an edge of } C_n \\ 1 & \text{otherwise} \end{cases} \quad (3.26)$$

where W is a constant in $[0, 1]$ (the case $W = 0$ corresponds to the independent-set polynomial of C_n , while the case $W = 1$ corresponds to the independent-set polynomial of the edgeless graph on n vertices). Then the univariate polynomial $Z_W(w)$ is given by a “transfer matrix” [7, Section 2.1] [14, Sections 2.2–2.4] [107, Section 4.7]:

$$Z_W(w) = \text{tr} \begin{pmatrix} 1 & 1 \\ w & wW \end{pmatrix}^n \quad (3.27)$$

where the first (resp. second) row or column corresponds to an empty (resp. occupied) site, and the fugacity w is attributed to the row only. It follows that

$$Z_W(w) = \lambda_+^n + \lambda_-^n, \quad (3.28)$$

where

$$\lambda_{\pm} = \frac{1 + wW \pm \sqrt{(1 - wW)^2 + 4w}}{2} \quad (3.29)$$

are the eigenvalues of the transfer matrix. Simple algebra then shows that all of the zeros of Z_W are real and negative (when $0 \leq W \leq 1$).

Example 3.2'. *The “soft” n -vertex path P_n .* Let us now generalize Example 3.2 analogously. The univariate polynomial $Z_W(w)$ is again given by a transfer matrix:

$$Z_W(w) = (1 \ 1) \begin{pmatrix} 1 & 1 \\ w & wW \end{pmatrix}^{n-1} \begin{pmatrix} 1 \\ w \end{pmatrix}. \quad (3.30)$$

After some algebra we find

$$Z_W(w) = A_+ \lambda_+^{n+2} + A_- \lambda_-^{n+2}, \quad (3.31)$$

with λ_{\pm} given by (3.29) and

$$A_{\pm} = \frac{2}{\pm(2 - W + wW^2)\sqrt{(1 - wW)^2 + 4w} + W[(1 - wW)^2 + 4w]} . \quad (3.32)$$

Example 3.4. *The star $K_{1,r}$.* Let x be the center vertex of the star and let y_1, \dots, y_r be the leaves. The independent-set polynomial is

$$Z_{K_{1,r}}(\mathbf{w}) = \prod_{i=1}^r (1 + w_{y_i}) + w_x . \quad (3.33)$$

More generally, suppose we have a pair interaction matrix $W(x, y_i)$ [with $W(y_i, y_j) = 1$ for all $i \neq j$ and $W(x, x) = W(y_i, y_i) = 0$]. Then

$$Z_W(\mathbf{w}) = \prod_{i=1}^r (1 + w_{y_i}) + w_x \prod_{i=1}^r [1 + W(x, y_i)w_{y_i}] . \quad (3.34)$$

Example 3.5. *The complete bipartite graph $K_{m,n}$.* Let $X = \{x_1, \dots, x_m\}$ and $Y = \{y_1, \dots, y_n\}$ be the bipartition. Then the independent-set polynomial is

$$Z_{K_{m,n}}(\mathbf{w}) = \prod_{i=1}^m (1 + w_{x_i}) + \prod_{j=1}^n (1 + w_{y_j}) - 1 . \quad (3.35)$$

Example 3.6. *The complete r -ary rooted tree [94, 98].* Let $T_n^{(r)}$ be the complete rooted tree with branching factor r and depth n . We limit attention to the univariate independent-set polynomial. Fix $r \geq 1$; and to lighten the notation, let us write Z_n as a shorthand for $Z_{T_n^{(r)}}$. Applying the fundamental identity (3.5) to the root vertex, we obtain the nonlinear recursion

$$Z_n(w) = Z_{n-1}(w)^r + wZ_{n-2}(w)^{r^2} , \quad (3.36)$$

which is valid for all $n \geq 0$ if we set $Z_{-1} \equiv Z_{-2} \equiv 1$. By defining

$$Y_n(w) = \frac{Z_n(w)}{Z_{n-1}(w)^r} , \quad (3.37)$$

we can convert the second-order recursion (3.36) to a first-order recursion

$$Y_n(w) = 1 + \frac{w}{Y_{n-1}(w)^r} \quad (3.38)$$

with initial condition $Y_{-1} \equiv 1$. The polynomials $Z_n(w)$ can be reconstructed from the rational functions $Y_n(w)$ by

$$Z_n(w) = \prod_{k=0}^n Y_k(w)^{r^{n-k}} . \quad (3.39)$$

Let $w_n < 0$ be the negative real root of Z_n of smallest magnitude (set $w_n = -\infty$ if Z_n has no negative real root). Note that $w_{-1} = -\infty$ and $w_0 = -1$. Let us prove by induction that $w_{n-1} < w_n$ for $n \geq 0$. It is true for $n = 0$. For $n \geq 1$ we have

$$Z_n(w_{n-1}) = Z_{n-1}(w_{n-1})^r + w_{n-1}Z_{n-2}(w_{n-1})^{r^2} < 0 \quad (3.40)$$

since $Z_{n-1}(w_{n-1}) = 0$, $w_{n-1} < 0$ and $Z_{n-2}(w_{n-1}) > 0$ by the inductive hypothesis. Therefore Z_n vanishes somewhere between w_{n-1} and 0.

It follows that the w_n increase to a limit $w_\infty \leq 0$ as $n \rightarrow \infty$. Let us show, following Shearer [98], that

$$w_\infty = -\frac{r^r}{(r+1)^{r+1}} \quad (3.41)$$

by proving the two inequalities:

PROOF OF \geq : If $w \in [w_\infty, 0)$, we have $Z_n(w) > 0$ for all n and hence also $Y_n(w) > 0$ for all n . Since $Y_{-1} > Y_0$, it follows from the monotonicity of (3.38) that $\{Y_n(w)\}_{n \geq 0}$ is a strictly decreasing sequence of positive numbers, hence converges to a limit $y_* \geq 0$ satisfying the fixed-point equation $y_* = 1 + w/y_*^r$, or equivalently $w = y_*^{r+1} - y_*^r$. Elementary calculus then shows that $w \geq -r^r/(r+1)^{r+1}$; taking $w = w_\infty$ we obtain $w_\infty \geq -r^r/(r+1)^{r+1}$.

PROOF OF \leq : If $-r^r/(r+1)^{r+1} \leq w < 0$, the equation $w = y_*^{r+1} - y_*^r$ has a unique solution $y_* \in [r/(r+1), 1)$. It then follows by induction [using (3.38) and the initial condition $Y_{-1} = 1$] that $1 = Y_{-1}(w) > Y_0(w) > \dots > Y_{n-1}(w) > Y_n(w) > \dots > y_*$ for all $n \geq 0$. In particular, $Y_n(w) > 0$ for all n , so that $w_n < w$ for all n . This shows that $w_\infty \leq -r^r/(r+1)^{r+1}$.

Let us conclude by observing that (3.38) defines a degree- r rational map $R_w: y \mapsto 1 + w/y^r$ parametrized by $w \in \mathbb{C} \setminus 0$. Moreover, the zeros of $Z_W(w)$ correspond to those values w for which R_w has a (superattractive) orbit $0 \mapsto \infty \mapsto 1 \mapsto 1 + w \mapsto \dots \mapsto 0$ of period $n+3$ (or some submultiple of $n+3$). As $n \rightarrow \infty$, these points accumulate on a ‘‘Mandelbrot-like’’ set in the complex w -plane. For further information on the maps $y \mapsto 1 + w/y^r$, see [81, 5, 18, 19, 20, 21].

3.3 Reduction formulae

Recall that, given W , we have defined a simple loopless graph $G = G_W$ (‘‘the support graph of W ’’) by setting $xy \in E(G)$ if and only if $W(x, y) \neq 1$ and $x \neq y$. When the support graph G has a very simple structure, the polynomial $Z_W(\mathbf{w})$ can be simplified:

1) *Disconnected graph.* Suppose that G can be written as the disjoint union of G_1 and G_2 . Set $\Lambda_i = V(G_i)$. Then clearly Z_W factorizes:

$$Z_{\Lambda_1 \cup \Lambda_2}(\mathbf{w}) = Z_{\Lambda_1}(\mathbf{w}) Z_{\Lambda_2}(\mathbf{w}). \quad (3.42)$$

Note that $Z_{\Lambda_i}(\mathbf{w})$ depends only on $\{w_x\}_{x \in \Lambda_i}$.

2) *Cut vertex.* Suppose that $G = G_1 \cup G_2$ where $V(G_1) \cap V(G_2) = \{x\}$. [If G_1 and G_2 have at least two vertices, this means that x is a cut vertex of G . But the

formulae below hold also in the trivial cases $V(G_1) = \{x\}$ or $V(G_2) = \{x\}$ or both.] Again set $\Lambda_i = V(G_i)$. The fundamental identity (3.3) asserts that

$$Z_{\Lambda_1 \cup \Lambda_2}(\mathbf{w}) = Z_{(\Lambda_1 \cup \Lambda_2) \setminus x}(\mathbf{w}) + w_x Z_{(\Lambda_1 \cup \Lambda_2) \setminus x}(W(x, \cdot) \mathbf{w}). \quad (3.43)$$

But since the graphs $G_1 \setminus x$ and $G_2 \setminus x$ are disjoint¹², we can apply the factorization (3.42) to the right-hand side of (3.43), yielding

$$Z_{\Lambda_1 \cup \Lambda_2}(\mathbf{w}) = Z_{\Lambda_1 \setminus x}(\mathbf{w}) Z_{\Lambda_2 \setminus x}(\mathbf{w}) + w_x Z_{\Lambda_1 \setminus x}(W(x, \cdot) \mathbf{w}) Z_{\Lambda_2 \setminus x}(W(x, \cdot) \mathbf{w}) \quad (3.44a)$$

$$= Z_{\Lambda_1 \setminus x}(\mathbf{w}) [Z_{\Lambda_2 \setminus x}(\mathbf{w}) + w_x^{\text{eff}, G_1} Z_{\Lambda_2 \setminus x}(W(x, \cdot) \mathbf{w})] \quad (3.44b)$$

$$= Z_{\Lambda_1 \setminus x}(\mathbf{w}) Z_{\Lambda_2}(\mathbf{w}_{\neq x}, w_x^{\text{eff}, G_1}) \quad (3.44c)$$

where we have defined

$$w_x^{\text{eff}, G_1} = w_x \frac{Z_{\Lambda_1 \setminus x}(W(x, \cdot) \mathbf{w})}{Z_{\Lambda_1 \setminus x}(\mathbf{w})} = w_x \mathcal{K}_{x, \Lambda_1}(\mathbf{w}). \quad (3.45)$$

This can be interpreted as “integrating out” the variables in $V(G_1) \setminus x$, leaving an “effective fugacity” w_x^{eff, G_1} for the vertex $x \in V(G_2)$.

Example 3.4 revisited. Integrating out the leaves y_1, \dots, y_r of the star $K_{1,r}$, we find from (3.34):

$$w_x^{\text{eff}, K_{1,r}} = w_x \prod_{i=1}^r \frac{1 + W(x, y_i) w_{y_i}}{1 + w_{y_i}}. \quad (3.46)$$

3.4 When G is a tree ...

When the support graph $G = G_W$ is a tree, every vertex of G is either a cut vertex or a leaf, so the partition function $Z_W(\mathbf{w})$ can be calculated by repeated use of the reduction formula (3.44)/(3.45). This can be done in many ways, but the simplest is probably to “roll up” the tree “from the leaves up”, as follows:

We say that a vertex $x \in V(G)$ is a *near-leaf* of G if it is adjacent to at least one leaf of G . If x is a near-leaf of G , then we can write $G = G_1 \cup G_2$ where $V(G_1) \cap V(G_2) = \{x\}$, G_1 is a star with center x consisting of x and some or all of the leaves of G adjacent to x , and G_2 is a tree. Integrating out $G_1 \setminus x$ using (3.46), we obtain

$$w_x^{\text{eff}, G_1} = w_x \prod_{y \in V(G_1 \setminus x)} \frac{1 + W(x, y) w_y}{1 + w_y} \quad (3.47)$$

and of course

$$Z_{\Lambda_1 \setminus x}(\mathbf{w}) = \prod_{y \in V(G_1 \setminus x)} (1 + w_y). \quad (3.48)$$

¹²We recall that $G \setminus x$ denotes the graph obtained from G by deleting the vertex x and all edges incident with it.

Applying this process repeatedly, we can obtain a linear-time algorithm for evaluating $Z_W(\mathbf{w})$ at any point $\mathbf{w} \in \mathbb{C}^X$ whenever the support graph G_W is a tree. (It is also an algorithm for computing $Z_W(\mathbf{w})$ as a polynomial in \mathbf{w} , but in this context it may no longer be linear-time, because of the need to multiply rational functions of \mathbf{w} .) The simplest approach is to orient the tree by choosing arbitrarily one vertex of the tree as the root, and then to work upwards from the leaves. We obtain the following algorithm:

Algorithm T. Given numbers $\{w_x\}_{x \in X}$ and $\{W(x, y)\}_{x, y \in X}$ for which the support graph $G = G_W$ is a tree, we compute $Z_W(\mathbf{w})$ as follows:

- 1) Pick (arbitrarily) a root $x_0 \in X$. Define

$$\begin{aligned} \text{depth}(x) &= \text{dist}(x, x_0) \\ D &= \max_{x \in X} \text{depth}(x) \\ \text{children}(x) &= \{y: xy \in E(G) \text{ and } \text{depth}(y) = \text{depth}(x) + 1\} \end{aligned}$$

- 2) For $d = D, D - 1, \dots, 0$, do:

For all vertices x of depth d , set

$$w_x^{\text{eff}} \longleftarrow w_x \prod_{y \in \text{children}(x)} \frac{1 + W(x, y)w_y^{\text{eff}}}{1 + w_y^{\text{eff}}}. \quad (3.49)$$

[Note that all the needed w_y^{eff} have been set at the previous iteration. If $w_y^{\text{eff}} = -1$ for any child y of x , the algorithm is declared to fail.]

- 3) Output

$$Z_W(\mathbf{w}) = \prod_{x \in V(G)} (1 + w_x^{\text{eff}}). \quad (3.50)$$

The correctness of this algorithm (when it succeeds) follows from (3.44b), (3.44c) and (3.48). Indeed, for any vertex x , the w_x^{eff} produced by this algorithm is w_x^{eff, G_x} where G_x is the subtree consisting of x and all its descendants.

This also provides an algorithm for testing whether a given vector \mathbf{R} lies in $\mathcal{R}(W)$:

Theorem 3.2 *Consider any repulsive lattice gas with hard-core self-repulsion for which the support graph G_W is a tree. Then a vector $\mathbf{R} \geq 0$ lies in $\mathcal{R}(W)$ if and only if Algorithm T with $\mathbf{w} = -\mathbf{R}$ produces $-1 < w_x^{\text{eff}} \leq 0$ for all vertices x .*

PROOF. If Algorithm T with $\mathbf{w} = -\mathbf{R}$ produces $-1 < w_x^{\text{eff}} \leq 0$ for all x , then by the monotonicity of (3.49) in each w_y^{eff} it also does so when $\mathbf{w} = -\mathbf{R}'$ for any vector $0 \leq \mathbf{R}' \leq \mathbf{R}$. By (3.50) this means that $Z_W(-\mathbf{R}') > 0$ for all such \mathbf{R}' , hence that $\mathbf{R} \in \mathcal{R}(W)$.

Conversely, let x be a vertex such that $w_x^{\text{eff}} \notin (-1, 0]$ but $w_y^{\text{eff}} \in (-1, 0]$ for all descendants y of x . Since $w_y^{\text{eff}} \in (-1, 0]$ for all children y of x , and $w_x = -R_x \leq 0$, it follows from (3.49) that $w_x^{\text{eff}} \leq 0$, so we must have $w_x^{\text{eff}} \leq -1$. Now set $R'_z = R_z$ for z in the subtree consisting of x and its descendants, and $R'_z = 0$ otherwise. Applying Algorithm T to $\mathbf{w}' = -\mathbf{R}'$, it follows from (3.50) that $Z_W(\mathbf{w}') \leq 0$. Thus $\mathbf{R}' \notin \mathcal{R}(W)$; and since $\mathcal{R}(W)$ is a down-set, we have $\mathbf{R} \notin \mathcal{R}(W)$. ■

Example 3.6 revisited. Consider again the tree $T_n^{(r)}$. Comparison of (3.38) with (3.49) shows that in the multivariate case $w_x^{\text{eff}} = Y_n(w) - 1$ for all vertices x of height n above the leaves. Then (3.39) is equivalent to (3.50).

3.5 Upper bounds on $\mathcal{R}(W)$ when G_W is a tree

We can use Theorem 3.2 to prove upper bounds on the set $\mathcal{R}(W)$ whenever the support graph G_W is a tree. Let us begin by considering the special case of a hard-core pair interaction, i.e. the independent-set polynomial for a tree G on the vertex set X . As before, pick (arbitrarily) a root $x_0 \in X$, and define $X_i = \{x \in X : \text{depth}(x) = i\}$ for $i = 0, \dots, D$. Given any vector $\mathbf{R} \geq 0$, define \tilde{R}_i to be the geometric mean of R_x over all vertices x of depth i :

$$\tilde{R}_i = \left(\prod_{x \in X_i} R_x \right)^{1/|X_i|}. \quad (3.51)$$

Now apply Algorithm T to $\mathbf{w} = -\mathbf{R}$, let $p_x = -w_x^{\text{eff}}$, and define

$$\tilde{p}_i = \left(\prod_{x \in X_i} p_x \right)^{1/|X_i|} \quad (3.52)$$

(we assume here that $w_x^{\text{eff}} \leq 0$ for all $x \in X_i$). By (3.49) we have

$$p_x = \frac{R_x}{\prod_{y \in \text{children}(x)} (1 - p_y)}, \quad (3.53)$$

so that

$$\tilde{p}_i = \frac{\tilde{R}_i}{\left(\prod_{y \in X_{i+1}} (1 - p_y) \right)^{1/|X_i|}}. \quad (3.54)$$

Now

$$\left(\prod_{y \in X_{i+1}} (1 - p_y) \right)^{1/|X_{i+1}|} \leq 1 - \frac{1}{|X_{i+1}|} \sum_{y \in X_{i+1}} p_y \quad (3.55a)$$

$$\leq 1 - \left(\prod_{y \in X_{i+1}} p_y \right)^{1/|X_{i+1}|} \quad (3.55b)$$

by two applications of the arithmetic-geometric mean inequality. It follows that

$$\tilde{p}_i \geq \frac{\tilde{R}_i}{(1 - \tilde{p}_{i+1})^{|X_{i+1}|/|X_i|}} . \quad (3.56)$$

So, suppose we fix the numbers $(\tilde{R}_i)_{i=0}^D$ and define $(\hat{p}_i)_{i=0}^D$ by the recursion

$$\hat{p}_i = \frac{\tilde{R}_i}{(1 - \hat{p}_{i+1})^{|X_{i+1}|/|X_i|}} \quad (3.57)$$

with initial condition $\hat{p}_{D+1} = 0$ (which is well-defined as long as \hat{p}_{i+1} remains < 1). Then it follows immediately from the monotonicity of the right-hand side of (3.56) that $\tilde{p}_i \geq \hat{p}_i$ as long as the former is well-defined. In particular, if for some level i we have $\hat{p}_i \geq 1$, then \tilde{p}_i is either ≥ 1 or else ill-defined, so that there exists $x \in X_i$ with $w_x^{\text{eff}} \notin (-1, 0]$. It then follows from Theorem 3.2 that $\mathbf{R} \notin \mathcal{R}(W)$. Moreover, since this calculation depends only on the numbers $(\tilde{R}_i)_{i=0}^D$, the same conclusion holds for *any* \mathbf{R} with the given geometric means on levels.

Let us now generalize this argument to the case of a soft-core pair interaction $0 \leq W(x, y) \leq 1$. It is easily verified that $1 - Wp \geq (1 - p)^W$ whenever $0 \leq p \leq 1$ and $0 \leq W \leq 1$. Therefore, defining $p_x = -w_x^{\text{eff}}$ as before and applying Algorithm T to $\mathbf{w} = -\mathbf{R}$, we have

$$p_x \geq \frac{R_x}{\prod_{y \in \text{children}(x)} (1 - p_y)^{1 - W(x, y)}} . \quad (3.58)$$

Now let us assign to each vertex x a weight

$$\alpha_x = \prod_{j=1}^d [1 - W(x_{j-1}, x_j)] \quad (3.59)$$

where $x_0, x_1, \dots, x_d \equiv x$ is the unique path in G_W connecting x to the root x_0 . [In particular, $\alpha_{x_0} = 1$ and if y is a child of x , then $\alpha_y = [1 - W(x, y)]\alpha_x$.] Setting

$$\tilde{\alpha}_i = \sum_{x \in X_i} \alpha_x , \quad (3.60)$$

let us define \tilde{R}_i and \tilde{p}_i to be the weighted geometric means

$$\tilde{R}_i = \left(\prod_{x \in X_i} R_x^{\alpha_x} \right)^{1/\tilde{\alpha}_i} \quad (3.61)$$

$$\tilde{p}_i = \left(\prod_{x \in X_i} p_x^{\alpha_x} \right)^{1/\tilde{\alpha}_i} \quad (3.62)$$

The preceding argument can then be repeated verbatim (using the weighted arithmetic-geometric mean inequality), yielding

$$\tilde{p}_i \geq \frac{\tilde{R}_i}{(1 - \tilde{p}_{i+1})^{\tilde{\alpha}_{i+1}/\tilde{\alpha}_i}}, \quad (3.63)$$

which can be analyzed analogously to (3.56). We have therefore proven:

Proposition 3.3 *Consider any repulsive lattice gas with hard-core self-repulsion for which the support graph G_W is a tree. Define weights $(\alpha_x)_{x \in X}$ and $(\tilde{\alpha}_i)_{i=0}^D$ by (3.59)/(3.60), and define numbers $(\hat{p}_i)_{i=0}^D$ by the recursion*

$$\hat{p}_i = \frac{\tilde{R}_i}{(1 - \hat{p}_{i+1})^{\tilde{\alpha}_{i+1}/\tilde{\alpha}_i}} \quad (3.64)$$

(as long as \hat{p}_{i+1} remains < 1), with initial condition $\hat{p}_{D+1} = 0$. Suppose that there exists an i for which $\hat{p}_i \geq 1$. Then $\mathbf{R} \notin \mathcal{R}(W)$ for all vectors $\mathbf{R} \geq 0$ satisfying $\prod_{x \in X_i} R_x^{\alpha_x/\tilde{\alpha}_i} \geq \tilde{R}_i$ for all i .

Just as Proposition 3.3 “homogenizes” each level of a tree, we can also “homogenize” *between* levels. Let us again begin by considering the special case of a hard-core pair interaction. We first define branching factors $b_i = |X_{i+1}|/|X_i|$. Choose any number $\bar{b} > 0$ and define

$$\gamma_i = |X_i|/\bar{b}^i. \quad (3.65)$$

Fix an integer $k \geq 1$ (the number of levels to be averaged together) and define

$$\gamma_{i,k} = \sum_{j=i}^{i+k-1} \gamma_j. \quad (3.66)$$

Then define the weighted inter-level geometric means

$$\tilde{p}_{i,k} = \left(\prod_{j=i}^{i+k-1} \tilde{p}_j^{\gamma_j} \right)^{1/\gamma_{i,k}} \quad (3.67)$$

$$\tilde{R}_{i,k} = \left(\prod_{j=i}^{i+k-1} \tilde{R}_j^{\gamma_j} \right)^{1/\gamma_{i,k}} \quad (3.68)$$

for $i = 0, \dots, D+1-k$, and $\tilde{p}_{i,k} = \tilde{R}_{i,k} = 0$ for $i \geq D+2-k$. It follows from (3.56) that

$$\begin{aligned} \tilde{p}_{i,k} &\geq \frac{\tilde{R}_{i,k}}{\prod_{j=i}^{i+k-1} (1 - \tilde{p}_{j+1})^{b_j \gamma_j / \gamma_{i,k}}} \\ &= \frac{\tilde{R}_{i,k}}{\prod_{j=i}^{i+k-1} (1 - \tilde{p}_{j+1})^{\bar{b} \gamma_{j+1} / \gamma_{i,k}}}. \end{aligned} \quad (3.69)$$

Now

$$\begin{aligned}
\prod_{j=i}^{i+k-1} (1 - \tilde{p}_{j+1})^{\gamma_{j+1}/\gamma_{i+1,k}} &\leq 1 - \sum_{j=i}^{i+k-1} \frac{\gamma_{j+1}}{\gamma_{i+1,k}} \tilde{p}_{j+1} \\
&\leq 1 - \prod_{j=i}^{i+k-1} \tilde{p}_{j+1}^{\gamma_{j+1}/\gamma_{i+1,k}} \\
&\equiv 1 - \tilde{\tilde{p}}_{i+1}
\end{aligned} \tag{3.70}$$

by two applications of the arithmetic-geometric mean inequality, so that

$$\tilde{p}_{i,k} \geq \frac{\tilde{R}_{i,k}}{(1 - \tilde{p}_{i+1,k})^{\tilde{b}\gamma_{i+1,k}/\gamma_{i,k}}}, \tag{3.71}$$

for $i = 0, \dots, D+1-k$. This holds true for an arbitrary choice of the numbers \tilde{b} and k .

In the special case in which the sequence $\{b_0, b_1, \dots, b_{D-1}\}$ is periodic of period k (where $D \geq k$), we can choose \tilde{b} to be the geometric mean of the b_i over one period:

$$\tilde{b} = \left(\prod_{i=0}^{k-1} b_i \right)^{1/k}. \tag{3.72}$$

Then the sequence $\{\gamma_0, \gamma_1, \dots, \gamma_D\}$ is also periodic of period k , so that $\gamma_{0,k} = \gamma_{1,k} = \dots = \gamma_{D+1-k,k}$, and (3.71) simplifies to

$$\tilde{p}_{i,k} \geq \frac{\tilde{R}_{i,k}}{(1 - \tilde{p}_{i+1,k})^{\tilde{b}}} \tag{3.73}$$

for $i = 0, \dots, D+1-k$ [note that for $i = D+1-k$ we have $\tilde{p}_{i+1,k} = 0$, so the value of the exponent $\tilde{b}\gamma_{i+1,k}/\gamma_{i,k}$ is in this case irrelevant].

So we can argue as before: suppose we fix the numbers $(\tilde{R}_{i,k})_{i=0}^{D+1-k}$ and define $(\hat{p}_{i,k})_{i=0}^{D+1-k}$ by the recursion

$$\hat{p}_{i,k} = \frac{\tilde{R}_{i,k}}{(1 - \hat{p}_{i+1,k})^{\tilde{b}\gamma_{i+1,k}/\gamma_{i,k}}} \tag{3.74}$$

(as long as $\hat{p}_{i+1,k}$ remains < 1), with initial condition $\hat{p}_{D+2-k,k} = 0$. Then it follows immediately from the monotonicity of the right-hand side of (3.71) that $\tilde{p}_{i,k} \geq \hat{p}_{i,k}$ as long as the former is well-defined. In particular, if for some level i we have $\hat{p}_{i,k} \geq 1$, then $\tilde{p}_{i,k}$ is either ≥ 1 or else ill-defined, so that there exists $x \in \bigcup_{j=i}^{i+k-1} X_j$ with $w_x^{\text{eff}} \notin (-1, 0]$. It then follows from Theorem 3.2 that $\mathbf{R} \notin \mathcal{R}(W)$. Moreover, since this calculation depends only on the numbers $(\tilde{R}_{i,k})_{i=0}^{D+1-k}$, the same conclusion holds for *any* \mathbf{R} with the given geometric means on-and-between levels.

The same argument works with soft-core pair interaction if we replace $|X_i|$ by $\tilde{\alpha}_i$. We have therefore proven:

Proposition 3.4 Consider any repulsive lattice gas with hard-core self-repulsion for which the support graph G_W is a tree. Define weights $(\alpha_x)_{x \in X}$ and $(\tilde{\alpha}_i)_{i=0}^D$ by (3.59)/(3.60). Fix an integer $k \geq 1$ and a real number $\bar{b} > 0$, and define $\gamma_i = \tilde{\alpha}_i / \bar{b}^i$. Then define $(\gamma_{i,k})_{i=0}^{D+1-k}$, $(\tilde{p}_{i,k})_{i=0}^{D+1-k}$ and $(\tilde{R}_{i,k})_{i=0}^{D+1-k}$ by (3.66)–(3.68). Finally, define numbers $(\hat{p}_{i,k})_{i=0}^D$ by the recursion

$$\hat{p}_{i,k} = \frac{\tilde{R}_{i,k}}{(1 - \hat{p}_{i+1,k})^{\bar{b}\gamma_{i+1,k}/\gamma_{i,k}}} \quad (3.75)$$

(as long as $\hat{p}_{i+1,k}$ remains < 1), with initial condition $\hat{p}_{D+2-k,k} = 0$. Suppose that there exists an i for which $\hat{p}_{i,k} \geq 1$. Then $\mathbf{R} \notin \mathcal{R}(W)$ for all vectors $\mathbf{R} \geq 0$ satisfying $\prod_{j=i}^{i+k-1} \left(\prod_{x \in X_j} R_x^{\alpha_x / \tilde{\alpha}_j} \right)^{\gamma_j / \gamma_{i,k}} \geq \tilde{R}_{i,k}$ for all $i = 0, \dots, D+1-k$.

We will return to these ideas in Section 8.2.

4 The Lovász local lemma

4.1 Hard-core version

Let $(A_x)_{x \in X}$ be a finite family of events on some probability space, and let G be a graph with vertex set X . We say that G is a *dependency graph* for the family $(A_x)_{x \in X}$ if, for each $x \in X$, the event A_x is independent from the σ -algebra $\sigma(A_y: y \in X \setminus \Gamma^*(x))$. [Here we have used the notation $\Gamma^*(x) = \Gamma(x) \cup \{x\}$, where $\Gamma(x)$ is the set of vertices of G adjacent to x .] Note that this is much stronger than requiring merely that A_x be independent of each such A_y separately.

A family of events typically has many possible dependency graphs: for instance, if G is a dependency graph for events $(A_x)_{x \in X}$, then any graph obtained by adding edges to G is also a dependency graph. In particular, if the events A_x are independent, then any graph on X is a dependency graph. Nor must there be a unique minimal dependency graph. Consider, for instance, the set of binary strings of length n with odd digit sum (giving each such string equal probability), and let A_i be the event that the i th digit is 1. Any graph without isolated vertices is a dependency graph for this collection of events.

There is also a stronger notion of a dependency graph G for a collection of events $(A_x)_{x \in X}$, where we demand that if Y and Z are disjoint subsets of X such that G contains no edges between Y and Z , then the σ -algebras $\sigma(A_y: y \in Y)$ and $\sigma(A_z: z \in Z)$ are independent. In this case we shall refer to G as a *strong dependency graph* for the events $(A_x)_{x \in X}$.¹³ Alternatively, the dependency-graph hypothesis can be replaced by a *weaker* hypothesis concerning conditional probabilities, as in the lopsided Lovász

¹³For instance, this situation arises in any statistical-mechanical model with variables living on the set X and pair interactions only on the edges of G , where each A_x depends only on the variable at x .

local lemma (Theorem 1.2). It will follow from Theorem 4.1 below that all three hypotheses lead to the same lower bounds on $\mathbb{P}(\bigcap_{x \in X} \overline{A}_x)$.

The various forms of the Lovász local lemma (e.g. Theorems 1.1 and 1.2) provide a *sufficient* (but not necessary) condition to have $\mathbb{P}(\bigcap_{x \in X} \overline{A}_x) > 0$, in terms of the existence of numbers $(r_x)_{x \in X}$ satisfying $p_x \leq r_x \prod_{y \in \Gamma(x)} (1 - r_y)$ for suitable probabilities or conditional probabilities $\mathbf{p} = (p_x)_{x \in X}$. As discussed in the Introduction, we shall approach the problem by dividing our analysis into two parts. First, in this section, we examine a best-possible condition to have $\mathbb{P}(\bigcap_{x \in X} \overline{A}_x) > 0$, in terms of the independent-set polynomial $Z_G(-\mathbf{p})$; then, in the next section, we discuss a *sufficient* condition to have $Z_G(-\mathbf{p}) > 0$, in terms of the existence of such numbers $(r_x)_{x \in X}$ or generalizations thereof.

The following result is a development of Shearer [98, Theorem 1].

Theorem 4.1 *Let $(A_x)_{x \in X}$ be a family of events on some probability space, and let G be a graph with vertex set X . Suppose that $(p_x)_{x \in X}$ are real numbers in $[0, 1]$ such that, for each x and each $Y \subseteq X \setminus \Gamma^*(x)$, we have*

$$\mathbb{P}(A_x | \bigcap_{y \in Y} \overline{A}_y) \leq p_x. \quad (4.1)$$

(a) *If $\mathbf{p} \in \mathcal{R}(G)$, then*

$$\mathbb{P}(\bigcap_{x \in X} \overline{A}_x) \geq Z_G(-\mathbf{p}) > 0 \quad (4.2)$$

and more generally

$$\mathbb{P}(\bigcap_{x \in Y} \overline{A}_x | \bigcap_{x \in Z} \overline{A}_x) \geq \frac{Z_G(-\mathbf{p} \mathbf{1}_{Y \cup Z})}{Z_G(-\mathbf{p} \mathbf{1}_Z)} > 0 \quad (4.3)$$

for any subsets $Y, Z \subseteq X$. Moreover, this lower bound is best possible in the sense that there exists a probability space on which there can be constructed a family of events $(B_x)_{x \in X}$ with probabilities $\mathbb{P}(B_x) = p_x$ and strong dependency graph G , such that $\mathbb{P}(\bigcap_{x \in X} \overline{B}_x) = Z_G(-\mathbf{p})$.

(b) *If $\mathbf{p} \notin \mathcal{R}(G)$, then there exists a probability space on which there can be constructed:*

- (i) *A family of events $(B_x)_{x \in X}$ with probabilities $\mathbb{P}(B_x) = p_x$ and strong dependency graph G , satisfying $\mathbb{P}(\bigcap_{x \in X} \overline{B}_x) = 0$; and*
- (ii) *A family of events $(B'_x)_{x \in X}$ with probabilities $\mathbb{P}(B'_x) = p'_x \leq p_x$ and strong dependency graph G , satisfying $\mathbb{P}(B'_x \cap B'_y) = 0$ for all $xy \in E(G)$ and $\mathbb{P}(\bigcap_{x \in X} \overline{B}'_x) = 0$.*

Remarks. 1. Please note that G is here an *arbitrary* graph with vertex set X ; it need not be a dependency graph for the events $(A_x)_{x \in X}$. Rather, given G , we can regard \mathbf{p} as *defined* by

$$p_x = \max_{Y \subseteq X \setminus \Gamma^*(x)} \mathbb{P}(A_x | \bigcap_{y \in Y} \overline{A}_y) \quad (4.4)$$

(this is clearly the minimal choice). There is then a tradeoff in the choice of G : adding more edges reduces p_x (since there are fewer conditional probabilities to control) but also shrinks the set $\mathcal{R}(G)$ (by the last sentence of Proposition 2.15).

2. Though (4.1) is the *weak* hypothesis of the lopsided Lovász local lemma (Theorem 1.2), we will prove in (a) and (b) that the extremal families $(B_x)_{x \in X}$ and $(B'_x)_{x \in X}$ have G as a *strong* dependency graph. Therefore, all three dependency hypotheses lead to the same optimal lower bound on $\mathbb{P}(\bigcap_{x \in X} \overline{A}_x)$.

3. The proofs given here of Theorems 4.1 and 4.2 are logically independent of nearly all of Theorem 2.10. More precisely, if we define $\mathcal{R}(G)$ and $\mathcal{R}(W)$ by condition (b) of Theorem 2.10, then the only part of Theorem 2.10 that is used in the proofs of Theorems 4.1 and 4.2 is the (relatively easy) implication (b) \implies (f).

PROOF. For $\mathbf{p} \in \mathcal{R}(G)$, we wish to define a family of events $(B_x)_{x \in X}$ [on a new probability space] such that the hypotheses of the theorem are satisfied and $\mathbb{P}(\bigcap_{x \in X} \overline{B}_x)$ is as small as possible. An intuitively reasonable way to do this is to make the events B_x as disjoint as possible, consistent with the condition (4.1) [or with either of the two stronger notions of dependency graph]. With this in mind, for $\Lambda \subseteq X$ let us define

$$\mathbb{P}\left(\bigcap_{x \in \Lambda} B_x\right) = \begin{cases} \prod_{x \in \Lambda} p_x & \text{if } \Lambda \text{ is independent in } G \\ 0 & \text{otherwise} \end{cases} \quad (4.5)$$

This defines a signed measure on the σ -algebra generated by $(B_x)_{x \in X}$; indeed, inclusion-exclusion gives

$$\mathbb{P}\left(\bigcap_{x \in \Lambda} B_x \cap \bigcap_{x \notin \Lambda} \overline{B}_x\right) = \sum_{I \supseteq \Lambda} (-1)^{|I| - |\Lambda|} \mathbb{P}\left(\bigcap_{x \in I} B_x\right) \quad (4.6a)$$

$$= \sum_{I \supseteq \Lambda, I \text{ independent}} (-1)^{|I| - |\Lambda|} \prod_{x \in I} p_x \quad (4.6b)$$

$$= (-1)^{|\Lambda|} Z_G(-\mathbf{p}; \Lambda), \quad (4.6c)$$

where $Z_G(-\mathbf{p}; \Lambda)$ is defined as in (2.37). In particular, taking $\Lambda = \emptyset$, we have $\mathbb{P}(\bigcap_{x \in X} \overline{B}_x) = Z_G(-\mathbf{p})$. Theorem 2.10(f) implies that (4.6c) is nonnegative for all Λ , so that (4.5) defines a probability measure on $\sigma(B_x: x \in X)$. [This is the probability measure defined in Theorem 2.10(g).]

If Y and Z are disjoint subsets of X such that G contains no edges between Y and Z , it follows from (4.5) that for $Y_0 \subseteq Y$ and $Z_0 \subseteq Z$ the events $\bigcap_{x \in Y_0} B_x$ and $\bigcap_{x \in Z_0} B_x$ are independent. This implies (see, for instance, [116, Theorem 4.2] or [16, Theorem 4.2]) that $\sigma(B_x: x \in Y)$ and $\sigma(B_x: x \in Z)$ are independent, and so G is a strong dependency graph.

We next show that $(B_x)_{x \in X}$ is a family minimizing $\mathbb{P}(\bigcap_{x \in X} \overline{B}_x)$. For $\Lambda \subseteq X$, we define

$$P_\Lambda = \mathbb{P}\left(\bigcap_{x \in \Lambda} \overline{A}_x\right) \quad (4.7)$$

$$Q_\Lambda = \mathbb{P}\left(\bigcap_{x \in \Lambda} \overline{B}_x\right). \quad (4.8)$$

Let us now prove by induction on $|\Lambda|$ that P_Λ/Q_Λ is monotone increasing in Λ . Note first that by inclusion-exclusion,

$$Q_\Lambda = \sum_{I \subseteq \Lambda} (-1)^{|I|} \mathbb{P}\left(\bigcap_{x \in I} B_x\right) \quad (4.9a)$$

$$= \sum_{I \subseteq \Lambda, I \text{ independent}} (-1)^{|I|} \prod_{x \in I} p_x \quad (4.9b)$$

$$= Z_G(-\mathbf{p} \mathbf{1}_\Lambda). \quad (4.9c)$$

Thus $Q_\Lambda > 0$ for all Λ , since $\mathbf{p} \in \mathcal{R}(G)$ and $\mathcal{R}(G)$ is a down-set. Furthermore, for $y \notin \Lambda$,

$$Q_{\Lambda \cup \{y\}} = \sum_{I \subseteq \Lambda \cup \{y\}, I \text{ independent}} (-1)^{|I|} \prod_{x \in I} p_x \quad (4.10a)$$

$$= Q_\Lambda - p_y \sum_{I \subseteq \Lambda \setminus \Gamma(y), I \text{ independent}} (-1)^{|I|} \prod_{x \in I} p_x \quad (4.10b)$$

$$= Q_\Lambda - p_y Q_{\Lambda \setminus \Gamma(y)}. \quad (4.10c)$$

[Note that this is just the fundamental identity (3.5) applied to $Z_G(-\mathbf{p} \mathbf{1}_\Lambda)$.] On the other hand,

$$P_{\Lambda \cup \{y\}} = P_\Lambda - \mathbb{P}(A_y \cap \bigcap_{x \in \Lambda} \bar{A}_x) \quad (4.11a)$$

$$\geq P_\Lambda - \mathbb{P}(A_y \cap \bigcap_{x \in \Lambda \setminus \Gamma(y)} \bar{A}_x) \quad (4.11b)$$

$$\geq P_\Lambda - p_y P_{\Lambda \setminus \Gamma(y)} \quad (4.11c)$$

by the hypothesis (4.1). Now we want to show that $P_{\Lambda \cup \{y\}}/Q_{\Lambda \cup \{y\}} \geq P_\Lambda/Q_\Lambda$, or equivalently that $P_{\Lambda \cup \{y\}}Q_\Lambda - Q_{\Lambda \cup \{y\}}P_\Lambda \geq 0$. By (4.10) and (4.11) we have

$$P_{\Lambda \cup \{y\}}Q_\Lambda - Q_{\Lambda \cup \{y\}}P_\Lambda \geq [P_\Lambda - p_y P_{\Lambda \setminus \Gamma(y)}]Q_\Lambda - [Q_\Lambda - p_y Q_{\Lambda \setminus \Gamma(y)}]P_\Lambda \quad (4.12a)$$

$$= p_y [P_\Lambda Q_{\Lambda \setminus \Gamma(y)} - Q_\Lambda P_{\Lambda \setminus \Gamma(y)}] \quad (4.12b)$$

$$\geq 0 \quad (4.12c)$$

since

$$\frac{P_\Lambda}{Q_\Lambda} \geq \frac{P_{\Lambda \setminus \Gamma(y)}}{Q_{\Lambda \setminus \Gamma(y)}} \quad (4.13)$$

by the inductive hypothesis.

Since P_Λ/Q_Λ is monotone increasing in Λ , we have $P_X/Q_X \geq P_\emptyset/Q_\emptyset = 1$, which proves (4.2). More generally, for any subsets $Y, Z \subseteq X$, we have $P_{Y \cup Z}/Q_{Y \cup Z} \geq P_Z/Q_Z$ and hence $P_{Y \cup Z}/P_Z \geq Q_{Y \cup Z}/Q_Z$, which gives (4.3).

For $\mathbf{p} \notin \mathcal{R}(G)$, choose a minimal vector $\mathbf{p}' \leq \mathbf{p}$ such that $\mathbf{p}' \geq 0$ and $Z_G(-\mathbf{p}') = 0$ [such a \mathbf{p}' is in general nonunique]. Then the family of events $(B'_x)_{x \in X}$ defined by (4.5) with p_x replaced by p'_x satisfies $\mathbb{P}(\bigcap_{x \in X} \bar{B}'_x) = Z_G(-\mathbf{p}') = 0$ [by (4.6c) with

$\Lambda = \emptyset$]. Since \mathbf{p}' is in the closure of $\mathcal{R}(G)$, it follows by the minimality of \mathbf{p}' and the continuity of Z_G that this is a well-defined probability measure; note that if x and y are adjacent then $\mathbb{P}(B'_x \cap B'_y) = 0$ by (4.5). Thus we have constructed a collection of events satisfying part (ii) of the Theorem.

To construct a collection of events satisfying part (i), let $(C_x)_{x \in X}$ be an (independent) collection of independent events satisfying

$$[1 - \mathbb{P}(B'_x)][1 - \mathbb{P}(C_x)] = 1 - p_x. \quad (4.14)$$

Then the events $B_x = B'_x \cup C_x$ satisfy $\mathbb{P}(B_x) = p_x$ and $\mathbb{P}(\bigcap \overline{B}_x) \leq \mathbb{P}(\bigcap \overline{B}'_x) = 0$. ■

Remarks. 1. If $(A_x)_{x \in X}$ is a family of events satisfying (4.2) with equality, then we have $P_X = Q_X$ in the foregoing proof; and since $P_\emptyset = Q_\emptyset = 1$, the monotonicity of P_Λ/Q_Λ implies that we have $P_\Lambda = Q_\Lambda$ for every $\Lambda \subseteq X$. Thus, if $(A_x)_{x \in X}$ is an extremal family, the probabilities of all events in $\sigma(A_x: x \in X)$ are completely determined and are given by (4.5)/(4.6).

2. The Lovász local lemma can be formulated more generally for families of events with a dependency *digraph*: each event A_x is independent from the σ -algebra $\sigma(A_y: y \in X \setminus \Gamma_+^*(x))$, where $\Gamma_+^*(x) = \Gamma_+(x) \cup \{x\}$ and $\Gamma_+(x)$ is the out-neighborhood of x . See e.g. [2, Lemma 5.1.1] or [23, Theorem 1.17]. It would be interesting to have a digraph analogue of Theorem 4.1, but we do not know how to do this.

3. There are other probabilistic inequalities that are expressed in terms of a dependency graph (see for instance Suen [109] or Janson [68]); it would be interesting to know if any of these have counterparts in statistical mechanics. One obstacle here is the need for a counterpart of Theorem 4.1. However, even without such a result, there may be scope for proving further inequalities in the presence of weak dependency conditions of the form discussed in the next subsection.

4.2 Soft-core version

Let us now consider how to extend Theorem 4.1 to the more general case of a soft-core pair interaction, i.e. to allow “soft edges” xy of strength $1 - W(x, y) \in [0, 1]$. The first step here is to replace the hard-core dependency condition (4.1) by an appropriate soft-core version.

Let $W: X \times X \rightarrow [0, 1]$ be symmetric and satisfy $W(x, x) = 0$ for all $x \in X$; and let $(A_x)_{x \in X}$ be a collection of events in some probability space. For each $x \in X$, let S_x be a random subset of X , independent of the σ -algebra $\sigma(A_x: x \in X)$, defined by the probabilities

$$\mathbb{P}(y \in S_x) = W(x, y) \quad (4.15)$$

independently for each $y \in X$. [Thus in the case of a hard-core pair interaction, we have $S_x = X \setminus \Gamma^*(x)$ with probability 1.] Let $(p_x)_{x \in X}$ be real numbers in $[0, 1]$. We say that $(A_x)_{x \in X}$ satisfies the *weak dependency conditions with interaction W and*

probabilities $(p_x)_{x \in X}$ if, for each $x \in X$ and each $Y \subseteq X \setminus x$ we have

$$\mathbb{E} \left(\mathbb{P}(A_x \cap \bigcap_{y \in Y \cap S_x} \bar{A}_y) \right) \leq p_x \mathbb{E} \left(\mathbb{P}(\bigcap_{y \in Y \cap S_x} \bar{A}_y) \right). \quad (4.16)$$

[Note that in the special case of a hard-core pair interaction, we have $Y \cap S_x = Y \setminus \Gamma^*(x)$ with probability 1, so that (4.16) reduces to (4.1).] Of course, the reference here to a random subset S_x can be replaced by an explicit expression for the probabilities $\mathbb{P}(Y \cap S_x = Y')$, so that (4.16) is equivalent to

$$\begin{aligned} \sum_{Y' \subseteq Y} \left(\prod_{y \in Y'} W(x, y) \right) \left(\prod_{y \in Y \setminus Y'} [1 - W(x, y)] \right) \mathbb{P}(A_x \cap \bigcap_{y \in Y'} \bar{A}_y) \leq \\ p_x \sum_{Y' \subseteq Y} \left(\prod_{y \in Y'} W(x, y) \right) \left(\prod_{y \in Y \setminus Y'} [1 - W(x, y)] \right) \mathbb{P}(\bigcap_{y \in Y'} \bar{A}_y). \end{aligned} \quad (4.17)$$

We can now state a soft-core version of Theorem 4.1:

Theorem 4.2 *Let $(A_x)_{x \in X}$ be a family of events in some probability space, and let $W: X \times X \rightarrow [0, 1]$ be symmetric and satisfy $W(x, x) = 0$ for all $x \in X$. Suppose that $(A_x)_{x \in X}$ satisfies the weak dependency conditions (4.16)/(4.17) with interaction W and probabilities $(p_x)_{x \in X}$.*

(a) *If $\mathbf{p} \in \mathcal{R}(W)$, then*

$$\mathbb{P}(\bigcap_{x \in X} \bar{A}_x) \geq Z_W(-\mathbf{p}) > 0 \quad (4.18)$$

and more generally

$$\mathbb{P}(\bigcap_{x \in Y} \bar{A}_x \mid \bigcap_{x \in Z} \bar{A}_x) \geq \frac{Z_W(-\mathbf{p} \mathbf{1}_{Y \cup Z})}{Z_W(-\mathbf{p} \mathbf{1}_Z)} > 0 \quad (4.19)$$

for any subsets $Y, Z \subseteq X$. Furthermore, this bound is best possible in the sense that there exists a family $(B_x)_{x \in X}$ with probabilities $\mathbb{P}(B_x) = p_x$ that satisfies the weak dependency conditions (4.16)/(4.17) with interaction W and probabilities $(p_x)_{x \in X}$, has strong dependency graph G_W , and has $\mathbb{P}(\bigcap_{x \in X} \bar{B}_x) = Z_W(-\mathbf{p})$.

(b) *If $\mathbf{p} \notin \mathcal{R}(W)$, then there exists a probability space on which there can be constructed:*

- (i) *A family of events $(B_x)_{x \in X}$ with probabilities $\mathbb{P}(B_x) = p_x$ and satisfying the weak dependency conditions (4.16)/(4.17) with interaction W , such that $\mathbb{P}(\bigcap_{x \in X} \bar{B}_x) = 0$; and*
- (ii) *A family of events $(B'_x)_{x \in X}$ with probabilities $\mathbb{P}(B'_x) = p'_x \leq p_x$ and satisfying the weak dependency conditions (4.16)/(4.17) with interaction W , such that $\mathbb{P}(B'_x \cap B'_y) = W(x, y)\mathbb{P}(B'_x)\mathbb{P}(B'_y)$ for all x, y and $\mathbb{P}(\bigcap_{x \in X} \bar{B}'_x) = 0$.*

PROOF. For $\mathbf{p} \in \mathcal{R}(W)$, we define a family of events $(B_x)_{x \in X}$ [on a new probability space] by

$$\mathbb{P}\left(\bigcap_{x \in \Lambda} B_x\right) = \left(\prod_{x \in \Lambda} p_x\right) \left(\prod_{\{x,y\} \subseteq \Lambda} W(x,y)\right) \quad (4.20)$$

[here the second product runs over all two-element subsets $\{x,y\} \subseteq X$ ($x \neq y$)]. As before, inclusion-exclusion gives

$$\mathbb{P}\left(\bigcap_{x \in \Lambda} B_x \cap \bigcap_{x \notin \Lambda} \overline{B}_x\right) = \sum_{I \supseteq \Lambda} (-1)^{|I| - |\Lambda|} \mathbb{P}\left(\bigcap_{x \in I} B_x\right) \quad (4.21a)$$

$$= \sum_{I \supseteq \Lambda} (-1)^{|I|} \prod_{x \in I} (-p_x) \prod_{\{x,y\} \subseteq I} W(x,y) \quad (4.21b)$$

$$= (-1)^{|\Lambda|} Z_W(-\mathbf{p}; \Lambda). \quad (4.21c)$$

In particular, we have $\mathbb{P}(\bigcap_{x \in X} \overline{B}_x) = Z_W(-\mathbf{p}) > 0$, and more generally for $\Lambda \subseteq X$ we have

$$\begin{aligned} \mathbb{P}\left(\bigcap_{x \in \Lambda} \overline{B}_x\right) &= \sum_{I \subseteq \Lambda} (-1)^{|I|} \mathbb{P}\left(\bigcap_{x \in I} \overline{B}_x\right) \\ &= \sum_{I \subseteq \Lambda} (-1)^{|I|} \left(\prod_{x \in I} p_x\right) \left(\prod_{\{x,y\} \subseteq \Lambda} W(x,y)\right) \\ &= Z_W(-\mathbf{p} \mathbf{1}_\Lambda) \\ &> 0. \end{aligned}$$

We define P_Λ and Q_Λ as in (4.7)/(4.8). Note that $Q_\Lambda = Z_W(-\mathbf{p} \mathbf{1}_\Lambda) > 0$, as $Q_X = Z_W(-\mathbf{p}) > 0$ and $\mathcal{R}(W)$ is a down-set. Let us now prove by induction on $|\Lambda|$ that P_Λ/Q_Λ is monotone increasing in Λ . For $\Lambda \subseteq X$ and $y \in X \setminus \Lambda$, we have

$$Q_{\Lambda \cup \{y\}} = Z_W(-\mathbf{p} \mathbf{1}_{\Lambda \cup \{y\}}) \quad (4.22a)$$

$$= Z_W(-\mathbf{p} \mathbf{1}_\Lambda) - p_y Z_W(-W(y, \cdot) \mathbf{p} \mathbf{1}_\Lambda) \quad (4.22b)$$

$$= Z_W(-\mathbf{p} \mathbf{1}_\Lambda) - p_y \sum_{Y \subset \Lambda} \left(\prod_{y \in Y} W(x,y)\right) \left(\prod_{y \in \Lambda \setminus Y} [1 - W(x,y)]\right) Z_W(-\mathbf{p} \mathbf{1}_Y) \quad (4.22c)$$

$$= Q_\Lambda - p_y \sum_{Y \subset \Lambda} \left(\prod_{y \in Y} W(x,y)\right) \left(\prod_{y \in \Lambda \setminus Y} [1 - W(x,y)]\right) Q_Y \quad (4.22d)$$

where we have used the fundamental identity (3.3)/(3.6). On the other hand,

$$P_{\Lambda \cup \{y\}} = P_\Lambda - \mathbb{P}\left(A_y \cap \bigcap_{x \in \Lambda} \overline{A}_x\right) \quad (4.23a)$$

$$= P_\Lambda - \sum_{Y \subseteq \Lambda} \left(\prod_{x \in Y} W(x, y) \right) \left(\prod_{x \in \Lambda \setminus Y} [1 - W(x, y)] \right) \mathbb{P}(A_y \cap \bigcap_{x \in \Lambda} \bar{A}_x) \quad (4.23b)$$

$$\geq P_\Lambda - \sum_{Y \subseteq \Lambda} \left(\prod_{x \in Y} W(x, y) \right) \left(\prod_{x \in \Lambda \setminus Y} [1 - W(x, y)] \right) \mathbb{P}(A_y \cap \bigcap_{x \in Y} \bar{A}_x) \quad (4.23c)$$

$$\geq P_\Lambda - p_y \sum_{Y \subseteq \Lambda} \left(\prod_{x \in Y} W(x, y) \right) \left(\prod_{x \in \Lambda \setminus Y} [1 - W(x, y)] \right) \mathbb{P}(\bigcap_{x \in Y} \bar{A}_x). \quad (4.23d)$$

where the last line uses the weak dependency condition (4.16)/(4.17). We then have

$$\frac{P_\Lambda}{Q_\Lambda} Q_{\Lambda \cup \{y\}} = P_\Lambda - p_y \sum_{Y \subseteq \Lambda} \left(\prod_{x \in Y} W(x, y) \right) \left(\prod_{x \in \Lambda \setminus Y} [1 - W(x, y)] \right) \frac{P_\Lambda}{Q_\Lambda} Q_Y \quad (4.24a)$$

$$\leq P_\Lambda - p_y \sum_{Y \subseteq \Lambda} \left(\prod_{x \in Y} W(x, y) \right) \left(\prod_{x \in \Lambda \setminus Y} [1 - W(x, y)] \right) P_Y \quad (4.24b)$$

$$\leq P_{\Lambda \cup \{y\}}, \quad (4.24c)$$

where the first inequality uses the inductive hypothesis and the second inequality uses (4.23). Hence $P_{\Lambda \cup \{y\}}/Q_{\Lambda \cup \{y\}} \geq P_\Lambda/Q_\Lambda$ as claimed.

The bounds (4.18) and (4.19) follow as in Theorem 4.1.

Part (b) follows as in Theorem 4.1. For $\mathbf{p} \notin \mathcal{R}(W)$ choose a minimal vector \mathbf{p}' with $\mathbf{0} \leq \mathbf{p}' \leq \mathbf{p}$ and $Z_W(-\mathbf{p}') = 0$. The events $(B'_x)_{x \in X}$ defined by (4.20) with p_x replaced by p'_x satisfy $\mathbb{P}(\bigcap_{x \in X} \bar{B}'_x) = 0$ and $\mathbb{P}(B'_x \cap B'_y) = W(x, y)\mathbb{P}(B'_x)\mathbb{P}(B'_y)$ for all x, y . A collection of events satisfying (i) is constructed as before. ■

Remark. The uniqueness of a family of events satisfying (4.18) with equality follows as before: since $P_X/Q_X = P_\emptyset/Q_\emptyset = 1$ and P_Λ/Q_Λ is increasing we have $P_\Lambda = Q_\Lambda$ for all $\Lambda \subseteq X$ and so the probabilities of events are given by (4.21).

5 Sufficient conditions for $Z_W \neq 0$ in a polydisc

In this section we shall exhibit sufficient conditions on a set of radii $\mathbf{R} = \{R_x\}_{x \in X}$ so that the partition function $Z_W(\mathbf{w})$ is nonvanishing in the closed polydisc $|\mathbf{w}| \leq \mathbf{R}$. We shall restrict attention to the case of a repulsive lattice gas with hard-core self-repulsion, i.e. $0 \leq W(x, y) \leq 1$ for all x, y and $W(x, x) = 0$ for all x , and we shall use the notation introduced in Section 3.1. Our main tool will be the fundamental identity (3.3), applied inductively.

5.1 Basic bound

Our first (and most basic) bound is due to Dobrushin [37, 38] in the case of a hard-core interaction; the generalization to a soft repulsive interaction was proven recently by one of us [101]. The method of proof is, however, already implicit (in more powerful form) in Shearer [98, Theorem 2].

Theorem 5.1 (Dobrushin [37, 38], Sokal [101]) *Let X be a finite set, and let W satisfy*

- (a) $0 \leq W(x, y) \leq 1$ for all $x, y \in X$
- (b) $W(x, x) = 0$ for all $x \in X$

Let $\mathbf{R} = \{R_x\}_{x \in X} \geq 0$. Suppose that there exist constants $\{K_x\}_{x \in X}$ satisfying $0 \leq K_x < 1/R_x$ and

$$K_x \geq \prod_{y \neq x} \frac{1 - W(x, y)K_y R_y}{1 - K_y R_y} \quad (5.1)$$

for all $x \in X$. Then, for each subset $\Lambda \subseteq X$, $Z_\Lambda(\mathbf{w})$ is nonvanishing in the closed polydisc $\bar{D}_{\mathbf{R}} = \{\mathbf{w} \in \mathbb{C}^X: |w_x| \leq R_x \text{ for all } x\}$ and satisfies there

$$\left| \frac{\partial \log Z_\Lambda(\mathbf{w})}{\partial w_x} \right| \leq \begin{cases} \frac{K_x}{1 - K_x |w_x|} & \text{for all } x \in \Lambda \\ 0 & \text{for all } x \in X \setminus \Lambda \end{cases} \quad (5.2)$$

Moreover, if $\mathbf{w}, \mathbf{w}' \in \bar{D}_{\mathbf{R}}$ and $w'_x/w_x \in [0, +\infty]$ for each $x \in \Lambda$, then

$$\left| \log \frac{Z_\Lambda(\mathbf{w}')}{Z_\Lambda(\mathbf{w})} \right| \leq \sum_{x \in \Lambda} \left| \log \frac{1 - K_x |w'_x|}{1 - K_x |w_x|} \right| \quad (5.3)$$

where on the left-hand side we take the standard branch of the log, i.e. $|\operatorname{Im} \log \cdots| \leq \pi$.

Remark. It follows from (5.1) that $K_x \geq 1$ and hence that $R_x < 1$.

PROOF. Note first that (5.2) for any given Λ implies (5.3) for the same Λ , by integration.

The proof is by induction on the cardinality of Λ . If $\Lambda = \emptyset$ the claims are trivial. So let us assume that (5.2) [and hence also (5.3)] holds for all sets of cardinality $< n$, and let a set Λ of cardinality n be given. Let x be any element of Λ . Let us apply the fundamental identity (3.3), and observe that $W(x, \cdot)\mathbf{w} \in \bar{D}_{\mathbf{R}}$ since $|W(x, y)| \leq 1$. Therefore, by the inductive hypothesis we have $Z_{\Lambda \setminus x}(\mathbf{w}) \neq 0$ and $Z_{\Lambda \setminus x}(W(x, \cdot)\mathbf{w}) \neq 0$; and from (3.3) we have

$$\frac{\partial}{\partial w_x} \log Z_\Lambda(\mathbf{w}) = \frac{\mathcal{K}_{x, \Lambda}(\mathbf{w})}{1 + \mathcal{K}_{x, \Lambda}(\mathbf{w})w_x} \quad (5.4)$$

where

$$\mathcal{K}_{x,\Lambda}(\mathbf{w}) = \frac{Z_{\Lambda \setminus x}(W(x, \cdot) \mathbf{w})}{Z_{\Lambda \setminus x}(\mathbf{w})}. \quad (5.5)$$

Now by the inductive hypothesis (5.3) for $\Lambda \setminus x$, and using the fact that $\mathbf{w}' = W(x, \cdot) \mathbf{w}$ satisfies $w'_y/w_y = W(x, y) \geq 0$, we have

$$|\mathcal{K}_{x,\Lambda}(\mathbf{w})| \leq \prod_{y \in \Lambda \setminus x} \frac{1 - W(x, y) K_y |w_y|}{1 - K_y |w_y|} \leq \prod_{y \in X \setminus x} \frac{1 - W(x, y) K_y |w_y|}{1 - K_y |w_y|}, \quad (5.6)$$

which is $\leq K_x$ by the hypothesis (5.1). This proves (5.2) for Λ , and hence completes the induction. ■

It is convenient to rewrite Theorem 5.1 in terms of the new variables $r_x = K_x R_x$:

Corollary 5.2 *Let X be a finite set, and let W satisfy $0 \leq W(x, y) \leq 1$ for all $x, y \in X$ and $W(x, x) = 0$ for all $x \in X$. Suppose that there exist constants $0 \leq r_x < 1$ satisfying*

$$R_x \leq r_x \prod_{y \neq x} \frac{1 - r_y}{1 - W(x, y) r_y} \quad (5.7)$$

for all $x \in X$. Then, for all \mathbf{w} satisfying $|\mathbf{w}| \leq \mathbf{R}$, the partition function Z_W satisfies

$$|Z_W(\mathbf{w})| \geq Z_W(-\mathbf{R}) \geq \prod_{x \in X} (1 - r_x) > 0 \quad (5.8)$$

and more generally

$$\left| \frac{Z_W(\mathbf{w} \mathbf{1}_{Y \cup Z})}{Z_W(\mathbf{w} \mathbf{1}_Z)} \right| \geq \prod_{x \in Y} (1 - r_x) > 0. \quad (5.9)$$

In particular, if we define the “maximum weighted degree”

$$\Delta_W = \max_{x \in X} \sum_{y \neq x} [1 - W(x, y)] \quad (5.10)$$

and write

$$F(\Delta_W) = \frac{2 + \Delta_W - \sqrt{\Delta_W^2 + 4\Delta_W}}{2} \quad (5.11)$$

$$R(\Delta_W) = F(\Delta_W) e^{-[1 - F(\Delta_W)]} \quad (5.12)$$

we have

$$|Z_W(\mathbf{w})| \geq [1 - F(\Delta_W)]^{|\mathbf{X}|} > 0 \quad (5.13)$$

whenever $|w_x| \leq R(\Delta_W)$ for all $x \in X$.

PROOF. Setting $r_x = K_x R_x$, we find that (5.1) becomes (5.7), and (5.3) with $\Lambda = X$ and $\mathbf{w}' = 0$ becomes (5.8).

To obtain the last claim, note first that

$$\frac{1-r}{1-Wr} = \frac{1-r}{1-r+(1-W)r} = \frac{1}{1+(1-W)\frac{r}{1-r}} \geq e^{-(1-W)r/(1-r)} \quad (5.14)$$

whenever $0 \leq W \leq 1$ and $0 \leq r \leq 1$. Therefore, if we set $r_x = r$ for all $x \in X$, we have

$$r_x \prod_{y \neq x} \frac{1-r_y}{1-W(x,y)r_y} \geq r e^{-\Delta_W r/(1-r)}. \quad (5.15)$$

We then choose r to maximize the right-hand side of (5.15); simple calculus yields $r = F(\Delta_W)$ and $\Delta_W r = (1-r)^2$, so that the right-hand side of (5.15) is bounded below by $R(\Delta_W)$. It follows that if we define $R_x = R(\Delta_W)$ and $r_x = F(\Delta_W)$ for all $x \in X$ then (5.7) and so (5.8) are satisfied. ■

Remarks. 1. The radius $R(\Delta_W)$ behaves as

$$R(\Delta_W) = \begin{cases} 1 - 2\Delta_W^{1/2} + \frac{5}{2}\Delta_W + O(\Delta_W^{3/2}) & \text{as } \Delta_W \rightarrow 0 \\ \frac{1}{\epsilon\Delta_W} \left[1 - \frac{1}{\Delta_W} + \frac{3}{2\Delta_W} + O(\Delta_W^{-3}) \right] & \text{as } \Delta_W \rightarrow \infty \end{cases} \quad (5.16)$$

Example 3.6 (the r -ary rooted tree) shows that this bound is sharp (to leading order) as $\Delta_W \rightarrow \infty$. At the other extreme, the $1 - \text{const} \times \Delta_W^{1/2}$ behavior at small Δ_W is also best possible, since the two-site lattice gas with $W(x,x) = W(y,y) = 0$ and $W(x,y) = 1 - \epsilon$ has $Z_W(w) = 1 + 2w + (1 - \epsilon)w^2$ and hence has a root at $w = -1/(1 + \sqrt{\epsilon})$. [However, the coefficient 2 rather than 1 in the $\Delta_W^{1/2}$ term of (5.16) may not be best possible.]

2. From Proposition 2.15(c) we know that the set $\mathcal{R}(W)$ is log-convex; it is therefore natural to ask whether the subset of $\mathcal{R}(W)$ produced by (5.7) is also log-convex. (If it weren't, then we could improve Corollary 5.2 by taking the log-convex hull.) It turns out that the set of vectors satisfying (5.7) is indeed always log-convex: in fact, if (\mathbf{r}, \mathbf{R}) and $(\mathbf{r}', \mathbf{R}')$ are two pairs of vectors satisfying (5.7), and $0 \leq \lambda \leq 1$, then $(\mathbf{r}^\lambda \mathbf{r}'^{1-\lambda}, \mathbf{R}^\lambda \mathbf{R}'^{1-\lambda})$ also satisfies (5.7). In the hard-core case this follows from the inequality $1 - r_y^\lambda (r'_y)^{1-\lambda} \geq (1 - r_y)^\lambda (1 - r'_y)^{1-\lambda}$, which is proven by two applications of the weighted arithmetic-geometric mean inequality as in (3.55). In the general case it can be shown by a similar argument (using four applications of the weighted arithmetic-geometric mean inequality!).

Specializing Corollary 5.2 to the case of a hard-core pair interaction for a graph G ,

$$W(x,y) = \begin{cases} 0 & \text{if } x = y \text{ or } xy \in E(G) \\ 1 & \text{if } x \neq y \text{ and } xy \notin E(G) \end{cases} \quad (5.17)$$

we have:

Corollary 5.3 *Let G be a finite graph with vertex set X , and let $\mathbf{R} = \{R_x\}_{x \in X} \geq 0$. Suppose that there exist constants $0 \leq r_x < 1$ satisfying*

$$R_x \leq r_x \prod_{y \in \Gamma(x)} (1 - r_y) \quad (5.18)$$

for all $x \in X$. Then, for all \mathbf{w} satisfying $|\mathbf{w}| \leq \mathbf{R}$, the independent-set polynomial Z_G satisfies

$$|Z_G(\mathbf{w})| \geq Z_G(-\mathbf{R}) \geq \prod_{x \in X} (1 - r_x) > 0 \quad (5.19)$$

and more generally

$$\left| \frac{Z_G(\mathbf{w} \mathbf{1}_{Y \cup Z})}{Z_G(\mathbf{w} \mathbf{1}_Z)} \right| \geq \prod_{x \in Y} (1 - r_x) > 0. \quad (5.20)$$

In particular, if G has maximum degree Δ , then $|Z_G(\mathbf{w})| \geq [\Delta/(\Delta + 1)]^{|X|} > 0$ whenever $|w_x| \leq \Delta^\Delta/(\Delta + 1)^{\Delta+1}$ for all $x \in X$.

PROOF. The last claim is obtained by setting $r_x = 1/(\Delta + 1)$ for all $x \in X$. ■

Remark. The radius $\Delta^\Delta/(\Delta + 1)^{\Delta+1}$ behaves for large Δ as

$$\frac{\Delta^\Delta}{(\Delta + 1)^{\Delta+1}} = \frac{1}{e\Delta} \left[1 - \frac{1}{2\Delta} + \frac{7}{24\Delta^2} - \frac{3}{16\Delta^3} + O(\Delta^{-4}) \right], \quad (5.21)$$

which agrees with (5.16) to leading order in $1/\Delta$ but is slightly larger (hence better) at order $1/\Delta^2$.

Combining Corollary 5.3 with Theorem 4.1, we immediately obtain the lopsided Lovász local lemma (Theorem 1.2). Combining Corollary 5.2 with Theorem 4.2, we obtain a “soft-core” version of the lopsided Lovász local lemma:

Theorem 5.4 *Let $(A_x)_{x \in X}$ be a family of events in some probability space, and let $W: X \times X \rightarrow [0, 1]$ be symmetric and satisfy $W(x, x) = 0$ for all $x \in X$. Suppose that $(A_x)_{x \in X}$ satisfies the weak dependency conditions (4.16)/(4.17) with interaction W and probabilities $(p_x)_{x \in X}$. Suppose further that $(r_x)_{x \in X}$ are real numbers in $[0, 1]$ satisfying*

$$p_x \leq r_x \prod_{y \in \Gamma(x)} (1 - r_y). \quad (5.22)$$

Then

$$\mathbb{P}\left(\bigcap_{x \in X} \bar{A}_x\right) \geq \prod_{x \in X} (1 - r_x) > 0, \quad (5.23)$$

and more generally for sets $Y, Z \subseteq X$, we have

$$\mathbb{P}\left(\bigcap_{x \in Y} \bar{A}_x \mid \bigcap_{x \in Z} \bar{A}_x\right) \geq \prod_{x \in Y \setminus Z} (1 - r_x) = 0. \quad (5.24)$$

Defining the weighted degree Δ_W as in (5.10), we obtain the following:

Lemma 5.5 *Let $(A_x)_{x \in X}$ satisfy the weak dependency conditions (4.16)/(4.17) with interaction W and probabilities $(p_x)_{x \in X}$. If $p_x < \Delta_W^{\Delta_W} / (\Delta_W + 1)^{\Delta_W + 1}$ for every $x \in X$, then $\mathbb{P}(\bigcap_{x \in X} \overline{A_x}) > 0$.*

PROOF. As in the proof of Corollary 5.3, set $r_x = r \equiv 1/(\Delta_W + 1)$ for all $x \in X$. Then check (5.7):

$$\begin{aligned} r_x \prod_{y \neq x} \frac{1 - r_y}{1 - W(x, y)r_y} &\leq r_x \prod_{y \neq x} (1 - r_y)^{1 - W(x, y)} \\ &\leq r(1 - r)^{\Delta_W} \\ &\leq \frac{\Delta_W^{\Delta_W}}{(\Delta_W + 1)^{\Delta_W + 1}}. \end{aligned} \tag{5.25}$$

In the first inequality we have used the fact that $1 - W(x, y)r_y \leq (1 - r_y)^{W(x, y)}$ for $0 \leq W(x, y) \leq 1$. ■

It would be interesting to see applications of Theorem 5.4 and Lemma 5.5.

5.2 Improved bound

Let us now attempt to improve the bound of Theorem 5.1. Note, first of all, that we need not insist that the bound (5.2) hold with the *same* constant K_x for all $\Lambda \ni x$; rather, we can use constants $K_{x, \Lambda}$ that depend on Λ . Inspection of the inductive argument shows that we can *define* the constants $K_{x, \Lambda} \in [0, +\infty]$ as a function of the family $\{R_x\}$ by the recursion

$$K_{x, \Lambda} = \begin{cases} \prod_{\substack{y \in \Lambda \setminus x \\ W(x, y) \neq 1 \\ R_y > 0}} \frac{1 - W(x, y)K_{y, \Lambda \setminus x}R_y}{1 - K_{y, \Lambda \setminus x}R_y} & \text{if } K_{y, \Lambda \setminus x}R_y < 1 \text{ for all terms in the product} \\ +\infty & \text{otherwise} \end{cases} \tag{5.26}$$

[Note that $K_{x, \{x\}} = 1$ for all x , because the product (5.26) is empty.] It follows immediately by induction that each $K_{x, \Lambda}$ is an increasing rational function of $\{R_y\}_{y \in \Lambda \setminus x}$ up to the first pole, and $+\infty$ thereafter. More precisely, suppose we define the rational functions $\widehat{K}_{x, \Lambda}(\mathbf{R})$ by the recursion (5.26) *without* the restrictions $K_{y, \Lambda \setminus x}R_y < 1$; then it is easy to see that

$$K_{x, \Lambda}(\mathbf{R}) = \begin{cases} \widehat{K}_{x, \Lambda}(\mathbf{R}) & \text{if } \widehat{K}_{x, \Lambda}(\mathbf{R}') < \infty \text{ for } 0 \leq \mathbf{R}' \leq \mathbf{R} \\ +\infty & \text{otherwise} \end{cases} \tag{5.27}$$

Moreover, it is easily proven by induction that all the partial derivatives of $\widehat{K}_{x, \Lambda}$ are nonnegative at $\mathbf{R} = 0$; it then follows from Proposition 2.11 that the Taylor series of

$\widehat{K}_{x,\Lambda}(\mathbf{R})$ about $\mathbf{0}$ converges throughout the region where the first case in (5.27) holds, and that all the partial derivatives of $K_{x,\Lambda} = \widehat{K}_{x,\Lambda}$ are nonnegative there. Finally, it is obvious from the definition (5.26) ff. that $K_{x,\Lambda}$ is an increasing function of Λ .

Let us now define a graph G with vertex set $V = \{x \in X: R_x > 0\}$ and edge set $E = \{x, y \in V: W(x, y) \neq 1\}$; and for each $\Lambda \subseteq X$, let G_Λ be the subgraph of G induced by $\Lambda \cap V$. Then only the connected component of G_Λ containing x plays any role in the definition of $K_{x,\Lambda}$: that is, if G_Λ has several connected components with vertex sets $\Lambda_1, \dots, \Lambda_k$ and $x \in \Lambda_i$, then $K_{x,\Lambda} = K_{x,\Lambda_i}$.

Let us now call a pair (x, Λ) “good” if $K_{x,\Lambda} < \infty$ and $K_{x,\Lambda}R_x < 1$. It follows immediately from the definition (5.26) ff. that if (x, Λ) is good, then $(y, \Lambda \setminus x)$ is also good whenever $y \in \Lambda \setminus x$ with $W(x, y) \neq 1$ and $R_y > 0$, i.e. whenever y is a neighbor of x in G_Λ . [Indeed, this follows under the weaker hypothesis that $K_{x,\Lambda} < \infty$.]

We then have:

Theorem 5.6 (improved Dobrushin–Shearer bound) *Let X be a finite set, and let W satisfy*

$$(a) \quad 0 \leq W(x, y) \leq 1 \text{ for all } x, y \in X$$

$$(b) \quad W(x, x) = 0 \text{ for all } x \in X$$

Let $\mathbf{R} = \{R_x\}_{x \in X} \geq 0$. Define the constants $K_{x,\Lambda} \in [0, +\infty]$ as above. Suppose that in each connected component of G_Λ there exists at least one vertex x for which the pair (x, Λ) is good. Then $Z_\Lambda(\mathbf{w})$ is nonvanishing in the closed polydisc $\bar{D}_\mathbf{R}$; and for every good pair (x, Λ) and every $\mathbf{w} \in \bar{D}_\mathbf{R}$, we have

$$\left| \frac{\partial \log Z_\Lambda(\mathbf{w})}{\partial w_x} \right| \leq \frac{K_{x,\Lambda}}{1 - K_{x,\Lambda}|w_x|}. \quad (5.28)$$

Moreover, if $\mathbf{w}, \mathbf{w}' \in \bar{D}_\mathbf{R}$ and $w'_x/w_x \in [0, +\infty]$ for each $x \in \Lambda$, and in addition the pair (x, Λ) is good whenever $w'_x \neq w_x$, then

$$\left| \log \frac{Z_\Lambda(\mathbf{w}')}{Z_\Lambda(\mathbf{w})} \right| \leq \sum_{\substack{x \in \Lambda \\ w'_x \neq w_x}} \left| \log \frac{1 - K_{x,\Lambda}|w'_x|}{1 - K_{x,\Lambda}|w_x|} \right|, \quad (5.29)$$

where on the left-hand side we take the standard branch of the log, i.e. $|\operatorname{Im} \log \dots| \leq \pi$.

PROOF. Note first that (5.28) for any given Λ implies (5.29) for the same Λ , by integration.

The proof is by induction on the cardinality of Λ . If $\Lambda = \emptyset$ the claims are trivial. So let us assume that (5.28) [and hence also (5.29)] holds for all sets of cardinality $< n$ satisfying the stated hypotheses, and let a set Λ of cardinality n satisfying these hypotheses be given. If G_Λ is not connected, i.e. has components with vertex sets $\Lambda_1, \dots, \Lambda_k$ ($k \geq 2$), then $Z_\Lambda(\mathbf{w})$ factorizes as

$$Z_\Lambda(\mathbf{w}) = \prod_{i=1}^k Z_{\Lambda_i}(\mathbf{w}); \quad (5.30)$$

moreover, each connected component has cardinality $< n$ and satisfies the hypotheses of the theorem, so it follows immediately from the inductive hypothesis that (5.28) and (5.29) hold also for Λ . We may therefore assume that G_Λ is connected.

Let x be any element of Λ for which the pair (x, Λ) is good. Since G_Λ is connected, the vertex x has at least one neighbor y in each connected component of $G_{\Lambda \setminus x}$. Moreover, as noted previously, the pair $(y, \Lambda \setminus x)$ is good whenever y is a neighbor of x in G_Λ . Therefore, the inductive hypothesis is applicable to $\Lambda \setminus x$. Let us now apply the fundamental identity (3.3), and observe that $W(x, \cdot)\mathbf{w} \in \bar{D}_{\mathbf{R}}$ since $|W(x, y)| \leq 1$. By the inductive hypothesis we have $Z_{\Lambda \setminus x}(\mathbf{w}) \neq 0$ and $Z_{\Lambda \setminus x}(W(x, \cdot)\mathbf{w}) \neq 0$; and from (3.3) we have

$$\frac{\partial}{\partial w_x} \log Z_\Lambda(\mathbf{w}) = \frac{\mathcal{K}_{x, \Lambda}(\mathbf{w})}{1 + \mathcal{K}_{x, \Lambda}(\mathbf{w})w_x} \quad (5.31)$$

where

$$\mathcal{K}_{x, \Lambda}(\mathbf{w}) = \frac{Z_{\Lambda \setminus x}(W(x, \cdot)\mathbf{w})}{Z_{\Lambda \setminus x}(\mathbf{w})}. \quad (5.32)$$

Now each $y \in \Lambda \setminus x$ with $w'_y \neq w_y$ necessarily has $W(x, y) \neq 1$ and $R_y > 0$ (i.e. is a neighbor of x in G_Λ), so that the pair $(y, \Lambda \setminus x)$ is good. We may therefore apply the inductive hypothesis (5.29); using the fact that $\mathbf{w}' = W(x, \cdot)\mathbf{w}$ satisfies $w'_y/w_y = W(x, y) \geq 0$, we have

$$|\mathcal{K}_{x, \Lambda}(\mathbf{w})| \leq \prod_{\substack{y \in \Lambda \setminus x \\ W(x, y) \neq 1 \\ R_y > 0}} \frac{1 - W(x, y)K_{y, \Lambda \setminus x}|w_y|}{1 - K_{y, \Lambda \setminus x}|w_y|} \leq K_{x, \Lambda} \quad (5.33)$$

since $|w_y| \leq R_y$. This proves (5.28) for (x, Λ) , and hence completes the induction. ■

Remark. It is unclear whether the set of vectors \mathbf{R} satisfying the hypotheses of Theorem 5.6 is log-convex. If it is not, then the conclusion of Theorem 5.6 can be improved by taking the log-convex hull.

As a corollary of Theorem 5.6, we can deduce a bound due originally (in the Lovász context) to Shearer [98, Theorem 2], which improves the last sentence of Corollary 5.3 by replacing Δ by $\Delta - 1$. Indeed, we can very slightly improve Shearer's bound by allowing *one* vertex x_0 to have a larger radius R_{x_0} :

Corollary 5.7 *Let $G = (X, E)$ be a finite graph of maximum degree $\Delta \geq 2$, and fix one vertex $x_0 \in X$. Suppose that $|w_{x_0}| \leq (\Delta - 1)^\Delta / \Delta^\Delta$ and that $|w_x| \leq (\Delta - 1)^{\Delta - 1} / \Delta^\Delta$ for all $x \neq x_0$. Then $Z_G(\mathbf{w}) \neq 0$.*

PROOF. Since Z_G factorizes over connected components, we can assume without loss of generality that G is connected. (Indeed, if G is disconnected, then we can allow one “ x_0 -like” vertex in *each* connected component.) Set $R_{x_0} = (\Delta - 1)^\Delta / \Delta^\Delta$ and $R_x = (\Delta - 1)^{\Delta - 1} / \Delta^\Delta$ for all $x \neq x_0$.

We first claim that if $x_0 \notin \Lambda$, and $x \in \Lambda$ is a vertex with at least one neighbor in $X \setminus \Lambda$, then

$$K_{x,\Lambda} < \left(\frac{\Delta}{\Delta-1} \right)^{\Delta-1} \quad (5.34)$$

(note the strict inequality). The proof is by induction on $|\Lambda|$, using the definition (5.26): it certainly holds if $\Lambda = \{x\}$; since every y appearing in the product on the right-hand side of (5.26) has at least one neighbor outside of $\Lambda \setminus x$ (namely, x itself), $K_{y,\Lambda \setminus x}$ satisfies (5.34) by the inductive hypothesis and so $K_{y,\Lambda \setminus x} R_y < 1/\Delta$; and finally, since x has at least one neighbor outside Λ , there are at most $\Delta - 1$ factors in the product. Thus

$$K_{x,\Lambda} < \left(\frac{1}{1-1/\Delta} \right)^{\Delta-1} = \left(\frac{\Delta}{\Delta-1} \right)^{\Delta-1}.$$

It then follows that

$$K_{x_0,X} < \left(\frac{\Delta}{\Delta-1} \right)^\Delta, \quad (5.35)$$

since the bound (5.34) applies to all the terms $K_{y,\Lambda \setminus x_0}$ appearing on the right-hand side of (5.26). We therefore have $K_{x_0,X} R_{x_0} < 1$, and so the pair (x_0, X) is good. The claim then follows from Theorem 5.6. ■

Replacing $\Delta^\Delta/(\Delta+1)^{\Delta+1}$ by $(\Delta-1)^{\Delta-1}/\Delta^\Delta$ may seem to be a negligible improvement, since both quantities have the same leading behavior $\approx 1/(e\Delta)$ as $\Delta \rightarrow \infty$, and differ only at higher order:

$$\frac{(\Delta-1)^{\Delta-1}}{\Delta^\Delta} = \frac{1}{e\Delta} \left[1 + \frac{1}{2\Delta} + \frac{7}{24\Delta^2} + \frac{3}{16\Delta^3} + O(\Delta^{-4}) \right] \quad (5.36)$$

[cf. (5.21)].¹⁴ But Shearer's bound $(\Delta-1)^{\Delta-1}/\Delta^\Delta$ has the great merit of being *best possible*: for, as he showed [98], if G is the complete rooted tree with branching factor $r = \Delta - 1$ and depth n , then $Z_G(w)$ has negative real zeros that tend to $w = -(\Delta-1)^{\Delta-1}/\Delta^\Delta$ as $n \rightarrow \infty$ (see Example 3.6 above).

We remark that Corollary 5.7 does not appear to extend naturally to the soft-core case (note that having one neighbor outside Λ in the argument around (5.34) need not reduce the weighted degree of a vertex in Λ by 1).

5.3 Optimal bound

Now let us try to further improve Theorem 5.6. Where in the proof did we lose equality? This happened only in (5.33), where we applied (5.29), which in turn arose from integrating (5.28): the point is that the bound (5.28) in general improves as we pass from \mathbf{w} downwards to $\mathbf{w}' = W(x, \cdot)\mathbf{w}$, but we failed to take advantage of this fact. The solution is to order (arbitrarily) the vertices y_1, \dots, y_k arising in the

¹⁴The amusing similarity of (5.21) and (5.36) arises from the fact that $-(-\Delta)^{-\Delta}/(-\Delta+1)^{-\Delta+1} = (\Delta-1)^{\Delta-1}/\Delta^\Delta$.

product (5.33) and to write $\mathcal{K}_{x,\Lambda}(\mathbf{w})$ as a telescoping product; then we will have a sharp bound, i.e. one that becomes equality when $\mathbf{w} = -\mathbf{R}$.

Let us show this first in the special case of a hard-core pair interaction: setting $\Gamma(x) \cap \Lambda = \{y_1, \dots, y_k\}$, we have

$$\mathcal{K}_{x,\Lambda}(\mathbf{w}) = \frac{Z_{\Lambda \setminus \Gamma^*(x)}(\mathbf{w})}{Z_{\Lambda \setminus x}(\mathbf{w})} = \prod_{i=1}^k \frac{Z_{\Lambda \setminus x \setminus \{y_1, \dots, y_i\}}(\mathbf{w})}{Z_{\Lambda \setminus x \setminus \{y_1, \dots, y_{i-1}\}}(\mathbf{w})}. \quad (5.37)$$

Therefore we can improve (5.33) by writing

$$|\mathcal{K}_{x,\Lambda}(\mathbf{w})| \leq \prod_{i=1}^k \frac{1}{1 - K_{y_i, \Lambda \setminus x \setminus \{y_1, \dots, y_{i-1}\}} R_{y_i}}, \quad (5.38)$$

and so we can replace the definition (5.26) by

$$K_{x,\Lambda}^{\text{opt}} = \prod_{i=1}^k \frac{1}{1 - K_{y_i, \Lambda \setminus x \setminus \{y_1, \dots, y_{i-1}\}}^{\text{opt}} R_{y_i}} \quad (5.39)$$

if $K_{y_i, \Lambda \setminus x \setminus \{y_1, \dots, y_{i-1}\}}^{\text{opt}} R_{y_i} < 1$ for all terms in the product, and $K_{x,\Lambda}^{\text{opt}} = +\infty$ otherwise (note that $K_{x,\{x\}}^{\text{opt}} = 1$). Moreover, if we use $K_{x,\Lambda}^{\text{opt}}$, then all the bounds in the proof become *equality* when $\mathbf{w} = -\mathbf{R}$.

In the general case of a “soft” interaction $W(x, y)$, things become slightly more complicated, since the vectors $\mathbf{w}' = W(x, \cdot)\mathbf{w}$ have their components depressed but not set to zero. To handle this case, we need to define the numbers $K_{x,\Lambda}^{\text{opt}}$ explicitly as functions of a vector $\mathbf{R}_{\neq x}$. We begin by writing $\mathcal{K}_{x,\Lambda}(w)$ as a telescoping product as in (3.10):

$$\mathcal{K}_{x,\Lambda}(\mathbf{w}) = \prod_{i=1}^k \frac{Z_{\Lambda \setminus x}(\tilde{\mathbf{w}}^{(i)})}{Z_{\Lambda \setminus x}(\tilde{\mathbf{w}}^{(i-1)})} \quad (5.40)$$

where the vectors $\tilde{\mathbf{w}}^{(i)}$ are defined by

$$(\tilde{\mathbf{w}}^{(i)})_y = \begin{cases} W(x, y)w_y & \text{if } y = y_j \text{ for some } j \leq i \\ w_y & \text{otherwise} \end{cases} \quad (5.41)$$

Therefore we can improve (5.33) by writing

$$|\mathcal{K}_{x,\Lambda}(\mathbf{w})| \leq \prod_{i=1}^k \frac{1 - W(x, y_i)K_{y_i, \Lambda \setminus x}(\tilde{\mathbf{R}}_{\neq x}^{(i-1)})R_{y_i}}{1 - K_{y_i, \Lambda \setminus x}(\tilde{\mathbf{R}}_{\neq x}^{(i-1)})R_{y_i}} \quad (5.42)$$

[where $\tilde{\mathbf{R}}_{\neq x}^{(i)}$ is defined by the obvious analogue of (5.41)], and so we could have replaced the definition (5.26) by

$$K_{x,\Lambda}^{\text{opt}} = \prod_{i=1}^k \frac{1 - W(x, y_i)K_{y_i, \Lambda \setminus x}^{\text{opt}}(\tilde{\mathbf{R}}_{\neq x}^{(i-1)})R_{y_i}}{1 - K_{y_i, \Lambda \setminus x}^{\text{opt}}(\tilde{\mathbf{R}}_{\neq x}^{(i-1)})R_{y_i}} \quad (5.43)$$

if $K_{y_i, \Lambda \setminus x}^{\text{opt}}(\tilde{\mathbf{R}}_{\neq x}^{(i-1)})R_{y_i} < 1$ for all terms in the product, and $K_{x, \Lambda}^{\text{opt}} = +\infty$ otherwise.

Another way of looking at all this is: We want to choose the constants $K_{x, \Lambda}$ so that the bound (5.28) holds. By (3.8) and (3.9), the optimal choice is manifestly

$$K_{x, \Lambda}^{\text{opt}} = \sup_{0 \leq \mathbf{R}' \leq \mathbf{R}} \mathcal{K}_{x, \Lambda}(-\mathbf{R}'), \quad (5.44)$$

where $\mathcal{K}_{x, \Lambda}(\mathbf{w})$ was defined in (3.7). We now claim that (5.44) is identical to (5.43) et seq. Indeed, the recursion (3.13) shows that the rational function $\mathcal{K}_{x, \Lambda}(-\mathbf{R})$ is identical to what one obtains from the recursion (5.43) if one *omits* the condition following the equation; and the monotonicity of (5.43) up to the first pole guarantees that implementing the condition following the equation is equivalent to taking the supremum over all \mathbf{R}' satisfying $0 \leq \mathbf{R}' \leq \mathbf{R}$.

In summary, this second improvement of Theorem 5.1 is *optimal* in the sense that it is equivalent to calculating the *exact* $\mathcal{K}_{x, \Lambda}(\mathbf{w})$ and hence [by (3.16)] the exact $Z_\Lambda(\mathbf{w})$. So this “optimally improved Dobrushin–Shearer theorem” is not usually going to be useful in practice; but it does give insight into what has been lost in Theorems 5.1 and 5.6.

6 Tree interpretation

Further insight into Theorem 5.6 and its “optimal” improvement à la (5.43)/(5.44) can be obtained by considering the tree structure underlying the recursions (5.26) and (5.43).

6.1 Tree interpretation of Theorem 5.6

Let us first consider (5.26), which provides a recursive definition of the functions $K_{x, \Lambda}(\mathbf{R})$ [or $\widehat{K}_{x, \Lambda}(\mathbf{R})$ if we ignore the conditions $K_{y, \Lambda \setminus x}R_y < 1$ following (5.26)]. This recursion can be encoded as a tree: at the top (root) of the tree is the pair (x, Λ) ; immediately underneath it are all the pairs $(y, \Lambda \setminus x)$ for which $W(x, y) \neq 1$ [that is, for which xy is an edge of the support graph $G = G_W$ restricted to Λ]; and so on. Another way of saying this is as follows:

Definition 6.1 (see also [78]) *Let G be a simple loopless graph, and let $x_1 \in V(G)$. Then the path-tree or self-avoiding-walk tree (SAW-tree) of G rooted at x_1 — let us call it $\text{SAW}(G, x_1)$ — is the graph whose vertex set is the set of paths in G starting at x_1 [i.e. paths $\overline{x_1 x_2 \cdots x_k}$ in G with $k \geq 1$] and whose edges connect P to P' whenever P' is a one-step extension of P [i.e. whenever $P' = \overline{x_1 \cdots x_k}$ with $k \geq 2$ and $P = \overline{x_1 \cdots x_{k-1}}$]. The root of $\text{SAW}(G, x_1)$ is the zero-step path $\overline{x_1}$. [Note that if G happens to be a tree, then $\text{SAW}(G, x_1)$ is isomorphic to G for any $x_1 \in V(G)$.]*

We then associate a path $\overline{x_1 \cdots x_k}$ with the pair (x_k, Λ) where $\Lambda = V(G) \setminus \{x_1, \dots, x_{k-1}\}$. Note that a given pair (x_k, Λ) can correspond to many paths $\overline{x_1 \cdots x_k}$, which arise from different orderings of the set $\{x_2, \dots, x_{k-1}\} = V(G) \setminus \Lambda \setminus x_1$.

Given a nonnegative vector $\mathbf{R} = \{R_x\}_{x \in V(G)}$, let us now assign to the vertices of $\text{SAW}(G, x_1)$ the fugacities

$$\widehat{w}_{\overline{x_1 \cdots x_k}} = -R_{x_k}, \quad (6.1)$$

and to the edges of $\text{SAW}(G, x_1)$ the weights

$$\widehat{W}(\overline{x_1 \cdots x_{k-1}}, \overline{x_1 \cdots x_k}) = W(x_{k-1}, x_k). \quad (6.2)$$

We can then apply Algorithm T from Section 3.4 to calculate the effective fugacities $\widehat{w}_{\overline{x_1 \cdots x_k}}^{\text{eff}}$ on the tree $\text{SAW}(G, x_1)$, working upwards from the leaves:

Proposition 6.2 *In the foregoing set-up, we have*

$$\widehat{w}_{\overline{x_1 \cdots x_k}}^{\text{eff}} = -R_{x_k} \widehat{K}_{x_k, V(G) \setminus \{x_1, \dots, x_{k-1}\}}(\mathbf{R}) \quad (6.3)$$

where the rational functions $\widehat{K}_{x, \Lambda}(\mathbf{R})$ are defined by the recursion (5.26) ignoring the conditions $K_{y, \Lambda \setminus x} R_y < 1$.

PROOF. The proof runs inductively up from the leaves.

If $\overline{x_1 \cdots x_k}$ is a leaf (i.e. a maximal SAW), then $\widehat{w}_{\overline{x_1 \cdots x_k}}^{\text{eff}} = \widehat{w}_{\overline{x_1 \cdots x_k}} = -R_{x_k}$ by Algorithm T and the definition (6.1); moreover, $\widehat{K}_{x_k, V(G) \setminus \{x_1, \dots, x_{k-1}\}} \equiv 1$ because x_k is an isolated vertex in $G \setminus \{x_1, \dots, x_{k-1}\}$ [since $\overline{x_1 \cdots x_k}$ is a maximal SAW], so that the product (5.26) is empty.

Let $P = \overline{x_1 \cdots x_k}$ be a non-leaf, and suppose that (6.3) holds for all one-step extensions $P' = \overline{x_1 \cdots x_k x_{k+1}}$. Then Algorithm T gives

$$\begin{aligned} \widehat{w}_{\overline{x_1 \cdots x_k}}^{\text{eff}} &= \widehat{w}_{\overline{x_1 \cdots x_k}} \prod_{x_{k+1} \in V(G) \setminus \{x_1, \dots, x_k\}} \frac{1 + W(x_k, x_{k+1}) \widehat{w}_{\overline{x_1 \cdots x_k x_{k+1}}}^{\text{eff}}}{1 + \widehat{w}_{\overline{x_1 \cdots x_k x_{k+1}}}^{\text{eff}}} \\ &= -R_{x_k} \prod_{x_{k+1} \in V(G) \setminus \{x_1, \dots, x_k\}} \frac{1 - W(x_k, x_{k+1}) R_{x_{k+1}} \widehat{K}_{x_{k+1}, V(G) \setminus \{x_1, \dots, x_k\}}}{1 - R_{x_{k+1}} \widehat{K}_{x_{k+1}, V(G) \setminus \{x_1, \dots, x_k\}}} \\ &= -R_{x_k} \widehat{K}_{x_k, V(G) \setminus \{x_1, \dots, x_{k-1}\}}, \end{aligned} \quad (6.4)$$

where the second equality uses the inductive hypothesis, and the last equality follows immediately from (5.26). \blacksquare

Now let us apply Theorem 3.2 to the tree $\text{SAW}(G, x_1)$ with the fugacities $\widehat{\mathbf{w}}$ defined by (6.1) and edge weights \widehat{W} defined by (6.2). We conclude that the vector $\widehat{\mathbf{R}}$ defined by

$$\widehat{R}_{\overline{x_1 \cdots x_k}} = R_{x_k} \quad (6.5)$$

lies in $\mathcal{R}(\widehat{W})$ if and only if $R_x \widehat{K}_{x, \Lambda} < 1$ for all $\Lambda \subseteq V(G)$ and all $x \in \Lambda$. [By (6.3), this latter condition is sufficient to have $\mathcal{R}(\widehat{W})$; indeed, only certain sets $\Lambda \subseteq V(G)$ actually arise. Conversely, given any $\Lambda \subseteq V(G)$ and any $x \in \Lambda$, we can choose

$x_1 = x$ and restrict to the subtree of $\text{SAW}(G, x_1)$ consisting of paths inside Λ . Using the monotonicity of $\mathcal{R}(\widehat{W})$ to restrict to this subtree, we conclude by Algorithm T that $\widehat{w}_{x_1}^{\text{eff}} \in (-1, 0]$ and hence by (6.3) that $R_x \widehat{K}_{x, \Lambda} < 1$.] On the other hand, by Theorem 5.6 this is a *sufficient* condition to have $\mathbf{R} \in \mathcal{R}(W)$.

In summary, the bounds produced by Theorem 5.6 for the lattice gas on G with edge weights W correspond to solving exactly (via Algorithm T) the lattice gas on the tree $\text{SAW}(G, x_1)$ with edge weights \widehat{W} [for any $x_1 \in V(G)$]. And this produces a *lower* bound on the set $\mathcal{R}(W)$.

6.2 Tree interpretation of the “optimal” bound: hard-core case

Next let us consider the “optimal” recursion (5.43) in the special case of a hard-core pair interaction, i.e. the independent-set polynomial for the graph G .

For each vertex $\overline{x_1 \cdots x_k}$ of the SAW-tree, let us choose (in any way we like) an ordering of the children $\overline{x_1 \cdots x_k x_{k+1}}$. We then define the *pruned SAW-tree* corresponding to this ordering to be the subtree of $\text{SAW}(G, x_1)$ whose vertex set consists of those paths $\overline{x_1 \cdots x_k}$ satisfying the rule that “you can’t use an elder sibling of a vertex of G that you have previously used”. More precisely, for each path $\overline{x_1 \cdots x_i}$, let us define the set $S(\overline{x_1 \cdots x_i})$ of “spurned vertices at step i ” to be the set of all $x \in V(G)$ such that $\overline{x_1 \cdots x_{i-1} x}$ is a path that precedes $\overline{x_1 \cdots x_{i-1} x_i}$ in the ordering of children of $\overline{x_1 \cdots x_{i-1}}$. A path $\overline{x_1 \cdots x_k}$ then belongs to the pruned SAW-tree if and only if for all $2 \leq i < j \leq k$ we have $x_j \notin S(\overline{x_1 \cdots x_i})$.

Let us remark that the pruned SAW-tree is in general much smaller than the full SAW-tree. For example, if $G = K_n$, then the SAW-tree has $(n-1)!$ leaves and $(n-1)! \sum_{i=0}^{n-1} 1/i!$ vertices, while the pruned SAW-tree (for any choice of ordering) has 2^{n-2} leaves and 2^{n-1} vertices.

Having defined the pruned SAW-tree, we then identify a path $\overline{x_1 \cdots x_k}$ in the pruned SAW-tree with the pair (x_k, Λ) where

$$\Lambda \equiv \Lambda(x_1, \dots, x_k) = V(G) \setminus \{x_1, \dots, x_{k-1}\} \setminus \bigcup_{i=2}^k S(\overline{x_1 \cdots x_i}), \quad (6.6)$$

that is, $V(G)$ minus the vertices already visited or already spurned. [Note that this differs from the definition $\Lambda = V(G) \setminus \{x_1, \dots, x_{k-1}\}$ used in the preceding subsection.]

Given a vector $\mathbf{w} = \{w_x\}_{x \in V(G)}$ of fugacities on G , let us now assign to the vertices of the pruned SAW-tree the fugacities

$$\widehat{w}_{\overline{x_1 \cdots x_k}} = w_{x_k} \quad (6.7)$$

and, as before, assign to the edges the weights

$$\widehat{W}(\overline{x_1 \cdots x_{k-1}}, \overline{x_1 \cdots x_k}) = W(x_{k-1}, x_k) \quad (6.8)$$

(which, in the hard-core case currently under consideration, takes the value 0 for each edge of the pruned SAW-tree). We can then apply Algorithm T from Section 3.4 to the pruned SAW-tree to calculate the effective fugacities $\widehat{w}_{x_1 \cdots x_k}^{\text{eff}}$:

Proposition 6.3 For all $\overline{x_1 \cdots x_k}$ belonging to the pruned SAW-tree, we have

$$\widehat{w}_{\overline{x_1 \cdots x_k}}^{\text{eff}} = w_{x_k} \mathcal{K}_{x_k, \Lambda(x_1, \dots, x_k)}(\mathbf{w}). \quad (6.9)$$

PROOF. The proof runs inductively up from the leaves.

If $\overline{x_1 \cdots x_k}$ is a leaf (i.e. a maximal *pruned* SAW), then $\widehat{w}_{\overline{x_1 \cdots x_k}}^{\text{eff}} = \widehat{w}_{\overline{x_1 \cdots x_k}} = w_{x_k}$ by Algorithm T and the definition (6.7); moreover, $\mathcal{K}_{x_k, \Lambda(x_1, \dots, x_k)} \equiv 1$ because x_k is an isolated vertex in the subgraph of G induced by $\Lambda(x_1, \dots, x_k)$ [since $\overline{x_1 \cdots x_k}$ is a maximal pruned SAW], so that all factors in the product (3.14) are 1.

Let $P = \overline{x_1 \cdots x_k}$ be a non-leaf, and suppose that (6.9) holds for all one-step extensions $P' = \overline{x_1 \cdots x_k x_{k+1}}$ that are pruned SAWs. These one-step extensions are children of $\overline{x_1 \cdots x_k}$, hence are ordered: let the i th such walk in order be $\overline{x_1 \cdots x_k y_i}$ ($1 \leq i \leq l$ for some $l \geq 1$). Note that $\{y_1, \dots, y_l\} = \Gamma(x_k) \cap \Lambda(x_1, \dots, x_k)$ and that

$$\Lambda(x_1, \dots, x_k, y_i) = \Lambda(x_1, \dots, x_k) \setminus \{x_k, y_1, \dots, y_{i-1}\}. \quad (6.10)$$

Then Algorithm T gives

$$\begin{aligned} \widehat{w}_{\overline{x_1 \cdots x_k}}^{\text{eff}} &= \widehat{w}_{\overline{x_1 \cdots x_k}} \prod_{i=1}^l \frac{1 + W(x_k, y_i) \widehat{w}_{\overline{x_1 \cdots x_k y_i}}^{\text{eff}}}{1 + \widehat{w}_{\overline{x_1 \cdots x_k y_i}}^{\text{eff}}} \\ &= \widehat{w}_{\overline{x_1 \cdots x_k}} \prod_{i=1}^l \frac{1}{1 + w_{y_i} \mathcal{K}_{y_i, \Lambda(x_1, \dots, x_k, y_i)}(\mathbf{w})} \\ &= \widehat{w}_{\overline{x_1 \cdots x_k}} \prod_{i=1}^l \frac{1}{1 + w_{y_i} \mathcal{K}_{y_i, \Lambda(x_1, \dots, x_k) \setminus \{x_k, y_1, \dots, y_{i-1}\}}(\mathbf{w})} \\ &= \widehat{w}_{\overline{x_1 \cdots x_k}} \mathcal{K}_{x_k, \Lambda(x_1, \dots, x_k)}(\mathbf{w}) \end{aligned} \quad (6.11)$$

where the second equality uses the inductive hypothesis and the fact that $W(x_k, y_i) = 0$, the third equality uses (6.10), and the final equality uses (3.14). ■

Now let us apply Theorem 3.2 to the pruned SAW-tree with the fugacities $\widehat{\mathbf{w}}$ defined by (6.7) with $\mathbf{w} = -\mathbf{R}$, and edge weights \widehat{W} defined by (6.8). We conclude that the vector $\widehat{\mathbf{R}}$ defined by

$$\widehat{R}_{\overline{x_1 \cdots x_k}} = R_{x_k} \quad (6.12)$$

lies in $\mathcal{R}(\widehat{W})$ if and only if $R_x \mathcal{K}_{x, \Lambda}(-\mathbf{R}) < 1$ for all $\Lambda \subseteq V(G)$ and all $x \in \Lambda$; and by the discussion surrounding (5.43)/(5.44), this is a *necessary and sufficient* condition to have $\mathbf{R} \in \mathcal{R}(W)$.

Therefore, we have shown that the “optimal” Dobrushin–Shearer bound à la (5.43)/(5.44) for the independent-set polynomial (= hard-core lattice gas) on G corresponds to computing exactly (via Algorithm T) the independent-set polynomial for the *pruned* SAW-tree [for any $x_1 \in V(G)$ and any choice of orderings of children]. And this produces an *exact computation* of the set $\mathcal{R}(W)$.

6.3 Tree interpretation of the “optimal” bound: general case

Finally, let us consider the “optimal” recursion (5.43) in the general case of a soft-core pair interaction W . Here we work again on the full SAW-tree $\text{SAW}(G, x_1)$, where $G = G_W$ is the support graph of W ; “pruning” will be replaced by a “soft suppression of spurned vertices”. As before, we begin by choosing (in any way we like) an ordering on the children of each vertex in the SAW-tree, and we use this ordering to define the set $S(\overline{x_1 \cdots x_i})$ of “spurned vertices at step i ”. Then, given a vector $\mathbf{w} = \{w_x\}_{x \in V(G)}$ of fugacities on G , we assign to the vertices of the SAW-tree the modified fugacities

$$\widehat{w}_{\overline{x_1 \cdots x_k}} = w_{x_k} \prod_{\substack{2 \leq i \leq k \\ x_k \in S(\overline{x_1 \cdots x_i})}} W(x_{i-1}, x_k). \quad (6.13)$$

[Note that the product in (6.13) could equally well be written $2 \leq i < k$, since $x_k \notin S(\overline{x_1 \cdots x_k})$.] As before, we assign to the edges the weights

$$\widehat{W}(\overline{x_1 \cdots x_{k-1}}, \overline{x_1 \cdots x_k}) = W(x_{k-1}, x_k). \quad (6.14)$$

We can then apply Algorithm T to calculate the effective fugacities $\widehat{w}_{\overline{x_1 \cdots x_k}}^{\text{eff}}$ on the tree $\text{SAW}(G, x_1)$. We obtain:

Proposition 6.4 *In the foregoing set-up, we have*

$$\widehat{w}_{\overline{x_1 \cdots x_k}}^{\text{eff}} = \widetilde{w}_{x_k}^{[x_1 \cdots x_k]} \mathcal{K}_{x_k, V(G) \setminus \{x_1, \dots, x_{k-1}\}}(\widetilde{\mathbf{w}}^{[x_1 \cdots x_k]}) \quad (6.15)$$

where $\widetilde{\mathbf{w}}^{[x_1 \cdots x_k]}$ is defined by

$$\widetilde{w}_y^{[x_1 \cdots x_k]} = w_y \prod_{\substack{2 \leq i \leq k \\ y \in S(\overline{x_1 \cdots x_i})}} W(x_{i-1}, y). \quad (6.16)$$

PROOF. The proof runs inductively up from the leaves.

If $\overline{x_1 \cdots x_k}$ is a leaf (i.e. a maximal SAW), then

$$\widehat{w}_{\overline{x_1 \cdots x_k}}^{\text{eff}} = \widehat{w}_{\overline{x_1 \cdots x_k}} = w_{x_k} \prod_{\substack{2 \leq i \leq k \\ x_k \in S(\overline{x_1 \cdots x_i})}} W(x_{i-1}, x_k) = \widetilde{w}_{x_k}^{[x_1 \cdots x_k]}, \quad (6.17)$$

where the first equality uses Algorithm T and the subsequent equalities are the definitions (6.13) and (6.16). On the other hand, $\mathcal{K}_{x_k, V(G) \setminus \{x_1, \dots, x_{k-1}\}} \equiv 1$ because $\overline{x_1 \cdots x_k}$ is a maximal SAW, so that the product (3.14) is empty.

Let $P = \overline{x_1 \cdots x_k}$ be a non-leaf, and suppose that (6.15) holds for all one-step extensions $P' = \overline{x_1 \cdots x_k x_{k+1}}$. These one-step extensions are children of $\overline{x_1 \cdots x_k}$,




Figure 1: The graphs G and \widehat{G}^{xy} .

hence are ordered: let the i th such walk in order be $\overline{x_1 \cdots x_k y_i}$ ($1 \leq i \leq l$ for some $l \geq 1$). Note that $\{y_1, \dots, y_l\} = \Gamma(x_k) \cap \{x_1, \dots, x_{k-1}\}$. Then Algorithm T gives

$$\begin{aligned} \widehat{w}_{x_1 \cdots x_k}^{\text{eff}} &= \widehat{w}_{x_1 \cdots x_k} \prod_{i=1}^l \frac{1 + W(x_k, y_i) \widehat{w}_{x_1 \cdots x_k y_i}^{\text{eff}}}{1 + \widehat{w}_{x_1 \cdots x_k y_i}^{\text{eff}}} \\ &= \widehat{w}_{x_k}^{[x_1 \cdots x_k]} \prod_{i=1}^l \frac{1 + W(x_k, y_i) \widetilde{w}_{y_i}^{[x_1 \cdots x_k y_i]} \mathcal{K}_{y_i, V(G) \setminus \{x_1, \dots, x_k\}}(\widetilde{\mathbf{w}}^{[x_1 \cdots x_k y_i]})}{1 + \widetilde{w}_{y_i}^{[x_1 \cdots x_k y_i]} \mathcal{K}_{y_i, V(G) \setminus \{x_1, \dots, x_k\}}(\widetilde{\mathbf{w}}^{[x_1 \cdots x_k y_i]})}. \end{aligned} \quad (6.18)$$

Note now that from (6.16) we have

$$\widetilde{w}_y^{[x_1 \cdots x_k y_i]} = \begin{cases} W(x_k, y) \widetilde{w}_y^{[x_1 \cdots x_k]} & \text{if } y = y_j \text{ for some } j < i \\ \widetilde{w}_y^{[x_1 \cdots x_k]} & \text{otherwise} \end{cases} \quad (6.19)$$

Therefore, the product on the right-hand side of (6.18) is identical to (3.13) if we set $x = x_k$ and $\Lambda = V(G) \setminus \{x_1, \dots, x_{k-1}\}$ and \mathbf{w} is replaced by $\widetilde{\mathbf{w}}^{[x_1 \cdots x_k]}$, for then (6.19) becomes precisely the vector $\widetilde{\mathbf{w}}^{(i-1)}$ defined in (3.11). Therefore, (6.18) equals the right-hand side of (6.15), as claimed. ■

7 Unfolding

In Section 6.2 we showed that the “optimal” Dobrushin–Shearer bound à la (5.43)/(5.44) can be interpreted, in the special case of the independent-set polynomial (= hard-core lattice gas) for a graph G , as an *exact computation* of the set $\mathcal{R}(G)$ based on computing exactly (via Algorithm T) the independent-set polynomial of the pruned SAW-tree of G . In this section we would like to show how this tree bound can be understood as arising from the repeated application of a single “unfolding” step.

Let $G = (V, E)$ be a graph, and let us select a pair of adjacent vertices x, y . As previously, $G \setminus x$ (resp. $G \setminus y$) denotes the graph obtained from G by deleting the vertex x (resp. y) and all edges incident with it. We then define \widehat{G}^{xy} to be the graph obtained from the disjoint union of $G' \equiv G \setminus y$ and $G'' \equiv G \setminus x$ by adjoining an extra edge connecting the vertex x in G' to the vertex y in G'' (Figure 1). To each vertex

z in $G \setminus x \setminus y \equiv (G \setminus x) \setminus y$, there corresponds a vertex z' in G' and a vertex z'' in G'' . Given a vector \mathbf{w} with index set $V(G)$, we define the corresponding “diagonal” vector $\widehat{\mathbf{w}}$ with index set $V(\widehat{G}^{xy})$ by

$$\widehat{w}_x = w_x \quad (7.1a)$$

$$\widehat{w}_y = w_y \quad (7.1b)$$

$$\widehat{w}_{z'} = w_z \quad (7.1c)$$

$$\widehat{w}_{z''} = w_z \quad (7.1d)$$

By considering the occupation of sites x and y , it is easily seen that

$$Z_G(\mathbf{w}) = Z_{G \setminus x \setminus y}(\mathbf{w}) + w_x Z_{G \setminus \Gamma^*(x)}(\mathbf{w}) + w_y Z_{G \setminus \Gamma^*(y)}(\mathbf{w}) \quad (7.2)$$

and

$$Z_{\widehat{G}^{xy}}(\widehat{\mathbf{w}}) = Z_{G \setminus x \setminus y}(\mathbf{w})^2 + w_x Z_{G \setminus x \setminus y}(\mathbf{w}) Z_{G \setminus \Gamma^*(x)}(\mathbf{w}) + w_y Z_{G \setminus x \setminus y}(\mathbf{w}) Z_{G \setminus \Gamma^*(y)}(\mathbf{w}) \quad (7.3a)$$

$$= Z_{G \setminus x \setminus y}(\mathbf{w}) Z_G(\mathbf{w}). \quad (7.3b)$$

We have the following fundamental result:

Theorem 7.1 $\mathbf{R} \in \mathcal{R}(G)$ if and only if $\widehat{\mathbf{R}} \in \mathcal{R}(\widehat{G}^{xy})$. In other words, $\mathcal{R}(G)$ is (isomorphic to) the “diagonal cross section” of $\mathcal{R}(\widehat{G}^{xy})$.

PROOF. Suppose first that $\mathbf{R} \in \mathcal{R}(G)$. Then we also have $\mathbf{R} \in \mathcal{R}(G \setminus x \setminus y)$ by the monotonicity statement at the end of Proposition 2.15. Therefore, if $-\mathbf{R} \leq \mathbf{w} \leq 0$ we have $Z_G(\mathbf{w}) > 0$ and $Z_{G \setminus x \setminus y}(\mathbf{w}) > 0$, and hence deduce by (7.3b) that $Z_{\widehat{G}^{xy}}(\widehat{\mathbf{w}}) > 0$. In particular, $Z_{\widehat{G}^{xy}}(-\lambda \widehat{\mathbf{R}}) > 0$ for $0 \leq \lambda \leq 1$. Applying Theorem 2.10(a) \implies (b) to the line segment connecting 0 to $-\widehat{\mathbf{R}}$, we conclude that $Z_{\widehat{G}^{xy}}(\mathbf{w}') > 0$ whenever $-\widehat{\mathbf{R}} \leq \mathbf{w}' \leq 0$ (whether or not \mathbf{w}' lies “on the diagonal”). Hence $\widehat{\mathbf{R}} \in \mathcal{R}(\widehat{G}^{xy})$.

Conversely, suppose that $\widehat{\mathbf{R}} \in \mathcal{R}(\widehat{G}^{xy})$. If $-\mathbf{R} \leq \mathbf{w} \leq 0$ we have $-\widehat{\mathbf{R}} \leq \widehat{\mathbf{w}} \leq 0$ and hence $Z_{\widehat{G}^{xy}}(\widehat{\mathbf{w}}) > 0$; by (7.3b) this implies $Z_G(\mathbf{w}) \neq 0$. ■

Assume now that G is connected. Let \widetilde{G}^{xy} be the component of \widehat{G}^{xy} containing x and y . Any other component of \widehat{G}^{xy} must be contained either in $G' \setminus x$ or in $G'' \setminus y$; either way, it corresponds to some component of $G \setminus x \setminus y$. Now, for any component H of $G \setminus x \setminus y$, there are three possibilities: either H is adjacent to x in G , or H is adjacent to y in G , or both. In the first case, the copy H' of H in G' is contained in \widetilde{G}^{xy} , while the copy H'' of H in G'' is disjoint from \widetilde{G}^{xy} ; in the second case, the reverse holds; in the third case, both copies of H are contained in \widetilde{G}^{xy} . Therefore, any component of \widehat{G}^{xy} other than \widetilde{G}^{xy} has a mirror image contained in \widetilde{G}^{xy} , and hence is isomorphic to a subgraph of \widetilde{G}^{xy} (with the same weights when we are “on-diagonal”).

Given a vector \mathbf{w} with index set $V(G)$, let us define the vector $\widetilde{\mathbf{w}}$ with index set \widetilde{G}^{xy} by restricting $\widehat{\mathbf{w}}$ from $V(\widehat{G}^{xy})$ to $V(\widetilde{G}^{xy})$.

Lemma 7.2 *Let H be a subgraph of G , and let $\mathbf{R} = \{R_x\}_{x \in V(G)} \in \mathcal{R}(G)$. Then $\mathbf{R}_H \equiv \{R_x\}_{x \in V(H)} \in \mathcal{R}(H)$.*

PROOF. Define first $\mathbf{R} \mathbf{1}_{V(H)}$ [with index set $V(G)$] by

$$(\mathbf{R} \mathbf{1}_{V(H)})_x = \begin{cases} R_x & \text{if } x \in V(H) \\ 0 & \text{if } x \notin V(H) \end{cases} \quad (7.4)$$

Since $\mathbf{R} \mathbf{1}_{V(H)} \leq \mathbf{R}$, we have $\mathbf{R} \mathbf{1}_{V(H)} \in \mathcal{R}(G)$ because $\mathcal{R}(G)$ is a down-set. Equivalently, $\mathbf{R}_H \in \mathcal{R}(G[V(H)])$, where $G[V(H)]$ is the induced subgraph of G with vertex set $V(H)$. But $\mathcal{R}(H) \supseteq \mathcal{R}(G[V(H)])$ by the monotonicity statement at the end of Proposition 2.15. ■

Corollary 7.3 *Assume that G is connected. Then $\mathbf{R} \in \mathcal{R}(G)$ if and only if $\tilde{\mathbf{R}} \in \mathcal{R}(\tilde{G}^{xy})$.*

PROOF. If $\mathbf{R} \in \mathcal{R}(G)$, then $\hat{\mathbf{R}} \in \mathcal{R}(\hat{G}^{xy})$ by Theorem 7.1. Then $\tilde{\mathbf{R}} \in \mathcal{R}(\tilde{G}^{xy})$ by Lemma 7.2, because \tilde{G}^{xy} is a subgraph of \hat{G}^{xy} .

Conversely, suppose that $\tilde{\mathbf{R}} \in \mathcal{R}(\tilde{G}^{xy})$. Let H_1, \dots, H_k be the components of \hat{G}^{xy} other than \tilde{G}^{xy} . Then

$$Z_{\hat{G}^{xy}}(\hat{\mathbf{w}}) = Z_{\tilde{G}^{xy}}(\tilde{\mathbf{w}}) \prod_{i=1}^k Z_{H_i}(\hat{\mathbf{w}} \upharpoonright H_i) \quad (7.5)$$

If $-\mathbf{R} \leq \mathbf{w} \leq 0$, then $-\tilde{\mathbf{R}} \leq \tilde{\mathbf{w}} \leq 0$ and hence $Z_{\tilde{G}^{xy}}(\tilde{\mathbf{w}}) > 0$. But since each component of H_i is isomorphic to a subgraph of \tilde{G}^{xy} (with the same weights in the mirror copy), Lemma 7.2 implies that $Z_{H_i}(\hat{\mathbf{w}} \upharpoonright H_i) > 0$ as well. Hence $Z_{\hat{G}^{xy}}(\hat{\mathbf{w}}) > 0$ by (7.5), so that $Z_G(\mathbf{w}) \neq 0$ by (7.3b). This shows that $\mathbf{R} \in \mathcal{R}(G)$. ■

Remark. It would be interesting to know whether there is a version of unfolding in the case of soft-core interaction.

8 Infinite graphs

In this section we discuss briefly the repulsive lattice gas on a countably infinite graph. We begin by deriving some general properties valid on an arbitrary countably infinite graph (Section 8.1). Next we discuss two cases of special interest: trees (Section 8.2) and regular lattices (Section 8.3). Finally we show how quantitative bounds on $\mathcal{R}(W)$ can be obtained, using as an example the square lattice \mathbb{Z}^2 (Section 8.4).

8.1 General properties

Let X be a countably infinite set, and let $W: X \times X \rightarrow [0, 1]$ be symmetric. For every nonempty finite subset $\Lambda \subset X$ we can consider the partition function $Z_\Lambda(\mathbf{w})$ defined for $\mathbf{w} \in \mathbb{C}^\Lambda$ in the obvious way, i.e. by considering the lattice gas on Λ with interaction $W \upharpoonright \Lambda$. Then, to each such Λ there corresponds a set $\mathcal{R}(W \upharpoonright \Lambda) \subseteq [0, \infty)^\Lambda$. From the Remark after Proposition 2.15, we observe that if $\Lambda' \subseteq \Lambda$ and $\mathbf{R} \in [0, \infty)^{\Lambda'}$, then

$$\mathbf{R} \in \mathcal{R}(W \upharpoonright \Lambda') \iff (\mathbf{R}, \mathbf{0}) \in \mathcal{R}(W \upharpoonright \Lambda) \quad (8.1)$$

(here $\mathbf{0}$ has index set $\Lambda \setminus \Lambda'$).

Now let us define, for each finite $\Lambda \subset X$, the set

$$\mathcal{R}_\Lambda(W) = \mathcal{R}(W \upharpoonright \Lambda) \times [0, \infty)^{X \setminus \Lambda} \quad (8.2)$$

in the infinite-dimensional space $[0, \infty)^X$. Otherwise put, a vector $\mathbf{R} \in [0, \infty)^X$ belongs to $\mathcal{R}_\Lambda(W)$ if and only if $\mathbf{R} \upharpoonright \Lambda$ lies in $\mathcal{R}(W \upharpoonright \Lambda)$. Note that

$$\partial \mathcal{R}_\Lambda(W) = \partial \mathcal{R}(W \upharpoonright \Lambda) \times [0, \infty)^{X \setminus \Lambda}. \quad (8.3)$$

Clearly $\mathcal{R}_\Lambda(W)$ is open (in the product topology) and is a down-set. Moreover, it follows immediately from (8.1) and the fact that $\mathcal{R}(W \upharpoonright \Lambda)$ is a down-set that if $\Lambda' \subseteq \Lambda$, then $\mathcal{R}_{\Lambda'}(W) \supseteq \mathcal{R}_\Lambda(W)$. In other words, the $\{\mathcal{R}_\Lambda(W)\}$ form a decreasing family of sets in $[0, \infty)^X$ [when Λ runs over the collection of finite subsets of X ordered by inclusion]. We define the limiting set

$$\mathcal{R}(W) = \bigcap_{\Lambda} \mathcal{R}_\Lambda(W). \quad (8.4)$$

Note that, because $\{\mathcal{R}_\Lambda(W)\}$ is a decreasing family, we also have

$$\mathcal{R}(W) = \bigcap_{n=1}^{\infty} \mathcal{R}_{\Lambda_n}(W) \quad (8.5)$$

for any increasing sequence $\Lambda_1 \subseteq \Lambda_2 \subseteq \dots$ whose union is all of X .

Remark. If $W(x, x) < 1$ for all $x \in X$, then it follows from Proposition 2.16 that there exist constants $C_x < \infty$ such that $\mathcal{R}(W) \subseteq \prod_{x \in X} [0, C_x]$, so that $\mathcal{R}(W)$ is relatively compact in the product topology.

In view of the fact that each set $\mathcal{R}_\Lambda(W)$ is open, it is perhaps surprising that the limiting set $\mathcal{R}(W)$ is “almost” closed:

Theorem 8.1 *Let X be countably infinite, and let $\mathbf{R} \in \overline{\mathcal{R}(W)}$ (where the closure is taken in the product topology). If every component of $G_W[\text{supp } \mathbf{R}]$ is infinite, then $\mathbf{R} \in \mathcal{R}(W)$.*

Corollary 8.2 *Let X be countably infinite, let $\mathbf{r} \in [0, \infty)^X$, and suppose that every component of $G_W[\text{supp } \mathbf{r}]$ is infinite. Then $\{\lambda \geq 0: \lambda \mathbf{r} \in \mathcal{R}(W)\}$ is a closed interval of $[0, \infty)$ (which may reduce to $\{0\}$ or to all of $[0, \infty)$).*

Remark. If $W(x, y) < 1$ for some $x, y \in \text{supp } \mathbf{r}$, then it follows from Proposition 2.16 that the interval $\{\lambda \geq 0: \lambda \mathbf{r} \in \mathcal{R}(W)\}$ is bounded, i.e. not all of $[0, \infty)$.

PROOF OF THEOREM 8.1. By hypothesis we have

$$\mathbf{R} \in \overline{\bigcap_{\Lambda} \mathcal{R}_{\Lambda}(W)} \subseteq \bigcap_{\Lambda} \overline{\mathcal{R}_{\Lambda}(W)}. \quad (8.6)$$

If $\mathbf{R} \in \mathcal{R}_{\Lambda}(W)$ for all Λ , we are done. So assume that $\mathbf{R} \in \partial \mathcal{R}_{\Lambda}(W)$ for some Λ , i.e. $\mathbf{R} \upharpoonright \Lambda \in \partial \mathcal{R}(W \upharpoonright \Lambda)$ and in particular $Z_{\Lambda}(-\mathbf{R}) = 0$ by Proposition 2.17(b). Let $\Lambda_1, \dots, \Lambda_k$ be the vertex sets of the components of $G_W[\text{supp } \mathbf{R} \cap \Lambda]$. Since $Z_{\Lambda}(\mathbf{w}) = \prod_{i=1}^k Z_{\Lambda_i}(\mathbf{w})$ whenever $\text{supp } \mathbf{w} \subseteq \text{supp } \mathbf{R}$, there must be at least one index i for which $Z_{\Lambda_i}(-\mathbf{R}) = 0$, so that $\mathbf{R} \upharpoonright \Lambda_i \in \partial \mathcal{R}(W \upharpoonright \Lambda_i)$ [recalling that $\mathbf{R} \upharpoonright \Lambda_i \in \overline{\mathcal{R}(W \upharpoonright \Lambda_i)}$ by (8.6)]. Now choose a vertex $x \in (\text{supp } \mathbf{R}) \setminus \Lambda_i$ that is adjacent to Λ_i in G_W [this is possible since each component of $G_W[\text{supp } \mathbf{R}]$ is infinite]; and define $\tilde{\Lambda}_i = \Lambda_i \cup \{x\}$. By (8.6) we have $\mathbf{R} \upharpoonright \tilde{\Lambda}_i \in \overline{\mathcal{R}(W \upharpoonright \tilde{\Lambda}_i)}$; and hence by Corollary 2.19 we have $(\mathbf{R} \upharpoonright \Lambda_i, 0) \in \mathcal{R}(W \upharpoonright \tilde{\Lambda}_i)$ [where 0 corresponds to the x th entry], i.e. $\mathbf{R} \upharpoonright \Lambda_i \in \mathcal{R}(W \upharpoonright \Lambda_i)$. But this contradicts $\mathbf{R} \upharpoonright \Lambda_i \in \partial \mathcal{R}(W \upharpoonright \Lambda_i)$ since $\mathcal{R}(W \upharpoonright \Lambda_i)$ is open. ■

8.2 Infinite trees

In Example 3.6 we considered the complete r -ary rooted tree and showed, following Shearer [98], that there are negative real roots of the univariate polynomial $Z_G(w)$ that tend to $w_{\infty} = -r^r/(r+1)^{r+1}$ as the number of levels tends to infinity. Let us now use the ‘‘homogenization’’ ideas of Section 3.5 to extend this result to trees that are not complete.

Proposition 8.3 *Let G be an infinite tree with (arbitrarily chosen) root vertex x_0 , and let $X_i = \{x: \text{dist}(x, x_0) = i\}$. Let $\bar{b} = \limsup_{i \rightarrow \infty} |X_i|^{1/i}$. If $R > \bar{b}^{\bar{b}}/(\bar{b}+1)^{\bar{b}+1}$, then*

any vector $\mathbf{R} \geq 0$ satisfying $\left(\prod_{x \in X_i} R_x\right)^{1/|X_i|} \geq R$ for all i does not lie in $\mathcal{R}(G)$. In particular, $R\mathbf{1} \notin \mathcal{R}(G)$.

PROOF. Let $\mathbf{R} \geq \mathbf{0}$ be a vector satisfying $\tilde{R}_i \equiv \left(\prod_{x \in X_i} R_x\right)^{1/|X_i|} \geq R$ for all i , and suppose that $\mathbf{R} \in \mathcal{R}(G)$. Then by definition we have $\mathbf{R} \in \mathcal{R}(G_D)$ for every D , where $G_D = G[\bigcup_{i \leq D} X_i]$ is the subtree consisting of the first $D+1$ levels of G .

Let us fix D , and apply Algorithm T of Section 3.4 to the tree G_D with $\mathbf{w} = -\mathbf{R}$. For $0 \leq i \leq D$, define \tilde{p}_i as in (3.52). Let $b_i = |X_{i+1}|/|X_i|$ and define \hat{p}_i by the recursion (3.57). Then it follows from Proposition 3.3 [or the discussion after (3.57)] that $\tilde{p}_i \geq \hat{p}_i$ for $0 \leq i \leq D$, since by hypothesis $\mathbf{R} \in \mathcal{R}(G_D)$. By monotonicity of (3.57) in the \tilde{R}_i , it follows that the sequence $(q_i^{(D)})_{i=1}^{\infty}$ defined by the recursion

$$q_i^{(D)} = \frac{R}{(1 - q_{i+1}^{(D)})^{b_i}}, \quad (8.7)$$

with initial condition $q_i^{(D)} = 0$ for $i \geq D + 1$ satisfies $0 \leq q_i^{(D)} \leq \tilde{p}_i$ for $0 \leq i \leq D$. In particular, we have $0 \leq q_i^{(D)} < 1$ for every i .

Now let $D \rightarrow \infty$. Monotonicity of (8.7) in $q_{i+1}^{(D)}$ implies that, for each i , $q_i^{(D)}$ is increasing in D . Since $q_i^{(D)}$ is bounded above by 1, taking $q_i = \lim_{D \rightarrow \infty} q_i^{(D)}$ gives a sequence $(q_i)_{i=0}^\infty$ satisfying

$$q_i = \frac{R}{(1 - q_{i+1})^{b_i}} \quad (8.8)$$

for every i . Furthermore, $0 \leq q_i \leq 1$ for every i , and so (8.8) implies that $0 \leq q_i < 1$ for $i \geq 1$.

The proposition is then a consequence of the following lemma. ■

Lemma 8.4 *Let $(b_i)_{i=1}^\infty$ be a sequence of positive real numbers, let $\bar{b} = \limsup_{i \rightarrow \infty} (b_1 b_2 \cdots b_{i-1})^{1/i}$, and let $R \geq 0$. Suppose that there exists a sequence $(q_i)_{i=1}^\infty$ satisfying $0 \leq q_i < 1$ and*

$$q_i \geq \frac{R}{(1 - q_{i+1})^{b_i}} \quad (8.9)$$

for all $i \geq 1$. Then $R \leq \bar{b}^{\bar{b}} / (\bar{b} + 1)^{\bar{b}+1}$.

As preparation for proving Lemma 8.4, let us prove an analogous result for finite sequences.

Lemma 8.5 *Let $k \geq 1$, let $b_0, b_1, \dots, b_{k-1} > 0$, and define $\bar{b} = \left(\prod_{i=0}^k b_i \right)^{1/k}$. Let $R \geq 0$. Suppose that there exist $q_0, q_1, \dots, q_k \in [0, 1)$ satisfying*

$$q_i \geq \frac{R}{(1 - q_{i+1})^{b_i}} \quad \text{for } i = 0, 1, \dots, k-1 \quad (8.10)$$

and

$$q_0 \leq q_k. \quad (8.11)$$

Then $R \leq \bar{b}^{\bar{b}} / (\bar{b} + 1)^{\bar{b}+1}$.

PROOF. When $k = 1$, the lemma is proved by straightforward calculus. So let us treat the case $k > 1$. Define weights

$$\gamma_i = \left(\prod_{j=0}^{i-1} b_j \right) / \bar{b}^i \quad (8.12)$$

for $i = 0, 1, \dots, k$ and note that $\gamma_0 = \gamma_k = 1$. Let $\Gamma = \sum_{i=1}^k \gamma_i$. Now define

$$\tilde{q}_0 = \left(\prod_{i=0}^{k-1} q_i^{\gamma_i} \right)^{1/\Gamma} \quad (8.13a)$$

$$\tilde{q}_1 = \left(\prod_{i=1}^k q_i^{\gamma_i} \right)^{1/\Gamma} \quad (8.13b)$$

The argument at (3.65)–(3.73) shows that

$$\tilde{q}_0 \geq \frac{R}{(1 - \tilde{q}_1)^{\bar{b}}}. \quad (8.14)$$

If $q_0 \leq q_k$, then $\tilde{q}_0 \leq \tilde{q}_1$; it then follows from the $k = 1$ case of the lemma that $R \leq \bar{b}^{\bar{b}}/(\bar{b} + 1)^{\bar{b}+1}$. ■

PROOF OF LEMMA 8.4. If $R > \bar{b}^{\bar{b}}/(\bar{b} + 1)^{\bar{b}+1}$, let us choose $\epsilon > 0$ and $\bar{c} < \bar{b}$ such that

$$\frac{R}{1 + \epsilon} > \frac{\bar{c}^{\bar{c}}}{(\bar{c} + 1)^{\bar{c}+1}} > \frac{\bar{b}^{\bar{b}}}{(\bar{b} + 1)^{\bar{b}+1}}. \quad (8.15)$$

Since $\limsup_{i \rightarrow \infty} (b_1 b_2 \cdots b_{i-1})^{1/i} > \bar{c}$, we can find integers $1 < j_1 < j_2 < \dots$ with

$$\prod_{i=j_t+1}^{j_{t+1}} b_i > \bar{c}^{j_{t+1}-j_t} \quad (8.16)$$

for $t = 1, 2, \dots$. Since the sequence $(q_{j_t})_{t=1}^{\infty}$ lies in the interval $[R, 1)$ and $R > 0$, there must exist t such that $q_{j_t} \leq (1 + \epsilon)q_{j_{t+1}}$. To simplify the notation, let us set $k = j_{t+1} - j_t$ and $\hat{q}_i = q_{j_t+i}$ for $i = 0, \dots, k$. By hypothesis we have

$$\hat{q}_i \geq \frac{R}{(1 - \hat{q}_{i+1})^{b_i}} \quad \text{for } i = 0, 1, \dots, k-1 \quad (8.17)$$

and $\hat{q}_0 \leq (1 + \epsilon)\hat{q}_k$. It follows that if we set $S = R/(1 + \epsilon)$, $\hat{q}_i = \hat{q}_i$ for $i = 1, \dots, k$, and $\hat{\hat{q}}_0 = \hat{q}_0/(1 + \epsilon)$, we have

$$\hat{\hat{q}}_i \geq \frac{S}{(1 - \hat{\hat{q}}_{i+1})^{b_i}} \quad \text{for } i = 0, 1, \dots, k-1 \quad (8.18)$$

and $\hat{\hat{q}}_0 \leq \hat{\hat{q}}_k$. Lemma 8.5 together with (8.16) then imply that $S \leq \bar{c}^{\bar{c}}/(\bar{c} + 1)^{\bar{c}+1}$, which is a contradiction. ■

8.3 Regular lattices

The most important situation in statistical mechanics is that of a model defined on a regular lattice (which we take for simplicity to be \mathbb{Z}^d) with a translation-invariant interaction [100]. In this case one expects to be able to prove that the free energy per site (or “pressure”) $F_\Lambda \equiv |\Lambda|^{-1} \log Z_\Lambda$ converges to an infinite-volume limit

$$F_\infty \equiv \lim_{\Lambda \nearrow \infty} F_\Lambda, \quad (8.19)$$

where $\Lambda \nearrow \infty$ denotes convergence in the Følner–van Hove sense, i.e. $|\Lambda| \rightarrow \infty$ in such a way that the surface-to-volume ratio tends to zero (see [113, Section 2.4.1 and Appendix A.3.1] for a variety of equivalent conditions). Indeed, there are several standard arguments for proving such convergence:

- 1) *Almost additivity.* A large volume Λ is subdivided into smaller (but still large) cubes Λ_i separated by wide “corridors”, and $|\log Z_\Lambda - \sum_i \log Z_{\Lambda_i}|$ is shown to be suitably small [67, Section I.2]. This method applies to all models with bounded interaction energies, and yields Følner–van Hove convergence [67, Theorems I.2.3–I.2.5].
- 2) *Superadditivity.* In some cases it can be shown that $\log Z_{\Lambda \cup \Lambda'} \geq \log Z_\Lambda + \log Z_{\Lambda'}$ whenever Λ and Λ' are disjoint. This occurs, in particular, for systems with certain symmetries, for ferromagnets, and for systems with negative interaction energies. Moreover, arbitrary models with bounded interaction energies can be reduced to the case of negative interaction energies simply by adding suitable constants to the interaction terms [100, Theorem II.2.4 and Examples 1–4 following it; see also Section II.4]. However, this method does not yield Følner–van Hove convergence, but only a slightly weaker convergence in which $|\Lambda|/\text{diam}(\Lambda)^d$ must be bounded below away from zero [113, Appendix A.3.3]. Moreover, the limiting free energy F_∞ could be $+\infty$; a separate argument is needed to exclude this possibility.
- 3) *Subadditivity.* In some cases it can be shown that $\log Z_{\Lambda \cup \Lambda'} \leq \log Z_\Lambda + \log Z_{\Lambda'}$ whenever Λ and Λ' are disjoint. This occurs, in particular, for systems with positive interaction energies. Moreover, arbitrary models with bounded interaction energies can be reduced to the case of positive interaction energies by adding suitable constants to the interaction terms. Like superadditivity, this method does not yield Følner–van Hove convergence, but only the slightly weaker convergence in which $|\Lambda|/\text{diam}(\Lambda)^d$ is bounded below away from zero. Moreover, the limiting free energy F_∞ could be $-\infty$; a separate argument is needed to exclude this possibility.

Remark. Although we shall concentrate here on the case of a regular lattice (namely, \mathbb{Z}^d), arguments of the foregoing types can usually be generalized to handle arbitrary quasi-transitive amenable infinite graphs [69, 91]; we expect that the same should be true for our results.

Let us therefore consider the lattice gas on $X = \mathbb{Z}^d$ with a translation-invariant interaction $W(x, y) = W(x - y)$ satisfying $0 \leq W(x, y) \leq 1$. Let us place the same fugacity $w \geq 0$ at each site. [More generally, we could consider a periodic or quasiperiodic fugacity $\mathbf{w} = \{w_x\}_{x \in \mathbb{Z}^d}$; but let us stick to a constant fugacity for simplicity.] Under these assumptions we can prove that the infinite-volume limit of the free energy exists. Indeed, the subadditivity argument applies immediately to our model, thanks to the hypothesis of *repulsive* interactions; moreover, since $Z_\Lambda \geq 1$ for all Λ (thanks to the contribution of the empty configuration), it follows that $F_\infty \geq 0$ and in particular that $F_\infty \neq -\infty$. But this method does not yield Følner–van Hove convergence.

To prove Følner–van Hove convergence, we use the almost-additivity argument. The standard theorems [67, Theorems I.2.3–I.2.5] do not apply to our model, because the interaction energies can be unbounded, indeed for either of two reasons:

- (a) the infinite interaction energy when $W(x, y) = 0$ for some $x \neq y$; and/or
- (b) the unboundedness of the interaction energies in the absence of hard-core self-repulsion [i.e. for $W(x, x) \neq 0$], arising from the fact that the occupation number n_x at a site can be arbitrary large.

We can nevertheless make slight adaptations in the standard almost-additivity argument so as to make the proof go through. We assume that the total interaction of any site with the rest of the world is finite, i.e.

$$\sum_{y \in \mathbb{Z}^d} [1 - W(x, y)] < \infty. \quad (8.20)$$

In particular, there is a number $R < \infty$ such that $W(x, y) > 0$ whenever $|x - y| > R$.

Let us begin with the case of hard-core self-repulsion, so that we need only deal with problem (a). This is handled by the following lemma:

Lemma 8.6 *Consider any repulsive lattice gas with hard-core self-repulsion and fugacities $w_x \geq 0$. Suppose that $\Lambda = \bigcup_{i=1}^n \Lambda_i \cup \Lambda_0$ (disjoint union). Then*

$$\left(\prod_{i=1}^n Z_{\Lambda_i} \right) \left(\prod_{\substack{1 \leq i < j \leq n \\ x \in \Lambda_i \\ y \in \Lambda_j}} W(x, y) \right) \leq Z_{\Lambda} \leq \left(\prod_{i=1}^n Z_{\Lambda_i} \right) \left(\prod_{x \in \Lambda_0} (1 + w_x) \right). \quad (8.21)$$

PROOF. We make use of the definition

$$Z_{\Lambda} = \sum_{X' \subseteq \Lambda} \prod_{x \in X'} w_x \prod_{\{x, y\} \subseteq X'} W(x, y). \quad (8.22)$$

The lower bound on Z_{Λ} is obtained by considering only those configurations in which Λ_0 is empty (i.e. $X' \cap \Lambda_0 = \emptyset$) and writing

$$\prod_{\{x, y\} \subseteq X'} W(x, y) \geq \left(\prod_{i=1}^n \prod_{\{x, y\} \subseteq X' \cap \Lambda_i} W(x, y) \right) \left(\prod_{\substack{1 \leq i < j \leq n \\ x \in \Lambda_i \\ y \in \Lambda_j}} W(x, y) \right), \quad (8.23)$$

which is valid since $0 \leq W(x, y) \leq 1$. The upper bound on Z_{Λ} is obtained by writing

$$\prod_{\{x, y\} \subseteq X'} W(x, y) \leq \prod_{i=1}^n \prod_{\{x, y\} \subseteq X' \cap \Lambda_i} W(x, y). \quad (8.24)$$

■

We then have:

Theorem 8.7 (infinite-volume limit, hard-core case) *Consider a translation-invariant repulsive lattice gas on \mathbb{Z}^d , with hard-core self-repulsion, satisfying $\sum_{y \in \mathbb{Z}^d} [1 - W(x, y)] < \infty$, with the same fugacity $w \geq 0$ at each site. Then $F_\infty(w) \equiv \lim_{\Lambda \nearrow \infty} |\Lambda|^{-1} \log Z_\Lambda(w)$ exists in Følner–van Hove sense and satisfies $0 \leq F_\infty(w) \leq \log(1 + w)$. Moreover, $F_\infty(w)$ is an increasing and convex function of $\log w$; in particular, it is continuous on $0 \leq w < \infty$.*

PROOF. Fix integers $a, c > 0$ and consider the paving of \mathbb{Z}^d by disjoint cubes of side $a + c$ with corners located at $(a + c)\mathbb{Z}^d$, i.e. cubes $C_{\mathbf{n}} = [0, a + c)^d + (a + c)\mathbf{n}$ for $\mathbf{n} \in \mathbb{Z}^d$. Consider also the subcubes $C'_{\mathbf{n}} \subset C_{\mathbf{n}}$ of side a with the same lowermost corner, i.e. $C'_{\mathbf{n}} = [0, a)^d + (a + c)\mathbf{n}$. For any finite subset $\Lambda \subset \mathbb{Z}^d$, let \mathcal{C}_Λ be the collection of all cubes $C_{\mathbf{n}}$ that are contained in Λ , and let $\mathcal{C}'_\Lambda = \{C'_{\mathbf{n}} : C_{\mathbf{n}} \in \mathcal{C}_\Lambda\}$. By Lemma 8.6,

$$\left| \log Z_\Lambda - \sum_{C' \in \mathcal{C}'_\Lambda} \log Z_{C'} \right| \leq \alpha(c) |\Lambda| + \beta \left| \Lambda \setminus \bigcup_{C' \in \mathcal{C}'_\Lambda} C' \right|, \quad (8.25)$$

where $\alpha(c) = - \sum_{\substack{x \in \mathbb{Z}^d \\ |x| > c}} \log W(x)$ and $\beta = \log(1 + w)$. By translation invariance,

$\log Z_{C'} = \log Z_{C'_0}$ for all $C' \in \mathcal{C}'_\Lambda$. Moreover, $|\mathcal{C}'_\Lambda| = |\mathcal{C}_\Lambda|$ by construction. Dividing (8.25) by $|\Lambda|$, we get

$$\left| \frac{1}{|\Lambda|} \log Z_\Lambda - \frac{a^d |\mathcal{C}'_\Lambda|}{|\Lambda|} \frac{1}{a^d} \log Z_{C'_0} \right| \leq \alpha(c) + \beta \left(1 - \frac{a^d |\mathcal{C}_\Lambda|}{|\Lambda|} \right) \quad (8.26)$$

and hence

$$|F_\Lambda - F_{C'_0}| \leq \alpha(c) + \left(1 - \frac{a^d |\mathcal{C}_\Lambda|}{|\Lambda|} \right) \left(\beta + \frac{1}{a^d} |\log Z_{C'_0}| \right) \quad (8.27a)$$

$$\leq \alpha(c) + 2\beta \left(1 - \frac{a^d |\mathcal{C}_\Lambda|}{|\Lambda|} \right) \quad (8.27b)$$

since $1 \leq Z_{C'_0} \leq (1 + w)^{a^d}$. Taking (8.27b) for two finite subsets $\Lambda, \tilde{\Lambda} \subset \mathbb{Z}^d$ and subtracting, we get

$$|F_\Lambda - F_{\tilde{\Lambda}}| \leq 2\alpha(c) + 2\beta \left(1 - \frac{a^d |\mathcal{C}_\Lambda|}{|\Lambda|} \right) + 2\beta \left(1 - \frac{a^d |\mathcal{C}_{\tilde{\Lambda}}|}{|\tilde{\Lambda}|} \right). \quad (8.28)$$

Now let $(\Lambda_n) \nearrow \infty$ in Følner–van Hove sense; this implies that the fraction of volume of Λ_n contained in the cubes \mathcal{C}_{Λ_n} tends to 1, i.e. $\lim_{n \rightarrow \infty} (a + c)^d |\mathcal{C}_{\Lambda_n}| / |\Lambda_n| = 1$. Therefore,

$$\limsup_{m, n \rightarrow \infty} |F_{\Lambda_m} - F_{\Lambda_n}| \leq 2\alpha(c) + 4\beta \left[1 - \left(\frac{a}{a + c} \right) d \right]. \quad (8.29)$$

Now, given any $\epsilon > 0$, we can [by (8.20)] choose c large enough so that $\alpha(c) \leq \epsilon$; and we can then choose a large enough so that $1 - [a/(a+c)]^d \leq \epsilon$. Hence

$$\limsup_{m,n \rightarrow \infty} |F_{\Lambda_m} - F_{\Lambda_n}| \leq 2\epsilon + 4\beta\epsilon. \quad (8.30)$$

Since ϵ is arbitrary, we have $\limsup_{m,n \rightarrow \infty} |F_{\Lambda_m} - F_{\Lambda_n}| = 0$, so that $(|\Lambda_n|^{-1} \log Z_{\Lambda_n})_{n=1}^{\infty}$ is a Cauchy sequence and hence converges. (The limit is the same for all Følner–van Hove sequences, since interleaving two Følner–van Hove sequences yields another one.)

The bounds $0 \leq F_{\infty}(w) \leq \log(1+w)$ follow immediately from $1 \leq Z_{\Lambda} \leq (1+w)^{|\Lambda|}$. Finally, each $F_{\Lambda}(w)$ is manifestly increasing, and by Lemma 2.32 it is a convex function of $\log w$; and these properties are preserved under pointwise limits. ■

Now let us turn to problem (b), arising from the unboundedness of the occupation number n_x when $W(x,x) \neq 0$. We begin with a well-known fact about positive correlation of increasing functions on a totally ordered space:

Lemma 8.8 *Let $\mu \neq 0$ be a nonnegative measure on a totally ordered space Ω , and let f and g be increasing functions on Ω . Then (provided the functions concerned are integrable):*

- (a) $(\int fg d\mu) (\int \mathbf{1} d\mu) \geq (\int f d\mu) (\int g d\mu)$.
- (b) $\frac{\int f e^{\alpha g} d\mu}{\int e^{\alpha g} d\mu}$ is an increasing function of $\alpha \in \mathbb{R}$.

PROOF. (a) We have

$$(\int fg d\mu) (\int \mathbf{1} d\mu) - (\int f d\mu) (\int g d\mu) = \frac{1}{2} \int [f(x) - f(y)] [g(x) - g(y)] d\mu(x) d\mu(y), \quad (8.31)$$

and the integrand on the right-hand side is nonnegative both when $x \leq y$ and when $x \geq y$.

(b) We have

$$\frac{d}{d\alpha} \left(\frac{\int f e^{\alpha g} d\mu}{\int e^{\alpha g} d\mu} \right) = \frac{(\int f g e^{\alpha g} d\mu) (\int e^{\alpha g} d\mu) - (\int f e^{\alpha g} d\mu) (\int g e^{\alpha g} d\mu)}{(\int e^{\alpha g} d\mu)^2}. \quad (8.32)$$

Now apply part (a) with μ replaced by $e^{\alpha g} \mu$. ■

Now let Λ be a finite set; for each $x \in \Lambda$, let Ω_x be a totally ordered space; and let $\Omega = \prod_{x \in \Lambda} \Omega_x$. For each $x \in \Lambda$, let $\mu_x \neq 0$ be a nonnegative measure on Ω_x and let F_x be a nonnegative *decreasing* function on Ω_x . Finally, let H be an increasing function on Ω and let $\alpha \geq 0$. Define

$$Z_{\Lambda}^{\{\{F_x\}, \alpha, H\}} = \int \left(\prod_{x \in \Lambda} F_x(\varphi_x) \right) \exp[-\alpha H(\varphi)] \prod_{x \in \Lambda} d\mu_x(\varphi_x). \quad (8.33)$$

Lemma 8.9 *Under the above hypotheses,*

$$Z_{\Lambda}^{\{\{F_x\}, \alpha, H\}} \geq \left(\prod_{x \in \Lambda} \frac{\int F_x d\mu_x}{\int \mathbf{1} d\mu_x} \right) \times Z_{\Lambda}^{\{\{\mathbf{1}\}, \alpha, H\}}. \quad (8.34)$$

PROOF. Choose one site $z \in \Lambda$, and let us study the integral over φ_z with $\varphi_{\neq z}$ held fixed. By Lemma 8.8(b), the quantity

$$\frac{\int F_z(\varphi_z) \exp[-\alpha H(\varphi_z, \varphi_{\neq z})] d\mu_z(\varphi_z)}{\int \exp[-\alpha H(\varphi_z, \varphi_{\neq z})] d\mu_z(\varphi_z)} \quad (8.35)$$

is an increasing function of $\alpha \in \mathbb{R}$, so its value at $\alpha \geq 0$ is bounded below by its value at $\alpha = 0$, which is

$$\frac{\int F_z(\varphi_z) d\mu_z(\varphi_z)}{\int d\mu_z(\varphi_z)}. \quad (8.36)$$

Therefore,

$$\int F_z(\varphi_z) \exp[-\alpha H(\varphi_z, \varphi_{\neq z})] d\mu_z(\varphi_z) \geq \left(\frac{\int F_z d\mu_z}{\int \mathbf{1} d\mu_z} \right) \times \int \exp[-\alpha H(\varphi_z, \varphi_{\neq z})] d\mu_z(\varphi_z). \quad (8.37)$$

Now multiply both sides by $\prod_{x \in \Lambda \setminus z} F_x(\varphi_x)$ and integrate with respect to $\prod_{x \in \Lambda \setminus z} d\mu_x(\varphi_x)$; we obtain

$$Z_{\Lambda}^{\{\{F_x\}, \alpha, H\}} \geq \left(\frac{\int F_z d\mu_z}{\int \mathbf{1} d\mu_z} \right) \times Z_{\Lambda}^{\{\{\mathbf{1}_x, F_{\neq x}\}, \alpha, H\}}. \quad (8.38)$$

Applying the same argument successively to each site in Λ , we obtain (8.34). ■

Let us now specialize these results to the repulsive lattice gas, by taking $\Omega_x = \mathbb{N}$, $d\mu_x(n_x) = w_x^{n_x} / n_x!$,

$$H(\mathbf{n}) = \sum_{x \in \Lambda} [-\log W(x, x)] \frac{n_x(n_x - 1)}{2} + \sum_{\{x, y\} \subseteq \Lambda} [-\log W(x, y)] n_x n_y, \quad (8.39)$$

and $F_x(n_x) = \mathbf{1}(n_x \leq K_x)$ for arbitrarily chosen positive constants K_x . We then have

$$\frac{Z_{\Lambda}^{(\mathbf{K})}}{Z_{\Lambda}} \geq \prod_{x \in \Lambda} [1 - \gamma(K_x, w_x)], \quad (8.40)$$

where $Z_{\Lambda}^{(\mathbf{K})}$ denotes the sum (1.1b) restricted to the configurations satisfying $n_x \leq K_x$ for all x , and

$$\gamma(K, w) = e^{-w} \sum_{n=K}^{\infty} \frac{w^n}{n!}. \quad (8.41)$$

We can now prove a variant of Lemma 8.6:

Lemma 8.10 Consider any repulsive lattice gas with fugacities $w_x \geq 0$, and let $(K_x)_{x \in \Lambda}$ be arbitrary positive constants. Suppose that $\Lambda = \bigcup_{i=1}^n \Lambda_i \cup \Lambda_0$ (disjoint union).

Then

$$\begin{aligned} \left(\prod_{i=1}^n Z_{\Lambda_i} \right) \left(\prod_{\substack{1 \leq i \leq n \\ x \in \Lambda_i}} [1 - \gamma(K_x, w_x)] \right) & \left(\prod_{\substack{1 \leq i < j \leq n \\ x \in \Lambda_i \\ y \in \Lambda_j}} W(x, y)^{K_x K_y} \right) \\ & \leq Z_{\Lambda} \leq \left(\prod_{i=1}^n Z_{\Lambda_i} \right) \left(\prod_{x \in \Lambda_0} e^{w_x} \right). \end{aligned} \quad (8.42)$$

PROOF. The upper bound is proved exactly as in Lemma 8.6. For the lower bound, we begin from the trivial fact that $Z_{\Lambda} \geq Z_{\Lambda}^{(\mathbf{K})}$. Now consider only those configurations satisfying $\mathbf{n} \leq \mathbf{K}$ and, in addition, $n_x = 0$ for all $x \in \Lambda_0$; for such configurations we have

$$\prod_{\{x, y\} \subseteq \Lambda \setminus \Lambda_0} W(x, y)^{n_x n_y} \geq \left(\prod_{i=1}^n \prod_{\{x, y\} \subseteq \Lambda_i} W(x, y)^{n_x n_y} \right) \left(\prod_{1 \leq i < j \leq n} \prod_{\substack{x \in \Lambda_i \\ y \in \Lambda_j}} W(x, y)^{K_x K_y} \right). \quad (8.43)$$

It follows that

$$Z_{\Lambda}^{(\mathbf{K})} \geq \left(\prod_{i=1}^n Z_{\Lambda_i}^{(\mathbf{K}|_{\Lambda_i})} \right) \left(\prod_{1 \leq i < j \leq n} \prod_{\substack{x \in \Lambda_i \\ y \in \Lambda_j}} W(x, y)^{K_x K_y} \right). \quad (8.44)$$

Now use (8.40) for each set Λ_i . ■

Theorem 8.11 (infinite-volume limit, general case) Consider a translation-invariant repulsive lattice gas on \mathbb{Z}^d , satisfying $\sum_{y \in \mathbb{Z}^d} [1 - W(x, y)] < \infty$, with the same fugacity $w \geq 0$ at each site. Then $F_{\infty}(w) \equiv \lim_{\Lambda \nearrow \infty} |\Lambda|^{-1} \log Z_{\Lambda}(w)$ exists in Følner–van Hove sense and satisfies $0 \leq F_{\infty}(w) \leq w$. Moreover, $F_{\infty}(w)$ is an increasing and convex function of $\log w$; in particular, it is continuous on $0 \leq w < \infty$.

PROOF. We begin by defining families of cubes \mathcal{C}_{Λ} and \mathcal{C}'_{Λ} as in the proof of Theorem 8.7. By Lemma 8.10 with $K_x = K$ for all x , we have

$$\left| \log Z_{\Lambda} - \sum_{C' \in \mathcal{C}'_{\Lambda}} \log Z_{C'} \right| \leq \alpha(c) K^2 |\Lambda| + w \left| \Lambda \setminus \bigcup_{C' \in \mathcal{C}'_{\Lambda}} C' \right| + |\Lambda| |\log[1 - \gamma(K, w)]| \quad (8.45)$$

where $\alpha(c) = - \sum_{\substack{x \in \mathbb{Z}^d \\ |x| > c}} \log W(x)$. By the same arguments as before, we have

$$|F_\Lambda - F_{C'_0}| \leq \alpha(c)K^2 + 2w \left(1 - \frac{a^d |C_\Lambda|}{|\Lambda|}\right) + |\log[1 - \gamma(K, w)]|. \quad (8.46)$$

Taking (8.46) for two finite subsets $\Lambda, \tilde{\Lambda} \subset \mathbb{Z}^d$ and subtracting, we get

$$|F_\Lambda - F_{\tilde{\Lambda}}| \leq 2\alpha(c)K^2 + 2w \left(1 - \frac{a^d |C_\Lambda|}{|\Lambda|}\right) + 2w \left(1 - \frac{a^d |C_{\tilde{\Lambda}}|}{|\tilde{\Lambda}|}\right) + 2|\log[1 - \gamma(K, w)]|. \quad (8.47)$$

Now let $(\Lambda_n) \nearrow \infty$ in Følner–van Hove sense, so that $\lim_{n \rightarrow \infty} (a+c)^d |C_{\Lambda_n}| / |\Lambda_n| = 1$. It follows that

$$\limsup_{m, n \rightarrow \infty} |F_{\Lambda_m} - F_{\Lambda_n}| \leq 2\alpha(c)K^2 + 4w \left[1 - \left(\frac{a}{a+c}\right)^d\right] + 2|\log[1 - \gamma(K, w)]|. \quad (8.48)$$

Now, given any $\epsilon > 0$, we first choose K large enough so that $|\log[1 - \gamma(K, w)]| \leq \epsilon$; then we choose c large enough so that $\alpha(c)K^2 \leq \epsilon$; and finally we choose a large enough so that $1 - [a/(a+c)]^d \leq \epsilon$. Therefore,

$$\limsup_{m, n \rightarrow \infty} |F_{\Lambda_m} - F_{\Lambda_n}| \leq 4\epsilon + 4\epsilon w. \quad (8.49)$$

The remainder of the argument is exactly as in Theorem 8.7. ■

Historical remark. Results like Theorem 8.11 have been proven in the vastly more general context of “unbounded spin systems” by Lebowitz and Presutti [75], based on superstability estimates due to Ruelle [93]. See also [74, 9] for related work. We think that it is nevertheless useful to give an elementary and self-contained proof for our special case.

Let us now return to the case of hard-core self-repulsion, and consider the convergence of the finite-volume free energies $F_\Lambda(w)$ for *complex* fugacities w . By hypothesis (8.20) we have

$$\Delta_W \equiv \sum_{y \neq 0} [1 - W(0, y)] < \infty. \quad (8.50)$$

Therefore, by Corollary 5.2, all of the partition functions $Z_\Lambda(w)$ are nonvanishing in the disc $|w| < R(\Delta_W)$ where $R(\Delta_W)$ is defined by (5.12). Moreover, they satisfy the trivial bounds $|Z_\Lambda(w)| \leq e^{|\Lambda||w|}$. It follows that the free energies $F_\Lambda(w) \equiv |\Lambda|^{-1} \log Z_\Lambda(w)$ are analytic in the disc $|w| < R(\Delta_W)$ and satisfy there $\operatorname{Re} F_\Lambda(w) \leq |w| \leq R(\Delta_W)$. Finally, Theorem 8.11 shows that the $F_\Lambda(w)$ converge to a limit when w lies in the *real* interval $[0, R(\Delta_W))$. These three facts are sufficient to imply the convergence of $F_\Lambda(w)$ to an analytic limit $F_\infty(w)$ everywhere in the disc $|w| < R(\Delta_W)$, using the following standard result on normal families of analytic functions:

Proposition 8.12 (exp log Vitali) *Let D be a domain in \mathbb{C} , let S be a subset of D having at least one accumulation point in D , let $M < \infty$, and let $(f_n)_{n=1}^\infty$ be analytic functions in D satisfying:*

- (a) $\operatorname{Re} f_n(z) \leq M$ for all n and all $z \in D$; and
- (b) $\lim_{n \rightarrow \infty} f_n(z)$ exists (and is finite) for all $z \in S$.

Then there exists an analytic function f_∞ on D such that $f_n(z) \rightarrow f_\infty(z)$ uniformly for z in compact subsets of D .

For a proof of Proposition 8.12, see e.g. [99, p. 343].

Corollary 8.13 *Consider a translation-invariant repulsive lattice gas on \mathbb{Z}^d , with hard-core self-repulsion, with*

$$\sum_{y \neq 0} [1 - W(0, y)] \equiv \Delta_W < \infty. \quad (8.51)$$

Then:

- (a) *Each function $F_\Lambda(w) \equiv |\Lambda|^{-1} \log Z_\Lambda(w)$ is analytic in the disc $|w| < R(\Delta_W)$ and satisfies there $\operatorname{Re} F_\Lambda(w) \leq |w|$.*
- (b) *There exists an analytic function $F_\infty(w)$ on the disc $|w| < R(\Delta_W)$ such that $\lim_{\Lambda \nearrow \infty} F_\Lambda(w) = F_\infty(w)$ uniformly for w in compact subsets of $|w| < R(\Delta_W)$.*

Remark. Proposition 8.12 is a very special case of a much more general “Vitali–Porter–type” theorem for normal families of analytic functions [103, Lemma 3.5], in which condition (a) can be weakened to “there exists a nonempty disc $\Delta \subset \mathbb{C}$ such that $f_n(z) \notin \Delta$ for all $z \in D$ ” [96, Example 2.3.9] or even to “there exist $w_1, w_2 \in \mathbb{C}$ with $w_1 \neq w_2$ such that $f_n(z) \notin \{w_1, w_2\}$ for all $z \in D$ ” [96, Section 2.7]. Moreover, it is sufficient for this hypothesis to hold locally in D . Detailed accounts of these results can be found in [84, 96].

8.4 Quantitative bounds for the lattice \mathbb{Z}^d

By Corollary 8.2, the set $\{\lambda \geq 0: \lambda \mathbf{1} \in \mathcal{R}(W)\}$ is a *closed* interval $[0, \lambda_c]$ provided that each component of G_W is infinite. In this section we sketch briefly some methods for finding reasonably sharp upper and lower bounds on λ_c . We shall illustrate these examples with reference to the hard-core lattice gas on \mathbb{Z}^d with nearest-neighbor edges, paying particular attention to the case of the square lattice \mathbb{Z}^2 .

8.4.1 Upper bounds on λ_c

Let G be any finite subgraph of \mathbb{Z}^d , and let $w_\star = -\lambda_\star$ be the negative real root of Z_G of smallest magnitude. Then, by the definition of $\mathcal{R}(W)$ for an infinite graph [cf. (8.4)] together with the monotonicity of $\mathcal{R}(G)$ in G [Proposition 2.15], we can conclude immediately that $\lambda_c(\mathbb{Z}^d) \leq \lambda_\star$. Moreover, these bounds converge to the exact value λ_c if we take any increasing sequence $G_1 \subseteq G_2 \subseteq \dots$ whose union is all of \mathbb{Z}^d .

More generally, as a consequence of the discussion in Section 6.2, we can take any finite subtree T of the pruned SAW-tree for \mathbb{Z}^d , and let $w_\star = -\lambda_\star$ be the negative real root of Z_T of smallest magnitude; we again have $\lambda_c(\mathbb{Z}^d) \leq \lambda_\star$.

Example 8.1. Clearly \mathbb{Z}^d contains finite paths of all lengths. By Example 3.2 (or alternatively Example 3.6 with $r = 1$), these give bounds λ_\star tending to $1/4$ as the path length tends to infinity. Thus $\lambda_c(\mathbb{Z}^d) \leq 1/4$.

Example 8.2. Now consider a finite subgraph $G \subseteq \mathbb{Z}^d$ consisting of a long path P along one axis (the *spine*) and, radiating from each vertex of P , $2d - 2$ disjoint long paths perpendicular to P (the *antennae*). Let us assign fugacity $w \leq 0$ to every vertex of G . Using Algorithm T on the antennae, we must first iterate the recursion $w_{\text{eff}} \mapsto w/(1 + w_{\text{eff}})$. If the antennae were infinitely long, we would approach the fixed point $w' = -\frac{1}{2} + \sqrt{w + \frac{1}{4}}$ [provided that $-1/4 \leq w \leq 0$; otherwise we are outside $\mathcal{R}(G)$]. By taking the antennae sufficiently long, we can get as close to this value as we wish. We are now left with a “caterpillar” consisting of the spine (with fugacity w on each vertex) along with $2d - 2$ pendant vertices (each with fugacity w') attached to each vertex of the spine. Applying Algorithm T to this graph, we get the recursion $w_{\text{eff}} \mapsto w/[(1 + w_{\text{eff}})(1 + w')^{2d-2}] \equiv w_\star/(1 + w_{\text{eff}})$, where

$$w_\star \equiv \frac{w}{(1 + w')^{2d-2}} = \frac{w}{\left(\frac{1}{2} + \sqrt{w + \frac{1}{4}}\right)^{2d-2}}. \quad (8.52)$$

We require $-\frac{1}{4} \leq w_\star \leq 0$ in order to stay within $\mathcal{R}(G)$. For $d = 2$ this yields the bound $\lambda_c(\mathbb{Z}^2) \leq 4/25$. For $d \rightarrow \infty$ it yields $\lambda_c(\mathbb{Z}^d) \leq (\log d)/(2d) + O((\log \log d)/d)$.

Example 8.3. To get the correct bound $\lambda_c(\mathbb{Z}^d) = O(1/d)$, one can argue using a subtree of the pruned SAW-tree. Note first that any walk using only steps in the positive coordinate directions is guaranteed to be self-avoiding. Moreover, if we define the pruning such that at each vertex all steps in the positive coordinate directions are preferred to all steps in the negative coordinate directions, then every walk using only positive coordinate steps appears as a vertex in the pruned SAW-tree of \mathbb{Z}^d . Thus, the complete d -ary rooted tree (consisting of these walks) is a subtree of the pruned SAW-tree of \mathbb{Z}^d , and so by Example 3.6 we have

$$\lambda_c(\mathbb{Z}^d) \leq \frac{d^d}{(d+1)^{d+1}} \sim \frac{1}{ed}. \quad (8.53)$$

Example 8.4. Asymptotically correct upper bounds on $\lambda_c(\mathbb{Z}^d)$ can be obtained by using large cylinders, for which λ_c can be computed by the transfer-matrix method.¹⁵ Let us illustrate the method for \mathbb{Z}^2 . Consider the strip $S_L = \{(x, y) \in \mathbb{Z}^2: 0 \leq x < L\}$. Since $S_1 \subseteq S_2 \subseteq \dots \subseteq \mathbb{Z}^2$, we have $\lambda_c(S_1) \geq \lambda_c(S_2) \geq \dots \geq \lambda_c(\mathbb{Z}^2)$. [In particular, $\lim_{L \rightarrow \infty} \lambda_c(S_L)$ exists.] On the other hand, since each finite subgraph of \mathbb{Z}^2 is contained (modulo translation) in S_L for all sufficiently large L , we have $\lim_{L \rightarrow \infty} \lambda_c(S_L) \leq \lambda_c(\mathbb{Z}^2)$. It follows that $\lambda_c(S_L) \downarrow \lambda_c(\mathbb{Z}^2)$ as $L \rightarrow \infty$.

An analogous argument can also be made using strips with periodic boundary conditions, which are more convenient for computation [110]. To see this, let \tilde{S}_L be the strip S_L with an extra edge added from $(L-1, y)$ to $(0, y)$ for each y . As before, each finite subgraph of \mathbb{Z}^2 is contained (modulo translation) in \tilde{S}_L for all sufficiently large L , so that $\limsup_{L \rightarrow \infty} \lambda_c(\tilde{S}_L) \leq \lambda_c(\mathbb{Z}^2)$. On the other hand, the pruned SAW-tree of \tilde{S}_L (for any given choice of ordering) is a subtree of the pruned SAW-tree of \mathbb{Z}^2 (provided we make an appropriately compatible choice of ordering): just map each path in \tilde{S}_L to its “universal cover” in \mathbb{Z}^2 , i.e. to the path in \mathbb{Z}^2 obtained by making the same sequence of north, south, east or west steps. Thus $\lambda_c(\tilde{S}_L) \geq \lambda_c(\mathbb{Z}^2)$ for every L (though we do not necessarily have monotonicity in L). It follows that $\lambda_c(S_L) \rightarrow \lambda_c(\mathbb{Z}^2)$ as $L \rightarrow \infty$.

Example 8.5. Let us also remark that $\lambda_c(\mathbb{Z}^d)$ is *strictly* decreasing in d ; indeed, we can derive an inequality bounding $\lambda_c(\mathbb{Z}^{d+1})$ above in terms of $\lambda_c(\mathbb{Z}^d)$. To see this, note that \mathbb{Z}^{d+1} contains a copy of the graph G_d obtained from \mathbb{Z}^d by attaching two semi-infinite paths (“antennae”) to every vertex. If we place fugacity $w \leq 0$ on every vertex of G_d and integrate out the antennae as in Example 8.2, we are left with a copy of \mathbb{Z}^d with effective fugacities

$$w_{\text{eff}} = \frac{w}{(1+w')^2} = \frac{w}{w + \frac{1}{2} + \sqrt{w + \frac{1}{4}}}. \quad (8.54)$$

If $w \in (-\lambda_c(\mathbb{Z}^{d+1}), 0]$, then w_{eff} must be in $(-\lambda_c(\mathbb{Z}^d), 0]$. It follows that

$$\lambda_c(\mathbb{Z}^d) \geq \frac{\lambda_c(\mathbb{Z}^{d+1})}{\frac{1}{2} - \lambda_c(\mathbb{Z}^{d+1}) + \sqrt{\frac{1}{4} - \lambda_c(\mathbb{Z}^{d+1})}}, \quad (8.55)$$

or equivalently

$$\lambda_c(\mathbb{Z}^{d+1}) \leq \frac{\lambda_c(\mathbb{Z}^d)}{[1 + \lambda_c(\mathbb{Z}^d)]^2}, \quad (8.56)$$

which is the desired bound.

Using (8.56) together with the initial condition $\lambda_c(\mathbb{Z}^1) = 1/4$, it is easy to show by induction that $\lambda_c(\mathbb{Z}^d) \leq 1/(2d+2)$. However, this bound is less sharp than the bound (8.53) obtained in Example 8.3.

¹⁵See e.g. [110] for a brief discussion of transfer matrices for the hard-core lattice gas.

8.4.2 Lower bounds on λ_c

We now turn to proving lower bounds on λ_c . Corollary 5.3 gives

$$\lambda_c(\mathbb{Z}^d) \geq \frac{\Delta^\Delta}{(\Delta + 1)^{\Delta+1}} = \frac{(2d)^{2d}}{(2d + 1)^{2d+1}} \sim \frac{1}{2ed}, \quad (8.57)$$

while Corollary 5.7 gives the slightly better bound

$$\lambda_c(\mathbb{Z}^2) \geq \frac{(\Delta - 1)^{\Delta-1}}{\Delta^\Delta} = \frac{(2d - 1)^{2d-1}}{(2d)^{2d}} \sim \frac{1}{2ed}. \quad (8.58)$$

For $d = 2$ the latter bound yields $\lambda_c(\mathbb{Z}^2) \geq 27/256$. Either of these bounds shows, when combined with Example 8.3, that $\lambda_c(\mathbb{Z}^d) = \Theta(1/d)$ as $d \rightarrow \infty$.

To get quantitatively better bounds, we can take a supertree of the pruned SAW-tree of \mathbb{Z}^d and calculate exactly for it. In particular, if we choose a supertree that is eventually periodic, the computation reduces to finding the fixed point of a recursion.

Example 8.6. The pruned SAW-tree of \mathbb{Z}^d is obviously contained in the $2d$ -branching tree, for which the computation has been performed in Example 3.6. This yields the bound (8.57). [This argument obviously works for any infinite graph of maximum degree Δ , not just \mathbb{Z}^d .]

Example 8.7. A smaller supertree of the pruned SAW-tree is obtained by giving the root $2d$ children and every other vertex $2d - 1$ children. Applying Algorithm T, the recursion (until we reach children of the root) is $w' \mapsto w/(1 + w')^{\Delta-1}$. We end up with the bound (8.58). [The argument again works for any infinite graph of maximum degree Δ .]

Example 8.8. Another possibility (not the only one) is to take a large finite piece T of the pruned SAW-tree of \mathbb{Z}^2 , and then repeat it periodically (attaching to each leaf of T a copy of T starting at the root, and repeating this). It is easily seen that this gives a supertree of the pruned SAW-tree. We therefore use the recursion $w_{\text{eff,leaves}} \mapsto w_{\text{eff,root}}$ and demand that there exist an attractive fixed point in $(-1, 0]$. We have various choices about how to order the edges to define the pruning. For instance, we can choose an ordering of the edges at each vertex in a translation-invariant way, find the first k levels of the pruned SAW-tree, and then repeat these periodically. Alternatively, we can order the vertices arbitrarily at the root of the SAW-tree, and at all other vertices of the SAW-tree order the edges according to the angle the path turns through (e.g. $0, +\pi/2, -\pi/2$) in taking the step. This defines a pruned SAW-tree, and we can take the first k levels and repeat periodically as before. We conjecture that either method gives bounds converging to $\lambda_c(\mathbb{Z}^2)$ as k grows, but we do not know how to prove this.

Remark. In Examples 8.2 and 8.7, we have proven for \mathbb{Z}^2 the rigorous bounds

$$0.105468\dots = \frac{27}{256} \leq \lambda_c(\mathbb{Z}^2) \leq \frac{4}{25} = 0.16. \quad (8.59)$$

By extensions of those arguments and with a little calculation, these bounds can be narrowed further; it would be interesting to see how far one can go. It is worth noting that Todo [110] has given the extraordinarily precise numerical estimate

$$\lambda_c(\mathbb{Z}^2) = 0.119\,338\,881\,88(1), \quad (8.60)$$

obtained by using transfer matrices and the phenomenological-renormalization method (a variant of finite-size scaling). Furthermore, his computations up to $L = 38$ show [111] that¹⁶

$$\lambda_c(\tilde{S}_{38}) = 0.119\,365(1), \quad (8.61)$$

which by Example 8.4 provides an upper bound on $\lambda_c(\mathbb{Z}^2)$. See also Guttmann [54] for an earlier and only slightly less precise estimate of $\lambda_c(\mathbb{Z}^2)$, obtained by series analysis.

Acknowledgments

We wish to thank Keith Ball for informing us of the important work of Shearer [98], and Joel Lebowitz for informing us of the work of Ginibre [46]. We also thank Keith Ball, Pierre Leroux and Joel Spencer for helpful discussions, and Synge Todo for communicating to us some of his unpublished numerical results.

One of us (Sokal) wishes to thank the Department of Mathematics at University College London and the Theoretical Physics Group at Imperial College for hospitality during his stays in London.

This research was supported in part by U.S. National Science Foundation grants PHY-9900769 and PHY-0099393.

¹⁶What Todo [111] actually computed is the number λ_L^\times for which

- (a) the transfer matrix has a unique eigenvalue of largest modulus for $w \in (-\lambda_L^\times, 0]$, and
- (b) the transfer matrix has two dominant eigenvalues of largest modulus for $w = -\lambda_L^\times$.

Let us show that $\lambda_L^\times = \lambda_c(\tilde{S}_L)$. Let $\tilde{S}_L^{(n)}$ be the cylinder of width L and length n . By monotonicity, $\lambda_c(\tilde{S}_L^{(n)})$ decreases in n and hence has a limit as $n \rightarrow \infty$; it is easy to see that this limit is $\lambda_c(\tilde{S}_L)$. Now, the Beraha–Kahane–Weiss theorem [10, 103] tells us that

- (a') for every $\epsilon > 0$, there exist $\delta > 0$ and $n_0 < \infty$ such that $Z_{\tilde{S}_L^{(n)}}$ has no (real or complex) zeros within a distance δ from the interval $[-\lambda_L^\times + \epsilon, 0]$ when $n \geq n_0$; and
- (b') there exist (possibly complex) zeros of $Z_{\tilde{S}_L^{(n)}}$ tending to $-\lambda_L^\times$ as $n \rightarrow \infty$.

Since $-\lambda_c(\tilde{S}_L^{(n)})$ is the closest zero to the origin of $Z_{\tilde{S}_L^{(n)}}$, it follows easily from (a') and (b') that $\lambda_L^\times = \lambda_c(\tilde{S}_L)$.

References

- [1] M. Aigner, *Combinatorial Theory*, Grundlehren der mathematischen Wissenschaften #234 (Springer-Verlag, Berlin–New York, 1979).
- [2] N. Alon and J. Spencer, *The Probabilistic Method*, 2nd ed. (Wiley, New York, 2000).
- [3] N. Alon, A parallel algorithmic version of the local lemma, *Random Structures and Algorithms* **2** (1991), 367–378.
- [4] J.L. Arocha, Propiedades del polinomio independiente de un grafo, *Revista Ciencias Matemáticas [Cuba]* **5** (1984), 103–110.
- [5] R. Bamón and J. Bobenrieth, The rational maps $z \mapsto 1 + 1/\omega z^d$ have no Herman rings. *Proc. Amer. Math. Soc.* **127** (1999), 633–636.
- [6] R.J. Baxter, Rogers-Ramanujan identities in the hard hexagon model, *J. Statist. Phys.* **26** (1981), 427–452.
- [7] R.J. Baxter, *Exactly Solved Models in Statistical Mechanics* (Academic Press, London–New York, 1982).
- [8] J. Beck, An algorithmic approach to the Lovász local lemma, I, *Random Structures and Algorithms* **2** (1991), 343–365.
- [9] J. Bellissard and R. Høegh-Krohn, Compactness and the maximal Gibbs state for random Gibbs fields on a lattice, *Commun. Math. Phys.* **84** (1982), 297–327.
- [10] S. Beraha, J. Kahane and N.J. Weiss, Limits of zeroes of recursively defined families of polynomials, in *Studies in Foundations and Combinatorics* (Advances in Mathematics Supplementary Studies, Vol. 1), edited by G.-C. Rota (Academic Press, New York, 1978), pp. 213–232.
- [11] C.A. Berenstein and R. Gay, *Complex Variables: An Introduction* (Springer-Verlag, New York, 1991).
- [12] J. van den Berg and J.E. Steif, Percolation and the hard-core lattice gas model, *Stoch. Proc. Appl.* **49** (1994), 179–197.
- [13] F. Bergeron, G. Labelle and P. Leroux, *Combinatorial Species and Tree-Like Structures* (Cambridge University Press, Cambridge–New York, 1998).
- [14] N.L. Biggs, *Interaction Models*, London Mathematical Society Lecture Note Series #30 (Cambridge University Press, Cambridge, 1977).
- [15] N. Biggs, *Algebraic Graph Theory*, 2nd ed. (Cambridge University Press, Cambridge–New York, 1993).

- [16] P. Billingsley, *Probability and Measure*, 2nd ed. (Wiley, New York, 1986).
- [17] A. Björner, The homology and shellability of matroids and geometric lattices, in N. White (editor), *Matroid Applications* (Encyclopedia of Mathematics and its Applications #40), Chapter 7, pp. 226–283 (Cambridge University Press, Cambridge, 1992).
- [18] J.R. Bobenrieth, Algunos aspectos de la dinámica de las funciones racionales $z \mapsto 1 + 1/\omega z^d$, Ph.D. thesis, Universidad de Chile (June 1997).
- [19] J. Bobenrieth, Hyperbolicity is dense in the real family $z \mapsto 1 + 1/wz^d$ ($w \in \mathbb{R}$), where d is odd, *Scientia, Series A, Mathematical Sciences* (N.S.) **7** (2001), 7–11.
- [20] J. Bobenrieth, Parabolic perturbation in the family $z \mapsto 1 + 1/wz^d$, *Proyecciones* **21** (2002), 1–7.
- [21] J. Bobenrieth, Hyperbolic components of the family $z \mapsto 1 + 1/wz^d$, preprint (2000).
- [22] B. Bollobás, *Modern Graph Theory* (Springer-Verlag, New York, 1998).
- [23] B. Bollobás, *Random Graphs*, 2nd ed. (Cambridge University Press, Cambridge, 2001).
- [24] C. Borgs, *Expansion Methods in Combinatorics*, to be published in the Conference Board of the Mathematical Sciences book series (preprint, 2003).
- [25] A. Bovier and M. Zahradník, A simple inductive approach to the problem of convergence of cluster expansions of polymer models, *J. Statist. Phys.* **100** (2000), 765–778.
- [26] G.R. Brightwell, O. Häggström and P. Winkler, Nonmonotonic behavior in hard-core and Widom-Rowlinson models, *J. Statist. Phys.* **94** (1999), 415–435.
- [27] J.I. Brown, K. Dilcher and R.J. Nowakowski, Roots of independence polynomials of well covered graphs, *J. Algebraic Combin.* **11** (2000), 197–210.
- [28] J.I. Brown and R.J. Nowakowski, Bounding the roots of independence polynomials, *Ars Combin.* **58** (2001), 113–120.
- [29] D.C. Brydges, A short course on cluster expansions, in K. Osterwalder and R. Stora (editors), *Phénomènes Critiques, Systèmes Aléatoires, Théories de Jauge / Critical Phenomena, Random Systems, Gauge Theories* (Les Houches summer school, Session XLIII, 1984), pp. 129–183 (Elsevier/North-Holland, Amsterdam, 1986).
- [30] D.C. Brydges and T. Kennedy, Mayer expansions and the Hamilton-Jacobi equation, *J. Statist. Phys.* **48** (1987), 19–49.

- [31] D.C. Brydges and Ph.A. Martin, Coulomb systems at low density: A review, *J. Statist. Phys.* **96** (1999), 1163–1330, cond-mat/9904122 at arXiv.org.
- [32] D.C. Brydges and J.D. Wright, Mayer expansions and the Hamilton-Jacobi equation. II. Fermions, dimensional reduction formulas. *J. Statist. Phys.* **51** (1988), 435–456; erratum **97** (1999), 1027.
- [33] C. Cammarota, Decay of correlations for infinite range interactions in unbounded spin systems, *Commun. Math. Phys.* **85** (1982), 517–528.
- [34] Y.-B. Choe, J.G. Oxley, A.D. Sokal and D.G. Wagner, Homogeneous multivariate polynomials with the half-plane property, *Adv. Appl. Math.* **32** (2004), 88–187, math.CO/0202034 at arXiv.org.
- [35] M. Chudnovsky and P. Seymour, The roots of the stable set polynomial of a claw-free graph, preprint (February 2004).
- [36] R.L. Dobrushin, The problem of uniqueness of a Gibbs random field and the problem of phase transition, *Funct. Anal. Appl.* **2** (1968), 302–312.
- [37] R.L. Dobrushin, Estimates of semi-invariants for the Ising model at low temperatures, in *Topics in Statistical and Theoretical Physics*, American Mathematical Society Translations, Ser. 2, **177**, 59–81 (1996).
- [38] R.L. Dobrushin, Perturbation methods of the theory of Gibbsian fields, in P. Bernard (editor), *Lectures on Probability Theory and Statistics*, Ecole d’Eté de Probabilités de Saint-Flour XXIV – 1994, Lecture Notes in Mathematics #1648, pp. 1–66. Springer–Verlag, Berlin, 1996.
- [39] P. Erdős and L. Lovász, Problems and results on 3-chromatic hypergraphs and some related questions, in *Infinite and finite sets*, Vol. II, Colloq. Math. Soc. Janos Bolyai, Vol. 10, North-Holland, Amsterdam, 1975, pp. 609–627.
- [40] P. Erdős and J. Spencer, Lopsided Lovász local lemma and Latin transversals, *Discrete Appl. Math.* **30** (1991), 151–154.
- [41] D.C. Fisher and J.M. Nonis, Bounds on the “growth factor” of a graph, *Congr. Numer.* **77** (1990), 113–119.
- [42] D.C. Fisher and A.E. Solow, Dependence polynomials, *Discrete Math.* **82** (1990), 251–258.
- [43] D. Galvin and J. Kahn, On phase transition in the hard-core model on \mathbf{Z}^d , *Combin. Probab. Comput.* **13** (2004), 137–164.
- [44] M.R. Garey and D.S. Johnson, *Computers and Intractability: A Guide to the Theory of NP-Completeness* (Freeman, San Francisco, 1979).

- [45] I.M. Gessel and B.E. Sagan, The Tutte polynomial of a graph, depth-first search, and simplicial complex partitions, *Electronic J. Combinatorics* **3**, no. 2, #R9 (1996).
- [46] J. Ginibre, On some recent work of Dobrushin, in *Systèmes à un nombre infini de degrés de liberté*, Colloques Internationaux du Centre National de la Recherche Scientifique, no. 181, Éditions du Centre National de la Recherche Scientifique, Paris, 1970, pp. 163–175.
- [47] J. Glimm and A. Jaffe, *Quantum Physics: A Functional Integral Point of View*, 2nd ed. (Springer-Verlag, New York, 1987).
- [48] J. Groeneveld, Two theorems on classical many-particle systems, *Phys. Lett.* **3** (1962), 50–51.
- [49] R.C. Gunning, *Introduction to Holomorphic Functions of Several Variables*, Vol. I: Function Theory (Wadsworth & Brooks/Cole, Pacific Grove, California, 1990).
- [50] W. Guo and H.W.J. Blöte, Finite-size analysis of the hard-square lattice gas, *Phys. Rev. E* **66** (2002), 046140.
- [51] I. Gutman, An identity for the independence polynomials of trees, *Publ. Inst. Math. (Beograd) (N.S.)* **50(64)** (1991), 19–23.
- [52] I. Gutman, Independent vertex sets in some compound graphs, *Publ. Inst. Math. (Beograd) (N.S.)* **52(66)** (1992), 5–9.
- [53] I. Gutman, Some analytical properties of the independence and matching polynomials, *Match* **28** (1992), 139–150.
- [54] A.J. Guttmann, Comment on ‘The exact location of partition function zeros, a new method for statistical mechanics’, *J. Phys. A: Math. Gen.* **20** (1987), 511–512.
- [55] O. Häggström, Ergodicity of the hard-core model on \mathbf{Z}^2 with parity-dependent activities, *Ark. Mat.* **35** (1997), 171–184.
- [56] Y.O. Hamidoune, On the numbers of independent k -sets in a claw free graph, *J. Combin. Theory B* **50** (1990), 241–244.
- [57] O.J. Heilmann, Use of reflection as symmetry operation in connection with Peierls argument, *Commun. Math. Phys.* **36** (1974), 91–114.
- [58] O.J. Heilmann and E.H. Lieb, Theory of monomer-dimer systems, *Commun. Math. Phys.* **25** (1972), 190–232.
- [59] O.J. Heilmann and E. Præstgaard, Phase transition of hard hexagons on a triangular lattice, *J. Stat. Phys.* **9** (1973), 23–44.

- [60] O.J. Heilmann and E. Præstgaard, Phase transition in a lattice gas with third nearest neighbour exclusion on a square lattice, *J. Phys. A: Math. Gen.* **7** (1974), 1913–1917.
- [61] J.R. Heringa, H.W.J. Blöte and E. Luijten, High-dimensional lattice gases, *J. Phys. A: Math. Gen.* **33** (2000), 2929–2941.
- [62] E. Hille, *Analytic Function Theory*, 2nd ed. (Chelsea, New York, 1973).
- [63] C. Hoede and X.-L. Li, Clique polynomials and independent set polynomials of graphs, *Discrete Math.* **125** (1994), 219–228.
- [64] D.G.C. Horrocks, The numbers of dependent k -sets in a graph are log concave, *J. Combin. Theory B* **84** (2002), 180–185.
- [65] T.W. Hungerford, *Algebra* (Springer-Verlag, New York, 1974, fifth printing 1989).
- [66] A. Iserles, On the generalized pantograph functional-differential equation, *European J. Appl. Math.* **4** (1993), 1–38.
- [67] R.B. Israel, *Convexity in the Theory of Lattice Gases* (Princeton University Press, Princeton NJ, 1979).
- [68] S. Janson, New versions of Suen’s correlation inequality, *Random Structures and Algorithms* **13** (1998), 467–483.
- [69] J. Jonasson and J.E. Steif, Amenability and phase transition in the Ising model, *J. Theoret. Probab.* **12** (1999), 549–559.
- [70] J. Kahn, An entropy approach to the hard-core model on bipartite graphs, *Combin. Probab. Comput.* **10** (2001), 219–237.
- [71] J. Kahn, Entropy, independent sets and antichains: A new approach to Dedekind’s problem, *Proc. Amer. Math. Soc.* **130** (2002), 371–378.
- [72] R. Kotecký and D. Preiss, Cluster expansion for abstract polymer models, *Commun. Math. Phys.* **103** (1986), 491–498.
- [73] S.G. Krantz, *Function Theory of Several Complex Variables*, 2nd ed. (Wadsworth & Brooks/Cole, Pacific Grove, California, 1992).
- [74] H. Künsch, Almost sure entropy and the variational principle for random fields with unbounded state space, *Z. Wahrsch. verw. Gebiete* **58** (1981), 69–85.
- [75] J.L. Lebowitz and E. Presutti, Statistical mechanics of systems of unbounded spins, *Commun. Math. Phys.* **50** (1976), 195–218; erratum **78** (1980/81), 151.
- [76] B.Ja. Levin, *Distribution of Zeros of Entire Functions* (American Mathematical Society, Providence, 1964).

- [77] V.E. Levit and E. Mandrescu, On the roots of independence polynomials of almost all very well-covered graphs, preprint (2003), math.CO/0305227 at arXiv.org.
- [78] L. Lovász and M.D. Plummer, *Matching Theory*, North-Holland Mathematics Studies #121 / Annals of Discrete Mathematics #29 (North-Holland, Amsterdam-New York / Akadémiai Kiadó, Budapest, 1986).
- [79] W.S. Massey, *Algebraic Topology: An Introduction*, 4th ed. (Springer-Verlag, New York, 1990).
- [80] C. Merino, A. de Mier and M. Noy, Irreducibility of the Tutte polynomial of a connected matroid, *J. Combin. Theory B* **83** (2001), 298–304.
- [81] J. Milnor, On rational maps with two critical points, *Experiment. Math.* **9** (2000), 481–522.
- [82] S. Miracle-Solé, On the convergence of cluster expansions, *Physica A* **279** (2000), 244–249.
- [83] M. Molloy and B. Reed, *Graph Colouring and the Probabilistic Method* (Springer-Verlag, Berlin–New York, 2002).
- [84] P. Montel, *Leçons sur les familles normales de fonctions analytiques et leurs applications* (Gauthier-Villars, Paris, 1927). Reprinted by Chelsea Publishing Co., New York, 1974.
- [85] G.R. Morris, A. Feldstein and E.W. Bowen, The Phragmén–Lindelöf principle and a class of functional differential equations, In L. Weiss (editor), *Ordinary Differential Equations: 1971 NRL-MRC Conference* (Academic Press, New York, 1972), pp. 513–540.
- [86] J.G. Oxley, *Matroid Theory* (Oxford University Press, New York, 1992).
- [87] O. Penrose, Convergence of fugacity expansions for fluids and lattice gases, *J. Math. Phys.* **4** (1963), 1312–1320.
- [88] O. Penrose, Convergence of fugacity expansions for classical systems, in T.A. Bak (editor), *Statistical Mechanics: Foundations and Applications*, pp. 101–109 (Benjamin, New York–Amsterdam, 1967).
- [89] A. Procacci, B.N.B. de Lima and B. Scoppola, A remark on high temperature polymer expansion for lattice systems with infinite range pair interactions, *Lett. Math. Phys.* **45** (1998), 303–322.
- [90] A. Procacci and B. Scoppola, Polymer gas approach to N -body lattice systems, *J. Statist. Phys.* **96** (1999), 49–68.

- [91] A. Procacci, B. Scoppola and V. Gerasimov, Potts model on infinite graphs and the limit of chromatic polynomials, *Commun. Math. Phys.* **235** (2003), 215–231.
- [92] R.M. Range, *Holomorphic Functions and Integral Representations in Several Complex Variables* (Springer-Verlag, New York, 1986).
- [93] D. Ruelle, Probability estimates for continuous spin systems, *Commun. Math. Phys.* **50** (1976), 189–194.
- [94] L.K. Runnels, Phase transition of a Bethe lattice gas of hard molecules. *J. Math. Phys.* **10** (1967), 2081–2087.
- [95] L.K. Runnels, Phase transitions of hard sphere lattice gases, *Commun. Math. Phys.* **40** (1975), 37–48.
- [96] J.L. Schiff, *Normal Families* (Springer-Verlag, New York, 1993).
- [97] E.G. Seiler, *Gauge Theories as a Problem of Constructive Quantum Field Theory and Statistical Mechanics*, Lecture Notes in Physics #159 (Springer-Verlag, Berlin–New York, 1982).
- [98] J. B. Shearer, On a problem of Spencer, *Combinatorica* **5** (1985), 241–245.
- [99] B. Simon, *The $P(\varphi)_2$ Euclidean (Quantum) Field Theory* (Princeton University Press, Princeton NJ, 1974).
- [100] B. Simon, *The Statistical Mechanics of Lattice Gases* (Princeton University Press, Princeton NJ, 1993).
- [101] A.D. Sokal, Bounds on the complex zeros of (di)chromatic polynomials and Potts-model partition functions, *Combin. Probab. Comput.* **10** (2001), 41–77, [cond-mat/9904146](https://arxiv.org/abs/cond-mat/9904146) at arXiv.org.
- [102] A.D. Sokal, A personal list of unsolved problems concerning lattice gases and antiferromagnetic Potts models, *Markov Process. Related Fields* **7** (2001), 21–38, [cond-mat/0004231](https://arxiv.org/abs/cond-mat/0004231) at arXiv.org.
- [103] A.D. Sokal, Chromatic roots are dense in the whole complex plane, *Combin. Probab. Comput.* **13** (2004), 221–261, [cond-mat/0012369](https://arxiv.org/abs/cond-mat/0012369) at arXiv.org.
- [104] A.D. Sokal, Tree limit of Mayer expansion, in preparation.
- [105] J. Spencer, Ramsey’s Theorem — a new lower bound, *J. Comb. Theory A* **18** (1975), 108–115.
- [106] J. Spencer, Asymptotic lower bounds for Ramsey functions, *Discrete Math.* **20** (1977/78), 69–76.
- [107] R.P. Stanley, *Enumerative Combinatorics*, vol. 1 (Wadsworth & Brooks/Cole, Monterey, California, 1986). Reprinted by Cambridge University Press, 1999.

- [108] R.P. Stanley, *Enumerative Combinatorics*, vol. 2 (Cambridge University Press, Cambridge–New York, 1999).
- [109] W.C. Suen, A correlation inequality and a Poisson limit theorem for nonoverlapping balanced subgraphs of a random graph, *Random Structures and Algorithms* **1** (1990), 231–242.
- [110] S. Todo, Transfer-matrix study of negative-fugacity singularity of hard-core lattice gas, *Int. J. Mod. Phys C* **10** (1999), 517–529, cond-mat/9703176 at arXiv.org.
- [111] S. Todo, private communication (August 2003).
- [112] G.E. Uhlenbeck and G.W. Ford, The theory of linear graphs with applications to the theory of the virial development of the properties of gases, in J. de Boer and G.E. Uhlenbeck (editors), *Studies in Statistical Mechanics*, vol. I, pp. 119–211 (North-Holland, Amsterdam, 1962).
- [113] A.C.D. van Enter, R. Fernández and A.D. Sokal, Regularity properties and pathologies of position-space renormalization-group transformations: Scope and limitations of Gibbsian theory, *J. Statist. Phys.* **72** (1993), 879–1167.
- [114] V.S. Vladimirov, *Methods of the Theory of Functions of Many Complex Variables* (MIT Press, Cambridge, Mass., 1966).
- [115] H.S. Wilf, *generatingfunctionology*, 2nd ed. (Academic Press, San Diego–London, 1994).
- [116] D. Williams, *Probability with Martingales* (Cambridge University Press, Cambridge, 1991).
- [117] G.M. Ziegler, *Lectures on Polytopes* (Springer-Verlag, New York, 1995).